



planetmath.org

Math for the people, by the people.

subanalytic set

Canonical name	SubanalyticSet
Date of creation	2013-03-22 16:46:16
Last modified on	2013-03-22 16:46:16
Owner	jirka (4157)
Last modified by	jirka (4157)
Numerical id	6
Author	jirka (4157)
Entry type	Definition
Classification	msc 32B20
Classification	msc 14P15
Related topic	TarskiSeidenbergTheorem
Related topic	SemialgebraicSet
Defines	subanalytic
Defines	semianalytic set
Defines	semianalytic
Defines	semianalytic function
Defines	subanalytic function
Defines	semianalytic mapping
Defines	subanalytic mapping
Defines	dimension of a subanalytic set

Let  $U \subset \mathbb{R}^n$ . Suppose  $\mathcal{A}(U)$  is any ring of real valued functions on  $U$ . Define  $\mathcal{S}(\mathcal{A}(U))$  to be the smallest set of subsets of  $U$ , which contain the sets  $\{x \in U \mid f(x) > 0\}$  for all  $f \in \mathcal{A}(U)$ , and is closed under finite union, finite intersection and complement.

**Definition.** A set  $V \subset \mathbb{R}^n$  is *semianalytic* if and only if for each  $x \in \mathbb{R}^n$ , there exists a neighbourhood  $U$  of  $x$ , such that  $V \cap U \in \mathcal{S}(\mathcal{O}(U))$ , where  $\mathcal{O}(U)$  denotes the real-analytic real valued functions.

Unlike for semialgebraic sets, there is no Tarski-Seidenberg theorem for semianalytic sets, and projections of semianalytic sets are in general not semianalytic.

**Definition.** We say  $V \subset \mathbb{R}^n$  is a *subanalytic* set if for each  $x \in \mathbb{R}^n$ , there exists a relatively compact semianalytic set  $X \subset \mathbb{R}^{n+m}$  and a neighbourhood  $U$  of  $x$ , such that  $V \cap U$  is the projection of  $X$  onto the first  $n$  coordinates.

In particular all semianalytic sets are subanalytic. On an open dense set subanalytic sets are submanifolds and hence we can define dimension. Hence at a point  $p$ , where a set  $A$  is a submanifold, the dimension  $\dim_p A$  is the dimension of the submanifold. The *dimension* of the subanalytic set is the maximum  $\dim_p A$  for all  $p$  where  $A$  is a submanifold. Semianalytic sets are contained in a real-analytic subvariety of the same dimension. However, subanalytic sets are not in general contained in any subvariety of the same dimension. We do have however the following.

**Theorem.** *A subanalytic set  $A$  can be written as a locally finite union of submanifolds.*

The set of subanalytic sets is still not completely closed under projections however. Note that a real-analytic subvariety that is not relatively compact can have a projection which is not a locally finite union of submanifolds, and hence is not subanalytic.

**Definition.** Let  $U \subset \mathbb{R}^n$ . A mapping  $f: U \rightarrow \mathbb{R}^m$  is said to be subanalytic (resp. semianalytic) if the graph of  $f$  (i.e. the set  $\{(x, y) \in U \times \mathbb{R}^m \mid x, y = f(x)\}$ ) is subanalytic (resp. semianalytic)

## References

- [1] Edward Bierstone and Pierre D. Milman, *Semianalytic and subanalytic sets*, Inst. Hautes Études Sci. Publ. Math. (1988), no. 67, 5–42. <http://www.ams.org/mathscinet-getitem?mr=89k:32011MR89k:32011>