



planetmath.org

Math for the people, by the people.

coherent analytic sheaf

Canonical name	CoherentAnalyticSheaf
Date of creation	2013-03-22 17:39:05
Last modified on	2013-03-22 17:39:05
Owner	jirka (4157)
Last modified by	jirka (4157)
Numerical id	4
Author	jirka (4157)
Entry type	Definition
Classification	msc 32C35
Defines	locally finitely generated sheaf
Defines	sheaf of relations

Let M be a complex manifold and \mathcal{F} be an analytic sheaf. For $z \in M$, denote by \mathcal{F}_z the stalk of \mathcal{F} at z . By \mathcal{O} denote the sheaf of germs of analytic functions. For a section f and a point $z \in M$ denote by f_z the germ of f at z .

\mathcal{F} is said to be *locally finitely generated* if for every $z \in M$, there is a neighbourhood U of z , a finite number of sections $f_1, \dots, f_k \in \Gamma(U, \mathcal{F})$ such that for each $w \in U$, \mathcal{F}_w is a finitely generated module (as an \mathcal{O}_w -module).

Let U be a neighbourhood in M and Suppose that f_1, \dots, f_k are sections in $\Gamma(U, \mathcal{F})$. Let $\mathcal{R}(f_1, \dots, f_k)$ be the subsheaf of \mathcal{O}^k over U consisting of the germs

$$\{(g_1, \dots, g_k) \in \mathcal{O}_z^k \mid \sum_{j=1}^k g_j(f_j)_z = 0\}.$$

$\mathcal{R}(f_1, \dots, f_k)$ is called the *sheaf of relations*.

Definition. \mathcal{F} is called a *coherent analytic sheaf* if \mathcal{F} is locally finitely generated and if for every open subset $U \subset M$, and $f_1, \dots, f_k \in \Gamma(U, \mathcal{F})$, the sheaf $\mathcal{R}(f_1, \dots, f_k)$ is locally finitely generated.

References

- [1] Lars Hörmander. , North-Holland Publishing Company, New York, New York, 1973.
- [2] Steven G. Krantz. , AMS Chelsea Publishing, Providence, Rhode Island, 1992.