



Math for the people, by the people.

# Kontinuitätssatz

Canonical name	Kontinuitatssatz
Date of creation	2013-03-22 15:49:15
Last modified on	2013-03-22 15:49:15
Owner	jirka (4157)
Last modified by	jirka (4157)
Numerical id	5
Author	jirka (4157)
Entry type	Theorem
Classification	msc 32T05
Synonym	Hartogs Kontinuitätssatz
Synonym	Kontinuitatssatz

**Theorem.**  $G \subset \mathbb{C}^n$  is pseudoconvex if and only if for any family of closed analytic discs  $\{d_\alpha\}_{\alpha \in I}$  in  $G$  with  $\cup_{\alpha \in I} \partial d_\alpha$  being a relatively compact set in  $G$  then  $\cup_{\alpha \in I} d_\alpha$  is also a relatively compact set in  $G$ .

This is the analogue of one of the definitions of a convex set. Just replace pseudoconvex with convex and closed analytic discs with closed line segments.

## References

- [1] M. Salah Baouendi, Peter Ebenfelt, Linda Preiss Rothschild. , Princeton University Press, Princeton, New Jersey, 1999.
- [2] Steven G. Krantz. , AMS Chelsea Publishing, Providence, Rhode Island, 1992.