



planetmath.org

Math for the people, by the people.

CR submanifold

Canonical name	CRSubmanifold
Date of creation	2013-03-22 14:49:04
Last modified on	2013-03-22 14:49:04
Owner	jirka (4157)
Last modified by	jirka (4157)
Numerical id	9
Author	jirka (4157)
Entry type	Definition
Classification	msc 32V05
Synonym	CR manifold
Synonym	Cauchy-Riemann submanifold
Related topic	GenericManifold
Related topic	TotallyRealSubmanifold
Related topic	TangentialCauchyRiemannComplexOfCinftySmoothForms
Related topic	ACRcomplex
Defines	CR bundle
Defines	CR vector field
Defines	complex tangent space
Defines	complex bundle
Defines	space of antiholomorphic vectors
Defines	antiholomorphic vector
Defines	CR dimension

Suppose that $M \subset \mathbb{C}^N$ is a real submanifold of real dimension n . Take $p \in M$, then let $T_p(\mathbb{C}^N)$ be the tangent vectors of \mathbb{C}^N at the point p . If we identify \mathbb{C}^N with \mathbb{R}^{2N} by $z_j = x_j + iy_j$, we can take the following vectors as our basis

$$\left. \frac{\partial}{\partial x_1} \right|_p, \left. \frac{\partial}{\partial y_1} \right|_p, \dots, \left. \frac{\partial}{\partial x_N} \right|_p, \left. \frac{\partial}{\partial y_N} \right|_p.$$

We define a real linear mapping $J: T_p(\mathbb{C}^N) \rightarrow T_p(\mathbb{C}^N)$ such that for any $1 \leq j \leq N$ we have

$$J \left(\left. \frac{\partial}{\partial x_1} \right|_p \right) = \left. \frac{\partial}{\partial y_1} \right|_p \quad \text{and} \quad J \left(\left. \frac{\partial}{\partial y_1} \right|_p \right) = - \left. \frac{\partial}{\partial x_1} \right|_p.$$

Where J is referred to as the complex structure on $T_p(\mathbb{C}^N)$. Note that $J^2 = -I$, that is applying J twice we just negate the vector.

Let $T_p(M)$ be the tangent space of M at the point p (that is, those vectors of $T_p(\mathbb{C}^N)$ which are tangent to M).

Definition. The subspace $T_p^c(M)$ defined as

$$T_p^c(M) := \{X \in T_p(M) \mid J(X) \in T_p(M)\}$$

is called the *complex tangent space* of M at the point p , and if the dimension of $T_p(M)$ is constant for all $p \in M$ then the corresponding vector bundle $T^c(M) := \bigcup_{p \in M} T_p^c(M)$ is called the *complex bundle* of M .

Do note that the complex tangent space is a real (not complex) vector space, despite its rather unfortunate name.

Let $\mathbb{C}T_p(M)$ and $\mathbb{C}T_p(\mathbb{C}^N)$ be the complexified vector spaces, by just allowing the coefficients of the vectors to be complex numbers. That is for $X = \sum a_j \left. \frac{\partial}{\partial x_1} \right|_p + b_j \left. \frac{\partial}{\partial y_1} \right|_p$ we allow a_j and b_j to be complex numbers. Next we can extend the mapping J to be \mathbb{C} -linear on these new vector spaces and still get that $J^2 = -I$ as before. We notice that the operator J has two eigenvalues, i and $-i$.

Definition. Let \mathcal{V}_p be the eigenspace of $\mathbb{C}T_p(M)$ corresponding to the eigenvalue $-i$. That is

$$\mathcal{V}_p := \{X \in \mathbb{C}T_p(M) \mid J(X) = -iX\}.$$

If the dimension of \mathcal{V}_p is constant for all $p \in M$, then we get a corresponding vector bundle \mathcal{V} which we call the *CR bundle* of M . A smooth section of the CR bundle is then called a *CR vector field*.

Definition. The submanifold M is called a *CR submanifold* (or just *CR manifold*) if the dimension of \mathcal{V}_p is constant for all $p \in M$. The complex dimension of \mathcal{V}_p will then be called the *CR dimension* of M .

An example of a CR submanifold is for example a hyperplane defined by $\text{Im } z_N = 0$ where the CR dimension is $N - 1$. Another less trivial example is the Lewy hypersurface.

Note that sometimes \mathcal{V}_p is written as $T_p^{0,1}(M)$ and referred to as the *space of antiholomorphic vectors*, where an *antiholomorphic vector* is a tangent vector which can be written in terms of the basis

$$\left. \frac{\partial}{\partial \bar{z}_j} \right|_p := \frac{1}{2} \left(\left. \frac{\partial}{\partial x_j} \right|_p + i \left. \frac{\partial}{\partial y_j} \right|_p \right).$$

The CR in the name refers to Cauchy-Riemann and that is because the vector space \mathcal{V}_p corresponds to differentiating with respect to \bar{z}_j .

References

- [1] M. Salah Baouendi, Peter Ebenfelt, Linda Preiss Rothschild. , Princeton University Press, Princeton, New Jersey, 1999.
- [2] Albert Boggess. , CRC, 1991.