



planetmath.org

Math for the people, by the people.

pseudoconvex

Canonical name	Pseudoconvex
Date of creation	2013-03-22 14:30:58
Last modified on	2013-03-22 14:30:58
Owner	jirka (4157)
Last modified by	jirka (4157)
Numerical id	5
Author	jirka (4157)
Entry type	Definition
Classification	msc 32T15
Classification	msc 32T05
Synonym	Hartogs pseudoconvex
Related topic	LeviPseudoconvex
Related topic	SolutionOfTheLeviProblem
Related topic	ExhaustionFunction

**Definition.** Let  $G \subset \mathbb{C}^n$  be a domain (open connected subset). We say  $G$  is *pseudoconvex* (or *Hartogs pseudoconvex*) if there exists a continuous plurisubharmonic function  $\varphi$  on  $G$  such that the sets  $\{z \in G \mid \varphi(z) < x\}$  are relatively compact subsets of  $G$  for all  $x \in \mathbb{R}$ . That is we say that  $G$  has a continuous plurisubharmonic exhaustion function.

When  $G$  has a  $C^2$  (twice continuously differentiable) boundary then this notion is the same as <http://planetmath.org/LeviPseudoconvex> Levi pseudoconvexity, which is easier to work with if you have such nice boundaries. If you don't have nice boundaries then the following approximation result can come in useful.

**Proposition.** *If  $G \subset \mathbb{C}^n$  is pseudoconvex then there exist bounded, strongly Levi pseudoconvex domains  $G_k \subset G$  with  $C^\infty$  (smooth) boundary which are relatively compact in  $G$ , such that  $G = \bigcup_{k=1}^\infty G_k$ .*

This is because once we have a  $\varphi$  as in the definition we can actually find a  $C^\infty$  exhaustion function.

The reason for the definition of pseudoconvexity is that it classifies domains of holomorphy. One thing to note then is that every open domain in one complex dimension (in the complex plane  $\mathbb{C}$ ) is then pseudoconvex.

## References

- [1] Steven G. Krantz. , AMS Chelsea Publishing, Providence, Rhode Island, 1992.