



Math for the people, by the people.

## biholomorphisms of strongly pseudoconvex domains extend to the boundary

Canonical name	BiholomorphismsOfStronglyPseudoconvexDomainsExtendToTheBoundary
Date of creation	2013-03-22 16:44:30
Last modified on	2013-03-22 16:44:30
Owner	jirka (4157)
Last modified by	jirka (4157)
Numerical id	4
Author	jirka (4157)
Entry type	Theorem
Classification	msc 32T15
Related topic	LeviPseudoconvex

It is a basic question in complex analysis to ask when does a biholomorphic mapping of two domains extend to the boundary. The following is a celebrated theorem of Fefferman for strongly pseudoconvex domains.

**Theorem** (Fefferman). *Let  $U, V \subset \mathbb{C}^n$  ( $n \geq 2$ ) be two strongly pseudoconvex domains with smooth ( $C^\infty$ ) boundaries (the boundaries are smooth submanifolds). Let  $f: U \rightarrow V$  be a biholomorphism. Then  $f$  extends to a smooth diffeomorphism of  $\bar{U}$  to  $\bar{V}$ .*

## References

- [1] Fefferman, Charles. . Invent. Math. **26** (1974), 1–65.
- [2] Steven G. Krantz. , AMS Chelsea Publishing, Providence, Rhode Island, 1992.