



planetmath.org

Math for the people, by the people.

Puiseux parametrization

Canonical name	PuiseuxParametrization
Date of creation	2013-03-22 15:20:32
Last modified on	2013-03-22 15:20:32
Owner	jirka (4157)
Last modified by	jirka (4157)
Numerical id	6
Author	jirka (4157)
Entry type	Theorem
Classification	msc 32B10
Synonym	Puiseux series parametrization
Synonym	Puiseux normalization
Synonym	Puiseux series normalization
Synonym	Puiseux parameterization
Synonym	Puiseux series parameterization
Related topic	PuiseuxSeries

Theorem. *Suppose that $V \subset U \subset \mathbb{C}^2$ is an irreducible complex analytic subset of (complex) dimension 1 where U is a domain. Suppose that $0 \in V$. Then there exists an analytic (holomorphic) map $f: \mathbb{D} \rightarrow V$, where \mathbb{D} is the unit disc, such that $f(0) = 0$ and $f(\mathbb{D}) = N$ where $N \subset V$ is a neighbourhood of 0 in V , f is one to one, and further $f|_{\mathbb{D} \setminus \{0\}}$ is a biholomorphism onto $N \setminus \{0\}$. In fact there exist suitable local coordinates (z, w) in \mathbb{C}^2 such that f is then given by $\xi \mapsto (z, w)$ where $z = \xi^k$, $w = \sum_{n=m}^{\infty} a_n \xi^n$ where $m > k$.*

This is sometimes written as

$$w = \sum_{n=m}^{\infty} a_n z^{n/k}$$

and hence the name *Puiseux series parametrization*. If you do however write it like this, it must be properly interpreted, as the Puiseux series is in general not single valued.

A similar result for arbitrary complex analytic sets with singularities of codimension 1 in higher dimensional spaces under further conditions on the singular set was obtained by Stutz, see Chirka [?] page 98.

References

- [1] E. M. Chirka. . Kluwer Academic Publishers, Dordrecht, The Netherlands, 1989.
- [2] Alexandru Dimca. . Vieweg, Braunschweig, Germany, 1987.