



WELCOME TO THE NARCOLEPSY TRIVIA GAME!

DO YOU HAVE A KNACK FOR NARCOLEPSY KNOWLEDGE?
PROVE IT!

[LEADERBOARD](#)[NEW GAME](#)

HEART DISEASE
DIABETES
HYPERTENSION

INVITE YOUR PEERS TO PLAY

FACEBOOK

LINKEDIN

TWITTER

Jazz Pharmaceuticals

REFERENCES

PRIVACY POLICY

?

©2020 Jazz Pharmaceuticals plc or its subsidiaries US-SLE-2000396 Rev0820

WELCOME TO THE NARCOLEPSY TRIVIA GAME!

DO YOU HAVE A KNACK FOR NARCOLEPSY KNOWLEDGE?

HOW TO PLAY

This game will test your narcolepsy knowledge. Let's see how you compare to your peers!

You will be asked 12 questions related to narcolepsy. Each has a maximum point value of 100, 200, or 300. You'll have a few seconds to read each question before the countdown begins. The faster you answer the question, the more points you can gain!

If you do well enough, you will have a chance to answer a bonus question. You can bet some or all of your points—and maybe double your score. Do well enough, fast enough, and you might make it onto the leaderboard.

Good luck!

INVITE YOUR PEERS TO PLAY



©2020 Jazz Pharmaceuticals plc or its subsidiaries US-SLE-2000396 Rev0820

REFERENCES

PRIVACY POLICY





HOW STRONG ARE YOUR SKILLS?

LEADERBOARD

3200 | A SMI
Medical School Name

3200 | A SMI
Medical School Name

3200 | A SMI
Medical School Name

3200 | A SMI
Medical School Name

3200 | A SMI
Medical School Name

3200 | A SMI
Medical School Name

3200 | A SMI
Medical School Name

3200 | A SMI
Medical School Name

3200 | A SMI
Medical School Name

3200 | A SMI
Medical School Name

INVITE YOUR PEERS TO PLAY



©2020 Jazz Pharmaceuticals plc or its subsidiaries US-SLE-2000396 Rev0820

REFERENCES

PRIVACY POLICY





REFERENCES

1. Narcolepsy Fact Sheet. National Institute of Neurological Disorders and Stroke website. <https://www.ninds.nih.gov/Disorders/Patient-Caregiver-Education/Fact-Sheets/Narcolepsy-Fact-Sheet>. Updated March 16, 2020. Accessed March 26, 2020.
2. American Academy of Sleep Medicine. Central disorders of hypersomnolence. In: *The International Classification of Sleep Disorders – Third Edition (ICSD-3) Online Version*. Darien, IL: American Academy of Sleep Medicine; 2014.
3. Carter LP, Acebo C, Kim A. Patients' journeys to a narcolepsy diagnosis: a physician survey and retrospective chart review. *Postgrad Med*. 2014;126(3):216-224.
4. Black J, Reaven NL, Funk SE, et al. Medical comorbidity in narcolepsy: findings from the Burden of Narcolepsy Disease (BOND) study. *Sleep Med*. 2017;33:13-18.
5. Ohayon MM. Narcolepsy is complicated by high medical and psychiatric comorbidities: a comparison with the general population. *Sleep Med*. 2013;14(6):488-492.
6. Cohen A, Mandrekar J, St Louis EK, Silber MH, Kotagal S. Comorbidities in a community sample of narcolepsy. *Sleep Med*. 2018;43:14-18.
7. Benjamin EJ, Muntner P, Alonso A, et al; American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2019 update: a report from the American Heart Association. *Circulation*. 2019;139(10):e56-e528.
8. Lloyd-Jones DM, Hong Y, Labarthe D, et al. Defining and setting national goals for cardiovascular health promotion and disease reduction: the American Heart Association's strategic Impact Goal through 2020 and beyond. *Circulation*. 2010;121(4):586-613.
9. Quader ZS, Zhao L, Gillespie C, et al. Sodium intake among persons aged ≥ 2 years—United States, 2013-2014. *MMWR Morb Mortal Wkly Rep*. 2017;66(12):324-328.
10. Perrin G, Korb-Savoldelli V, Karras A, Danchin N, Durieux P, Sabatier B. Cardiovascular risk associated with high sodium-containing drugs: a systematic review. *PLoS One*. 2017;12(7):e0180634.
11. Rodriguez CJ, Bibbins-Domingo K, Jin Z, Daviglius ML, Goff DC Jr, Jacobs DR Jr. Association of sodium and potassium intake with left ventricular mass: coronary artery risk development in young adults. *Hypertension*. 2011;58(3):410-416.
12. Gardener H, Rundek T, Wright CB, Elkind MS, Sacco RL. Dietary sodium and risk of stroke in the Northern Manhattan study. *Stroke*. 2012;43(5):1200-1205.
13. Strazzullo P, D'Elia L, Kandala NB, Cappuccio FP. Salt intake, stroke, and cardiovascular disease: meta-analysis of prospective studies. *BMJ*. 2009;339:b4567.
14. Institute of Medicine. *Sodium Intake in Populations: Assessment of Evidence*. Washington, DC: National Academies Press; 2013.
15. Johns M, Hocking B. Excessive daytime sleepiness and sleep habits of Australian workers. *Sleep*. 1997;20(10):844-849.
16. Green PM, Stillman MJ. Narcolepsy: signs, symptoms, differential diagnosis, and management. *Arch Fam Med*. 1998;7(5):472-478.
17. Overeem S, van Nues SJ, van der Zande WL, Donjacour CE, van Mierlo P, Lammers GJ. The clinical features of cataplexy: a questionnaire study in narcolepsy patients with and without hypocretin-1 deficiency. *Sleep Med*. 2011;12(1):12-18.



BETES

YOUR SCORE : 0

HOW STRONG ARE YOUR SKILLS?

NARCOLEPSY KNOWLEDGE	NARCOLEPSY AND COMORBIDITIES	OVERALL HEALTH	DIAGNOSING NARCOLEPSY
100	100	100	100
200	200	200	200
300	300	300	300
			BONUS QUESTION

TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 0

According to the NINDS and AASM,
onset of narcolepsy symptoms typically occurs
during which age range?

- A

>50 years old
- C

25-35 years old
- B

35-50 years old
- D

10-25 years old

YOUR SCORE : 100

Correct, the answer is D

According to the NINDS and AASM,
onset of narcolepsy symptoms typically occurs
during the age range of 10-25 years old.^{1,2}

CONTINUE

YOUR SCORE : 0

Actually, the answer is D

According to the NINDS and AASM,
onset of narcolepsy symptoms typically occurs
during the age range of 10-25 years old.^{1,2}

CONTINUE

TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 0

According to a quantitative survey of physicians, people with narcolepsy are often misdiagnosed with conditions including:

- A

Insomnia
- B

Depression
- C

Attention-deficit/
hyperactivity disorder
- D

All of the above

YOUR SCORE : 200

Correct, the answer is D

According to a quantitative survey of physicians, people with narcolepsy are often misdiagnosed with conditions including insomnia, depression, and attention-deficit/ hyperactivity disorder.³

CONTINUE

YOUR SCORE : 0

Actually, the answer is D

According to a quantitative survey of physicians, people with narcolepsy are often misdiagnosed with conditions including insomnia, depression, and attention-deficit/ hyperactivity disorder.³

CONTINUE

TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 100

Narcolepsy type 1 is believed to result from autoimmune destruction of _____ neurons, according to the AASM.

- A

Oxytocin
- C

Dopamine
- B

Hypocretin
- D

Norepinephrine

YOUR SCORE : 400

Correct, the answer is B

Narcolepsy type 1 is believed to result from autoimmune destruction of hypocretin neurons, according to the AASM.²

CONTINUE

YOUR SCORE : 0

Actually, the answer is B
Narcolepsy type 1 is believed to result from autoimmune destruction of hypocretin neurons, according to the AASM.²

CONTINUE

TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 400

According to analyses of medical claims data, patients with narcolepsy have a higher prevalence of cardiovascular events, such as stroke and myocardial infarction, relative to those without narcolepsy.

TRUE

FALSE

YOUR SCORE : 500

Correct, the answer is True

According to analyses of medical claims data, patients with narcolepsy have a higher prevalence of cardiovascular events, such as stroke and myocardial infarction, relative to those without narcolepsy.⁴⁻⁶


CONTINUE

YOUR SCORE : 0

Actually, the answer is True

According to analyses of medical claims data, patients with narcolepsy have a higher prevalence of cardiovascular events, such as stroke and myocardial infarction, relative to those without narcolepsy.⁴⁻⁶

CONTINUE



TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 500

According to analyses of medical claims data, patients with narcolepsy are more likely to have an increased risk of hypertension compared with the general population.

TRUE


FALSE

YOUR SCORE : 700

Correct, the answer is True

According to analyses of medical claims data, patients with narcolepsy are more likely to have an increased risk of hypertension compared with the general population.^{4,5}

CONTINUE



YOUR SCORE : 0

Actually, the answer is True

According to analyses of medical claims data, patients with narcolepsy are more likely to have an increased risk of hypertension compared with the general population.^{4,5}

CONTINUE

TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 700

According to analyses of medical claims data, diabetes is less prevalent among patients with narcolepsy relative to those without narcolepsy.

TRUE

FALSE

YOUR SCORE : 1000

Correct, the answer is False

According to analyses of medical claims data, diabetes is actually more prevalent among patients with narcolepsy relative to those without narcolepsy.⁷

CONTINUE

YOUR SCORE : 0

Actually, the answer is False

According to analyses of medical claims data, diabetes is actually more prevalent among patients with narcolepsy relative to those without narcolepsy.⁷

CONTINUE

TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 1000

What is the ideal daily target for sodium intake according to the American Heart Association?

- A

<1000 mg
- C

<2000 mg
- B

<1500 mg
- D

<2500 mg

YOUR SCORE : 1100

Correct, the answer is B

The ideal daily target for sodium intake according to the American Heart Association is <1500 mg.⁸

CONTINUE

YOUR SCORE : 0

Actually, the answer is B
The ideal daily target for sodium intake
according to the American Heart
Association is <1500 mg.⁸

CONTINUE



Jazz Pharmaceuticals

©2020 Jazz Pharmaceuticals plc or its subsidiaries US-SLE-2000396 Rev0820

REFERENCES

PRIVACY POLICY

?

TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 1100

According to a case-control study, in addition to sodium consumed in foods and beverages, high sodium-containing medications also contribute to cardiovascular complications.

TRUE

FALSE

YOUR SCORE : 1300

Correct, the answer is True

According to a case-control study, in addition to sodium consumed in foods and beverages, high sodium-containing medications also contribute to cardiovascular complications.^{9,10}

CONTINUE

YOUR SCORE : 0

Actually, the answer is True

According to a case-control study, in addition to sodium consumed in foods and beverages, high sodium-containing medications also contribute to cardiovascular complications.^{9,10}

CONTINUE

TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 1300

According to international, cross-sectional studies, daily sodium intake is associated with increase in blood pressure and risk for adverse CV outcomes, particularly stroke.

TRUE

FALSE

YOUR SCORE : 1600

Correct, the answer is True

According to international, cross-sectional studies, daily sodium intake is associated with increase in blood pressure and risk for adverse CV outcomes, particularly stroke.¹¹⁻¹⁴

CONTINUE

YOUR SCORE : 0

Actually, the answer is True

According to international, cross-sectional studies, daily sodium intake is associated with increase in blood pressure and risk for adverse CV outcomes, particularly stroke.¹¹⁻¹⁴

CONTINUE

TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 1600

According to peer-reviewed research,
narcolepsy should be considered in all
patients reporting excessive daytime sleepiness.

TRUE

FALSE

YOUR SCORE : 1700

Correct, the answer is True

According to peer-reviewed research,
narcolepsy should be considered in all
patients reporting excessive daytime sleepiness.^{15,16}

CONTINUE

YOUR SCORE : 0

Actually, the answer is True

According to peer-reviewed research,
narcolepsy should be considered in all
patients reporting excessive daytime sleepiness.^{15,16}

CONTINUE

TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 1700

According to the AASM, when considering a diagnosis of narcolepsy, which symptoms may help support the clinical diagnosis of narcolepsy and the need for sleep laboratory testing?

- A

Hypnagogic hallucinations
- B

Sleep paralysis
- C

Sleep disruption
- D

All of the above

Correct, the answer is D

YOUR SCORE : 2000

According to the AASM, when considering a diagnosis of narcolepsy, the symptoms of hypnagogic hallucinations, sleep paralysis, or sleep disruption may help support the clinical diagnosis of narcolepsy and the need for sleep laboratory testing.²

CONTINUE

Actually, the answer is D

YOUR SCORE : 0

According to the AASM, when considering a diagnosis of narcolepsy, the symptoms of hypnagogic hallucinations, sleep paralysis, or sleep disruption may help support the clinical diagnosis of narcolepsy and the need for sleep laboratory testing.²

CONTINUE



Jazz Pharmaceuticals

©2020 Jazz Pharmaceuticals plc or its subsidiaries US-SLE-2000396 Rev0820

REFERENCES

PRIVACY POLICY



TIME REMAINING	XX
POINTS	XX

YOUR SCORE : 2000

According to a study of patients with narcolepsy, cataplexy may be difficult to recognize because patients may associate the sudden muscle weakness with certain situations rather than specific emotions.

TRUE

FALSE

YOUR SCORE : 2200

Correct, the answer is True

According to a study of patients with narcolepsy, cataplexy may be difficult to recognize because patients may associate the muscle weakness with certain situations rather than specific emotions.¹⁷

CONTINUE



YOUR SCORE : 0

Actually, the answer is True

According to a study of patients with narcolepsy, cataplexy may be difficult to recognize because patients may associate the muscle weakness with certain situations rather than specific emotions.¹⁷

CONTINUE





YOUR SCORE : 2200

HOW STRONG ARE YOUR SKILLS? BONUS QUESTION

BET YOUR POINTS 

Bet up to your current point total.



Jazz Pharmaceuticals

©2020 Jazz Pharmaceuticals plc or its subsidiaries US-SLE-2000396 Rev0820

[REFERENCES](#)

[PRIVACY POLICY](#)



YOUR SCORE : 2200

BONUS QUESTION: 2050 POINTS

An analysis of NHANES data indicates that Americans have an average daily dietary sodium intake of:

- A

2100 mg
- B

2700 mg
- C

3100 mg
- D

3400 mg

YOUR SCORE : 3750

Correct, the answer is D

An analysis of NHANES data indicates that Americans have an average daily dietary sodium intake of 3400 mg.⁹

CONTINUE

YOUR SCORE : 2200

Actually, the answer is D

An analysis of NHANES data indicates that Americans have an average daily dietary sodium intake of 3400 mg.⁹

CONTINUE



YOUR SCORE : 2200

HOW STRONG ARE YOUR SKILLS?

FINAL SCORE 2200

POST MY SCORE

REPLAY



YOUR SCORE : 2200

HOW STRONG ARE YOUR SKILLS?

FIRST LETTER
OF FIRST NAME

FIRST INITIAL (LETTER ONLY)

FIRST THREE LETTERS
OF LAST NAME

LAST NAME (LETTERS ONLY)

MEDICAL SCHOOL

SCHOOL NAME (LETTERS ONLY PLEASE)

CONTINUE



Jazz Pharmaceuticals

©2020 Jazz Pharmaceuticals plc or its subsidiaries US-SLE-2000396 Rev0820

REFERENCES

PRIVACY POLICY





YOUR SCORE : 2200

HOW STRONG ARE



SHARE

Share the app with your colleagues, see how they compare!



HTTP://SLEEPTRIVIA.S3-WEBSITE-US-WEST-1.A



Jazz Pharmaceuticals

©2020 Jazz Pharmaceuticals plc or its subsidiaries US-SLE-2000396 Rev0820

REFERENCES

PRIVACY POLICY





YOUR SCORE : 3750

HOW STRONG ARE YOUR SKILLS?

**ARE YOU SURE YOU WANT TO
REPLAY THE GAME?**

CANCEL

CONTINUE