L^2-BETTI NUMBERS FROM ALGEBRAIC POINT OF VIEW

JIANRU DUAN

Abstract. Fifty years ago, Atiyah asked a question that is now known as the following Atiyah Conjecture: for any group G, are the L^2-Betti numbers of any G-CW-complex always rational?

This short course will focus on Linnell's proof of the (strong) Atiyah Conjecture for a large and concrete class of groups. His proof is based on an elegant algebraization of L^2-Betti numbers via Lück's dimension theory.

For a torsion-free group G, the Atiyah Conjecture predicts the existence of a canonical division ring $\mathcal{D}G$ extending the group ring $\mathbb{Z}G$. The torsion theory over $\mathcal{D}G$ yields the universal L^2-torsion. If time permits, we will demonstrate how the universal L^2-torsion detects the fiberedness of most 3-manifolds.

1. Lecture 1

One thing that a topologist, an algebraist and a group theorist would have in common is they love to study a group G by looking at its action on a mathematical object. Especially:

- $G \curvearrowright A$ where A is an abelian group. This makes A a $\mathbb{Z}G$*-module*.
- $G \curvearrowright X$ cellularly where X is a CW-complex. This makes X a *G-CW complex*. In particular, the homology group of X is a $\mathbb{Z}G$-module.

When G is infinite, the topology of X and its homology group group $H_*(X)$ is typically not of finite type. An idea is to construct the " $\mathbb{Z}G$-dimension" of such modules.

Question 1.1. *Define a "$\mathbb{Z}G$-dimension" for every $\mathbb{Z}G$-module M, such that:*

(1) $\dim_G(\mathbb{Z}G)^n = n$.

(2) Suppose $0 \to M_1 \to M \to M_2 \to 0$ *is exact, then*

$$\dim_G M = \dim_G M_1 + \dim_G M_2.$$

The desired $\mathbb{Z}G$-dimension does not exist for a simple reason:

Example 1.2. Consider the free group of rank two $F = \langle g_1, g_2 \rangle$. Identify F with the fundamental group of $S^1 \vee S^1$, then F naturally acts on the universal cover X, a infinite tree of degree 4.

The cellular chain complex of X is finitely generated as $\mathbb{Z}F$-module.

$$C_*(X) = 0 \to \mathbb{Z}F \oplus \mathbb{Z}F \xrightarrow{\begin{pmatrix} g_1 - 1 \\ g_2 - 1 \end{pmatrix}} \mathbb{Z}F \to 0.$$

It is easy to see that $\partial : \mathbb{Z}F \oplus \mathbb{Z}F \to \mathbb{Z}F$ is injective. In fact, any chain of $C_1(X)$ is supported on finitely many edges, hence have non-trivial endpoints. Similarly,

Date: April 17, 2026.

This is the lecture notes for the author's minicourse at SIMIS, April 2026.

there is an embedding of $\mathbb{Z}F$-modules $\mathbb{Z}F \oplus \mathbb{Z}F \hookrightarrow \mathbb{Z}F$ for any free group F of rank greater than 1.

In fact, we have the following theorem. The if-part is proved by Ceccherini-Silberstein and Coornaert [CSC06], the only-if-part is proved by Bartholdi [Bar19] and also Kropholler–Lorensen [KL19].

Theorem 1.3. *A group G is non-amenable if and only if there exists $n > 0$ such that $\mathbb{Z}G^{n+1} \hookrightarrow \mathbb{Z}G^n$.*

This suggests that the ring $\mathbb{Z}G$ is algebraically highly complicated. However, we do have a beautiful dimension theory if we pass to the Hilbert space completion of $\mathbb{Z}G$.

1.1. **von Neumann dimension.**

Definition 1.4. Let G be a group. Define $\ell^2(G)$ to be the Hilbert space spanned by elements of G. Namely,

$$\ell^2(G) = \Big\{ \sum_{g\in G} c_g \cdot g \;\Big|\; c_g \in \mathbb{C},\ \sum_{g\in G} |c_g|^2 < \infty \Big\}.$$

The inner product is given by

$$\Big\langle \sum_{g\in G} c_g \cdot g, \sum_{g\in G} d_g \cdot g \Big\rangle = \sum_{g\in G} c_g \overline{d_g}.$$

There is a natural isometric G-action on both sides of $\ell^2(G)$.

We wish to define the dimension of the following class of spaces:

Definition 1.5 (Hilbert G-module)**.** A finitely generated *Hilbert G-module* is a Hilbert space V together with an isometric left G-linear action, such that there exists an isometric G-embedding

$$V \hookrightarrow \ell^2(G)^n$$

for some n. The *morphisms* between Hilbert G-modules are bounded G-linear maps.

Example 1.6. Any closed G-invariant subspace of $\ell^2(G)^n$ is a f.g. Hilbert G-module. Conversely, any f.g. Hilbert G-module is G-linearly isomorphic to a closed G-invariant subspace of $\ell^2(G)^n$. Therefore

$$\{\text{Hilbert } G\text{-modules}\}/\sim \quad \overset{1:1}{\longleftrightarrow} \quad \{\text{closed } G\text{-invariant subspaces of } \ell^2(G)^n\}/\sim .$$

Definition 1.7 (von Neumann algebra)**.** Define the von Neumann algebra

$$\mathcal{N}G = \{\text{bounded } G\text{-linear operators } f : \ell^2(G) \to \ell^2(G)\}.$$

Remark 1.8. It is easy to verify the following facts:

(1) $\mathcal{N}G$ is a $\mathbb{C}$-algebra. In particular $\mathbb{Z}G \subset \mathcal{N}G$ by right multiplication.
(2) An operator $f \in \mathcal{N}G$ is determined by the value $f(e) \in \ell^2(G)$. Therefore, as a set,

$$\mathcal{N}G = \{x \in \ell^2(G) \mid ax \in \ell^2(G),\ \forall a \in \ell^2(G)\}.$$

(3) Note that $\ell^2(G)$ is in general not a ring.

Theorem 1.9 (L^p-conjecture, [Sae90])**.** *Suppose G is a infinite discrete group, then $\ell^p(G) \neq \ell^p(G) \cdot \ell^p(G)$ for any $p > 1$.*

Suppose we enumerate the element of G as $\{g_i\}_{i \in I}$, then the matrix $\mathcal{A}$ representing $f \in \mathcal{N}G$ has the form

$$\mathcal{A}_{ij} = \langle f(g_i), g_j \rangle = \langle g_i f(e), g_j \rangle = \langle f(e), g_i^{-1} g_j \rangle.$$

Hence the diagonal elements are constantly $\mathrm{tr}_{\mathcal{N}G}(f)$.

Definition 1.10 (von Neumann trace)**.** The von Neumann trace is a $\mathbb{C}$-linear map $\mathrm{tr}_{\mathcal{N}G} : \mathcal{N}G \to \mathbb{C}$ defined as

$$\mathrm{tr}_{\mathcal{N}G}\, f := \langle f(e), e \rangle, \quad \forall f \in \mathcal{N}G.$$

There are at least two natural topologies on $\mathcal{N}G$, all inherited from $\mathcal{B}(\ell^2(G))$: the norm topology, and the weak topology. The von Neumann trace is continuous with respect to both topology. Moreover, as a trace function, we certainly have

$$\mathrm{tr}_{\mathcal{N}G}(fg) = \mathrm{tr}_{\mathcal{N}G}(gf), \quad f, g \in \mathcal{N}G.$$

The trace can be used to characterize the dimension of a vector space. Similarly, we have the following definitions.

Definition 1.11 (von Neumann dimension)**.** Suppose $V \subset \ell^2(G)$ is a closed G-invariant subspace. Let $\mathrm{pr} : \ell^2(G) \to \ell^2(G)$ be the orthogonal projection onto V. Define the *von Neumann dimension* of V as

$$\dim_{\mathcal{N}G} V := \mathrm{tr}_{\mathcal{N}G}(\mathrm{pr}) \in [0, 1].$$

Similarly, we define the von Neumann dimension for any closed G-invariant subspace $V \subset \ell^2(G)^n$:

$$V \longmapsto \dim_{\mathcal{N}G} V \in [0, n].$$

Finally, define the von Neumann dimension of a fin. gen. Hilbert G-module M using the embedding $M \hookrightarrow \ell^2(G)^n$.

Remark 1.12. (1) The von Neumann dimension is independent of the choice of the embedding.

Proof. Assume that there are two isometric G-embeddings i, j and projections A, B, respectively.

$$\begin{tikzcd} \ell^2(G)^n \arrow[r, bend left, "A"] \arrow[r, shift left, "j"] & M \arrow[l, shift left, "i"] \arrow[l, shift right] \arrow[l, bend right, "B"] \end{tikzcd}$$

Then the isomorphism $\mathrm{im}\, A \cong \mathrm{im}\, B$ trivially extends to a morphism $U : \ell^2(G)^n \to \ell^2(G)^n$ by zero on the orthogonal complements. Therefore $j = U \circ i$, $A = B \circ U$, and.

$$\mathrm{tr}_{\mathcal{N}G}(j \circ B) = \mathrm{tr}_{\mathcal{N}G}(U \circ i \circ B) = \mathrm{tr}_{\mathcal{N}G}(i \circ B \circ U) = \mathrm{tr}_{\mathcal{N}G}(i \circ A).$$

□

(2) (Regularity) For any fin. gen. Hilbert G-module M,

$$\dim_{\mathcal{N}G} M = 0 \iff M = 0.$$

(3) (Additivity) If $0 \to M_1 \to M \to M_2 \to 0$ is weakly exact, then

$$\dim_{\mathcal{N}G} M = \dim_{\mathcal{N}G} M_1 + \dim_{\mathcal{N}G} M_2.$$

Example 1.13. Consider the free abelian group $\mathbb{Z}^n$. By standard Fourier transformation, we can identify $\ell^2(\mathbb{Z}^n)$ with $L^2(T^n)$, the complex Hilbert space of L^2-integrable functions on the n-dimensional torus $T^n = (S^1)^n$. The basis element $(k_1, \dots, k_n) \in \mathbb{Z}^n$ is sent to the function

$$f_{k_1,\dots,k_n} : T^n \to \mathbb{C}, \quad (z_1, \dots, z_n) \mapsto z_1^{k_1} \cdots z_n^{k_n}.$$

The inner product of $f, g \in L^2(T^n)$ is given by

$$\langle f, g \rangle := \frac{1}{(2\pi)^n} \int_{T^n} f(w)\overline{g(w)} dw.$$

The von Neumann algebra $\mathcal{N}\mathbb{Z}^n$ has the description

$$\begin{aligned} \mathcal{N}\mathbb{Z}^n &= \{x \in \ell^2(\mathbb{Z}^n) \mid ax \in \ell^2(\mathbb{Z}^n),\ \forall a \in \ell^2(\mathbb{Z}^n)\} \\ &= \{f \in L^2(T^n) \mid gf \in L^2(T^n),\ \forall g \in L^2(T^n)\} \\ &= L^\infty(T^n). \end{aligned}$$

The von Neumann trace $\mathrm{tr}_{\mathcal{N}\mathbb{Z}^n} : \mathcal{N}\mathbb{Z}^n \to \mathbb{C}$ is given by

$$\mathrm{tr}_{\mathcal{N}\mathbb{Z}^n} : L^\infty(T^n) \to \mathbb{C}, \quad \mathrm{tr}_{\mathcal{N}\mathbb{Z}^n}(f) := \langle f, 1 \rangle = \frac{1}{(2\pi)^n} \int_{T^n} f.$$

A closed G-invariant subspace $M \subset \ell^2(\mathbb{Z}^n)$ is characterized by an orthogonal projection $\mathrm{pr} \in \mathcal{N}\mathbb{Z}^n$. Namely, pr is an self-adjoint idempotent. These corresponds to the characteristic functions

$$\chi_A : T^2 \to \mathbb{C}, \quad A \subset T \text{ measurable set.}$$

The von Neumann dimension of M is given by $\mathrm{Vol}(A) \in [0, 1]$. In conclusion,

$\ell^2(\mathbb{Z}^n)$	$\mathcal{N}\mathbb{Z}^n$	$\mathrm{tr}_{\mathcal{N}\mathbb{Z}^n}$	submodule $M \subset \ell^2(\mathbb{Z}^n)$	$\dim_{\mathcal{N}\mathbb{Z}^n} M$
$L^2(T^n)$	$L^\infty(T^n)$	$f \mapsto \int_{T^n} f$	measurable set $A \subset T^n$	$\mathrm{Vol}(A)/(2\pi)^n$

Example 1.14. Recall from Example 1.2 the F-CW complex X for free group $F = \langle g_1, g_2 \rangle$.

$$C_*(X) = 0 \to \mathbb{Z}F \oplus \mathbb{Z}F \xrightarrow{\begin{pmatrix} g_1 - 1 \\ g_2 - 1 \end{pmatrix}} \mathbb{Z}F \to 0.$$

If we pass to the L^2-completion, i.e. we allow the cellular chains to be infinite:

$$C_*^{(2)}(X) = 0 \to \ell^2(F) \oplus \ell^2(F) \xrightarrow{\begin{pmatrix} g_1 - 1 \\ g_2 - 1 \end{pmatrix}} \ell^2(F) \to 0.$$

Observe that:

- $\operatorname{im}\partial$ contains the augmentation ideal $\mathcal{I} \subset \mathbb{Z}F \subset \ell^2(F)$, a dense subset of $\ell^2(F)$. Therefore
$$\overline{\operatorname{im}\partial} = \ell^2(F).$$
(In fact, the 0-th L^2-Betti number of any CW-complex with infinite fundamental group is zero.)
- $0 \to \ker\partial \to \ell^2(F) \oplus \ell^2(F) \xrightarrow{\partial} \overline{\operatorname{im}\partial} \to 0$ is weakly exact, therefore
$$\dim_{\mathcal{N}G}(\ker\partial) = 1.$$

Definition 1.15 (L^2-homology)**.** Let G acts freely, cellularly and cocompactly on a CW-complex X. Define
$$\begin{aligned} C_*^{(2)}(X) &= \ell^2(G) \otimes_{\mathbb{Z}G} C_*(X), \quad (L^2\text{-cellular chain complex}) \\ H_i^{(2)}(X) &= \ker\partial_i^{(2)}/\overline{\operatorname{im}\partial_{i+1}^{(2)}}, \quad (L^2\text{-homology, a Hilbert } G\text{-module}) \\ \beta_i^{(2)}(X) &= \dim_{\mathcal{N}G} H_i^{(2)}(X) \quad (L^2\text{-Betti numbers}). \end{aligned}$$
In the special case that X is the universal cover of a finite connected CW-complex Y with obvious $\pi_1(Y)$-action, we often abbreviate the above by the corresponding invariants of Y.

Example 1.16. When $Y =$  , we have $\beta_1^{(2)}(Y) = 1$, $\beta_0^{(2)}(Y) = 0$.

There are many standard topological properties of the L^2-homology: *homotopy invariance*, *Künneth formula*, *Poincaré duality*, *Euler–Hopf formula*, and so on. These can be found in any textbook on L^2-invariants. Actually one can even guess their meaning from the name.

1.2. **Extended von Neumann dimension.** There are two restrictions in the above definition of L^2-Betti numbers.

(1) We would like a more algebraic-flavored definition of L^2-Betti number, instead of caring about functional analysis.
(2) There are lots of interesting CW-complex with group action which is not cocompact, such as $G \curvearrowright EG$ where EG is the universal cover of the classifying space BG. Therefore, we have to study Hilbert G-modules which is *not* finitely generated.

In the above discussion, we were working in the following category:

Definition 1.17. The category $\mathcal{C}_1$ of finitely generated Hilbert G-module is defined as follows:
$$\begin{aligned} \operatorname{Obj}(\mathcal{C}_1) &= \{\text{fin. gen. Hilbert } G\text{-modules}\}, \\ \operatorname{Mor}(\mathcal{C}_1) &= \{\text{bounded } G\text{-linear maps}\}. \end{aligned}$$

View $\mathcal{N}G$ as a abstract ring, we have the following category:

Definition 1.18. The category $\mathcal{C}_2$ of finitely generated projective $\mathcal{N}G$-module is defined as follows:
$$\begin{aligned} \operatorname{Obj}(\mathcal{C}_2) &= \{\text{fin. gen. projective } \mathcal{N}G\text{-modules}\}, \\ \operatorname{Mor}(\mathcal{C}_2) &= \{\mathcal{N}G\text{-linear maps}\}. \end{aligned}$$

Theorem 1.19. *The categories $\mathcal{C}_1, \mathcal{C}_2$ are equivalent. Namely, there are functors*

$$\nu : \mathcal{C}_2 \to \mathcal{C}_1 \quad \text{and} \quad \nu^{-1} : \mathcal{C}_1 \to \mathcal{C}_2$$

inverses to each other up to natural equivalences.

Proof. The proof roughly goes as follows: construct the functor ν_1 that sends the regular Hilbert G-modules $\ell^2(G)^n$ to the free $\mathcal{N}G$-module $\mathcal{N}G^n$. A morphism $A : \ell^2(G)^n \to \ell^2(G)^m$ is a matrix over $\mathcal{N}G$, hence naturally a morphism $\nu^{-1}(A) : \mathcal{N}G^n \to \mathcal{N}G^m$. And vise versa.

Generally, a finitely generated Hilbert G-module can be identified with a pair $(\ell^2(G)^n, p)$, where $p : \ell^2(G)^n \to \ell^2(G)$ is an *idempotent*, namely p is a idempotent matrix over $\mathcal{N}G$. The functor ν sends the pair $(\ell^2(G)^n, p)$ to $(\mathcal{N}G^n, p)$, identifying a projective $\mathcal{N}G$-module $p(\mathcal{N}G^n)$. □

Definition 1.20 (Extended von Neumann dimension)**.** Define the *extended von Neumann dimension* $\dim_{\mathcal{N}G}(M)$ for arbitrary $\mathcal{N}G$-module M:

- If M is finitely generated projective, define
$$\dim_{\mathcal{N}G} M := \dim_{\mathcal{N}G}(\nu(M)).$$
- If M is any $\mathcal{N}G$-module. Define its *extended von Neumann dimension*
$$\dim_{\mathcal{N}G}(M) := \sup\{\dim_{\mathcal{N}G} P \mid P \subset M \text{ fin. gen. projective submodule}\} \in [0, +\infty].$$

Definition 1.21 (Extended L^2-Betti number)**.** Let X be a topological space, with continuous G-action. The homology of X with coefficient in $\mathcal{N}G$

$$H_*(X; \mathcal{N}G)$$

is the homology of the $\mathcal{N}G$-chain complex

$$\mathcal{N}G \otimes_{\mathbb{Z}G} C_*^{\text{sing}}(X).$$

The L^2-Betti number of X is given by

$$\beta_p^{(2)}(X) := \dim_{\mathcal{N}G}(H_p^G(X; \mathcal{N}G)) \in [0, \infty].$$

The L^2-Betti number of a discrete group G is defined by

$$\beta_p^{(2)}(G) := \beta_p^{(2)}(EG; \mathcal{N}G).$$

Remark 1.22. (1) When X is a free, cocompact G-CW complex, then the extended L^2-Betti number agrees with the original one.

(2) Given a chain complex

$$\cdots \to \ell^2(G)^{r_{n+1}} \xrightarrow{\partial_{n+1}} \ell^2(G)^{r_n} \xrightarrow{\partial_n} \ell^2(G)^{r_{n-1}} \to \cdots.$$

The usual homology is not a Hilbert space. We have to define

$$H_n^{(2)} := \ker \partial_n / \overline{\operatorname{im} \partial_{n+1}}.$$

However, when viewed as chain complex of von Neumann algebras

$$\cdots \to \mathcal{N}G^{r_{n+1}} \xrightarrow{\partial_{n+1}} \mathcal{N}G^{r_n} \xrightarrow{\partial_n} \mathcal{N}G^{r_{n-1}} \to \cdots.$$

The classical homology is a well-defined $\mathcal{N}G$-module (though not projective in general)

$$H_n := \ker \partial_n / \operatorname{im} \partial_{n+1}.$$

And we have $\dim_{\mathcal{N}G} H_n = \dim_{\mathcal{N}G} H_n^{(2)}$.

(3) The tradeoff is that for an arbitrary $\mathcal{N}G$-module M,

$$\dim_{\mathcal{N}G} M = 0 \quad \text{does not imply} \quad M = 0.$$

Lück [Lüc02, Section 6] had a belief that the von Neumann algebra is very similar to the ring $\mathbb{Z}$ of integers with two exceptions: (when G is infinite) $\mathcal{N}G$ is not Noetherian, and $\mathcal{N}G$ has non-trivial zero-divisors. The von Neumann dimension of a $\mathcal{N}G$-module is comparable with the $\mathbb{Z}$-rank of a $\mathbb{Z}$-module. In the next lecture we will consider the "localization" of $\mathcal{N}G$: the affiliated algebra $\mathcal{U}G$. Naturally, $\mathcal{U}G$ to $\mathcal{N}G$ is like $\mathbb{Q}$ to $\mathbb{Z}$.

Let me mention the following beautiful theorem.

Theorem 1.23 (Lück [Lüc98], Kropholler–Lorensen [KL19])**.** *A group G is amenable if and only if*

$$\dim_{\mathcal{N}G}(\mathrm{Tor}_p^{\mathbb{C}G}(M, \mathcal{N}G)) = 0$$

for all $p \geqslant 0$ and all $\mathbb{Z}G$-module M.

Corollary 1.24. *If G is an infinite amenable group, then the L^2-Betti numbers of G vanish.*

2. Lecture 2

2.1. **Atiyah Conjecture.** Fix a group G. Let's look at the set of all possible L^2-Betti numbers.

Proposition 2.1. *For any group G. The following sets are the same:*

(1) $\Lambda_1 := \{\beta_p^{(2)}(M; \mathcal{N}G) \mid M$ *is a free, cocompact G-manifold,* $p \in \mathbb{Z}\}$.
(2) $\Lambda_2 := \{\beta_p^{(2)}(X; \mathcal{N}G) \mid X$ *is a free, cocompact G-CW complex,* $p \in \mathbb{Z}\}$.
(3) $\Lambda_3 := \{\dim_{\mathcal{N}G}(\ker r_A) \mid r_A : \ell^2(G)^n \to \ell^2(G)^m$ *given by a matrix A over* $\mathbb{Z}G\}$.
(4) $\Lambda_4 := \{\dim_{\mathcal{N}G}(\mathcal{N}G \otimes_{\mathbb{Z}G} M) \mid M$ *is a finitely presented $\mathbb{Z}G$-module*$\}$.

Therefore, we denote this set by $\Lambda(G) \subset [0, +\infty)$.

Proof. We piece together proofs from [Lüc02, Section 10.1].

- $\Lambda_1 \subset \Lambda_2$: this is obvious.
- $\Lambda_2 \subset \Lambda_1$: embed $G\backslash X$ into $\mathbb{R}^N$ for N large enough. Let M be the regular neighborhood of $G\backslash X \subset \mathbb{R}^N$. Then ∂M is a closed manifold and the projection from the ∂M to $G\backslash X$ is $(N - \dim X - 2)$-connected.
- $\Lambda_2 = \Lambda_3$: the support of A is a finitely generated subgroup $H < G$. Realize A as a boundary map of an H-CW complex Y, then consider $X = G \times_H Y$.
- $\Lambda_3 = \Lambda_4$: suppose M is finitely presented $\mathbb{Z}G$-module with

$$\mathbb{Z}G^n \xrightarrow{r_A} \mathbb{Z}G^m \to M \to 0$$

exact. Then the $\mathcal{N}G$-complex

$$\mathcal{N}G^n \xrightarrow{r_A} \mathcal{N}G^m \to \mathcal{N}G \otimes_{\mathbb{Z}G} M \to 0$$

is also exact, with

$$\dim_{\mathcal{N}G}(\mathcal{N}G \otimes_{\mathbb{Z}G} M) = m - n + \dim_{\mathcal{N}G}(\ker r_A).$$

□

Remark 2.2. The proposition states for all groups, but essentially reduces to finitely generated groups.

Conjecture 2.3 (Atiyah [Ati76])**.**

$$\Lambda(G) \subset \mathbb{Q}.$$

Not surprisingly, this conjecture is not true if we put no restriction on the group.

Theorem 2.4 (Grabowski [Gra14], Pichot–Schick–Zuk [PSZ15])**.**

$$\bigcup_{\text{fin. gen. } G} \Lambda(G) = \mathbb{R}_{\geqslant 0}.$$

Conjecture 2.5 (Strong Atiyah Conjecture)**.** *Let N be the least common multiple of the order of finite subgroups of G. Then*

$$\Lambda(G) \subset \frac{1}{N} \cdot \mathbb{Z}.$$

In particular, if G is torsion-free, then

$$\Lambda(G) \subset \mathbb{Z}.$$

Conjecture 2.6 (Kaplansky zero divisor conjecture)**.** *Suppose G is torsion-free, then $\mathbb{Z}G$ contains no nontrivial zero divisors.*

For torsion-free groups, the Atiyah Conjecture implies the Kaplansky Conjecture as follows: for any $x \in \mathbb{Z}G$, if x is a non-zero divisor, then $r_x : \ell^2(G) \to \ell^2(G)$ has non-trivial kernel, hence

$$\dim_{\mathcal{N}G} \ker r \in \mathbb{Z}_+,$$

therefore $\dim_{\mathcal{N}G} \ker r = 1$. This shows that $r_x = 0$, and $x = r_x(e) = 0$.

2.2. Linnell's class $\mathcal{C}$.

Definition 2.7 (Linnell's class $\mathcal{C}$)**.** Let $\mathcal{C}$ be the smallest class of groups which contains all free groups and is closed under:

- directed unions, and
- extensions by elementary amenable groups.

We remark that $\mathcal{C}$ is closed under *taking subgroups*, closed under any *free products*, and contains all *3-manifold groups* [KL24].

Theorem 2.8 (Linnell [Lin93])**.** *Let G be a group in $\mathcal{C}$ and suppose there is a upper bound on the orders of finite subgroups of G. Then G satisfies the strong Atiyah Conjecture.*

The proof is complicated and the key is to deal with finite extensions within the class $\mathcal{C}$. In this lecture we aim to look at the Atiyah Conjecture for free groups (Theorem 2.20) and for groups extended by $\mathbb{Z}$ (Theorem 2.17). We remark that the following general assertion remains open:

Conjecture 2.9 ([LS07])**.** *Suppose K satisfies the Strong Atiyah Conjecture and G is a finite extension of K. Then G satisfies the Atiyah Conjecture.*

2.3. **The algebra of affiliated operators.** We will need the following important construction. Intuitively,

$$\text{Hilbert } \mathcal{N}G\text{-module} \rightsquigarrow \mathcal{N}G\text{-module} \rightsquigarrow \mathcal{U}G\text{-module}$$
$$\mathbb{Z}^n \rightsquigarrow \mathbb{Z}\text{-module} \rightsquigarrow \mathbb{Q}\text{-module}$$

Let V, W be a Hilbert space. An *(unbounded)* operator $f : V \to W$ is a linear map defined on its domain $\mathrm{dom}(f) \subset V$. The operator f is called *closed* if its graph $\{(x, f(x)) \in V \times W \mid x \in \mathrm{dom}(f)\}$ is a closed subset.

Theorem 2.10 (Closed graph theorem)**.** *If a closed linear operator $f : \mathrm{dom}(f) \subset U \to V$ is everywhere defined, then f is bounded.*

This explains the name "unbounded operator": if an closed operator is unbounded, then its domain must be a proper subset.

Given $f \in \mathcal{N}G$, recall that x is by definition a bounded G-linear operator

$$f : \ell^2(G) \to \ell^2(G).$$

Suppose f is injective, then f is a weak isomorphism and $\mathrm{im}\, f$ is dense. The inverse $f^{-1} : \ell^2(G) \to \ell^2(G)$ is an unbounded G-linear operator, defined on the dense subset $\mathrm{im}\, f$. Moreover, the graph of f^{-1} is the same as f, hence f^{-1} is closed.

Definition 2.11 (Affiliated operators)**.** Define $\mathcal{U}G$ to be the set of densely defined, closed, (unbounded) G-linear operators.

Proposition 2.12. *The basic properties of $\mathcal{U}G$:*

(1) *Addition and composition of affiliated operators are well-defined. For any $f, g \in \mathcal{U}G$, define $(f + g) \in \mathcal{U}G$ as the closure of*
$$x \longmapsto f(x) + g(x), \quad x \in \mathrm{dom}(f) \cap \mathrm{dom}(g),$$
define $f \circ g \in \mathcal{U}G$ as the closure of
$$x \longmapsto f(g(x)), \quad x \in g^{-1}(\mathrm{dom}(f)).$$
Verifying the well-definedness is not obvious, but standard. Therefore, $\mathcal{U}G$ becomes a ring.

(2) *Any $f \in \mathcal{U}G$ can be written as $f = ab^{-1}$, where $a, b \in \mathcal{N}G$ and b is injective.*

(3) *$\mathcal{U}G$ is von Neumann regular. Equivalently, any finitely presented submodule of $\mathcal{U}G^n$ is a direct summand* [Lüc02, Lemma 8.18].

The properties (2) and (3) appears natural if one compares $\mathcal{N}G \subset \mathcal{U}G$ with $\mathbb{Z} \subset \mathbb{Q}$. Next, we extend the von Neumann dimension from $\mathcal{N}G$-modules to $\mathcal{U}G$-modules.

Definition 2.13 (Extended von Neumann dimensions)**.** For any $\mathcal{U}G$-module M, define the von Neumann dimension

$$\dim_{\mathcal{U}G} M \in [0, +\infty]$$

as follows:

- It is known that any finitely generated projective $\mathcal{U}G$-module M can be expressed as $M = \mathcal{U}G \otimes_{\mathcal{N}G} N$ where N is a finitely generated projective $\mathcal{N}G$-module [Lüc02, Theorem 8.22]. Define $\dim_{\mathcal{U}G} M := \dim_{\mathcal{N}G} N$.

- For general $\mathcal{U}G$-module M, define

$$\dim_{\mathcal{U}G} M := \limsup\{\dim_{\mathcal{U}G} P \mid P \subset M \text{ fin. gen. projective}\}.$$

This extended von Neumann dimension has all nice properties like the one defined over $\mathcal{N}G$-modules. Importantly [Lüc02, Theorem 8.29], for any $\mathcal{N}G$-module M,

$$\dim_{\mathcal{U}G}(\mathcal{U}G \otimes_{\mathcal{N}G} M) = \dim_{\mathcal{N}G} M.$$

2.4. **Linnell's theorem.** In this note, a field always refers to a *skew field*, a ring whose non-zero elements are all units. Other authors may call it a *division ring* or *division algebra*.

We are ready to prove the following beautiful criterion of Linnell:

Theorem 2.14 (Linnell [Lin93])**.** *Let G be a torsion-free group. Then the Atiyah Conjecture holds for G if and only if there exists a field F with $\mathbb{Z}G \subset F \subset \mathcal{U}G$.*

Proof. Suppose there is a field F with $\mathbb{Z}G \subset F \subset \mathcal{U}G$. Given any finitely presented $\mathbb{Z}G$-module M, we have

$$\begin{aligned}\dim_{\mathcal{N}G}(\mathcal{N}G \otimes_{\mathbb{Z}G} M) &= \dim_{\mathcal{U}G}(\mathcal{U}G \otimes_{\mathbb{Z}G} M)\\ &= \dim_{\mathcal{U}G}(\mathcal{U}G \otimes_F F \otimes_{\mathbb{Z}G} M)\\ &= \dim_{\mathcal{U}G}(\mathcal{U}G \otimes_F F^n) \quad \text{for some } n \in \mathbb{N}\\ &= \dim_{\mathcal{U}G}(\mathcal{U}G^n) = n.\end{aligned}$$

Therefore $\Lambda(G) \subset \mathbb{Z}$.

On the other hand, suppose that G satisfies the Atiyah Conjecture, we prove that there is a field between $\mathbb{Z}G \subset \mathcal{U}G$.

Definition 2.15 (Division closure)**.** The *division closure* of $\mathbb{Z}G \subset \mathcal{U}G$ is the smallest ring $\mathcal{D}G$ such that

(1) $\mathbb{Z}G \subset \mathcal{D}G \subset \mathcal{U}G$, and
(2) $\mathcal{D}G$ is *division-closed*, i.e. any $x \in \mathcal{D}G$ that is invertible in $\mathcal{U}G$ is already invertible in $\mathcal{D}G$.

It is easy to see that a field between $\mathbb{Z}G \subset \mathcal{U}G$ exists if and only if $\mathcal{D}G$ is a field.

To prove $\mathcal{D}G$ is a field, suppose the contrary that $\exists x \in \mathcal{D}G \setminus \{0\}$ not invertible in $\mathcal{U}G$. Consider the right multiplication by x

$$r_x : \mathcal{U}G \to \mathcal{U}G.$$

Since we have exact sequence

$$0 \to \ker r_x \to \mathcal{U}G \to \operatorname{im} r_x \to 0,$$

the *finitely generated* submodule $\operatorname{im}(r_x)$ is direct summand of $\mathcal{U}G$ by von Neumann regular. Hence

$$\mathcal{U}G \cong \ker(r_x) \oplus \operatorname{im}(r_x).$$

Both direct summand is not trivial, therefore has nontrivial $\mathcal{U}G$-dimension in $(0,1)$. By Additivity of $\mathcal{U}G$-dimensions $\dim_{\mathcal{U}G}(\ker r_x) \in (0,1)$.

By the *non-commutative Cramer's Rule* [Sch85, Page 53], given $x \in \mathcal{D}G$ there exists invertible matrices P, Q over $\mathcal{U}G$ and a matrix A over $\mathbb{Z}G$, such that

$$\begin{pmatrix} x & 0 \\ 0 & I \end{pmatrix} = PAQ$$

for some identity matrix I. Therefore

$$\dim_{\mathcal{U}G}(\ker r_x) = \dim_{\mathcal{U}G}(\ker(PAV)) = \dim_{\mathcal{U}G}(\ker A) = \dim_{\mathcal{N}G}(\ker A)$$

(Implicitly using the fact that $\mathcal{U}G$ is flat over $\mathcal{N}G$ for the last equality) By assumption of the Atiyah Conjecture, this is an integer, a contradiction. This proves that $x \in \mathcal{D}G$ is invertible, hence $\mathcal{D}G$ is a field. □

Example 2.16. We have identified $\mathcal{N}\mathbb{Z}^n$ with $L^\infty(T^n)$ in Example 1.13. An operator $g \in \mathcal{N}\mathbb{Z}^n$ is injective if and only if $g \in L^\infty(T^n)$ is a.e. nonzero. The algebra of affiliated operators is

$$\begin{aligned}\mathcal{U}G &= \{fg^{-1} : T^n \to \mathbb{C} \mid f, g \in L^\infty(T^n) \text{ and } g \neq 0 \text{ a.e.}\} \\ &= \{\text{measurable functions } T^n \to \mathbb{C}\}.\end{aligned}$$

Finally,

$$\begin{aligned}\mathbb{Z}\mathbb{Z}^n &= \{\text{polynomial functions } p : T^n \to \mathbb{C}\}, \quad \text{and} \\ \mathcal{D}\mathbb{Z}^n &= \{\text{rational functions } p/q : T^n \to \mathbb{C}\}.\end{aligned}$$

Since $\mathcal{D}\mathbb{Z}^n$ is a field, we conclude that $\mathbb{Z}^n$ satisfies the Atiyah Conjecture.

2.5. **Groups extended by integers.**

Theorem 2.17. *Suppose we have an exact sequence of groups*

$$1 \to K \to G \to \mathbb{Z} \to 1$$

such that K is torsion-free and satisfies the Atiyah Conjecture. Then G is torsion-free and satisfies the Atiyah Conjecture.

This directly follows from Theorem 2.14 and the following Proposition:

Proposition 2.18. *Choose $t \in G$ which maps to the generator of $\mathbb{Z}$. Then $\mathcal{D}G$ is the field of fraction of the polynomial ring $\mathcal{D}K[t^\pm]$.*

Proof. Choose $t \in G$ which maps to the generator of $\mathbb{Z}$. It is easy to see that

$$\mathbb{Z}G = \mathbb{Z}K[t^\pm]$$

is the "polynomial ring" generated by $\mathbb{Z}K$ and t. Note that since the coefficient in $\mathbb{Z}K$ is not commutative with t, the multiplication law is as follows: for any $a_i, b_j \in \mathbb{Z}K$,

$$\begin{aligned}\sum_{i\in\mathbb{Z}} a_i t^i \cdot \sum_{j\in\mathbb{Z}} b_j t^j &= \sum_{i,j\in\mathbb{Z}} a_i t^i b_j t^j \\ &= \sum_{i,j\in\mathbb{Z}} (a_i t^i b_j t^{-i}) \cdot t^{i+j}.\end{aligned}$$

This non-commutative polynomial ring structure is sometimes called a *crossed product* of the ring $\mathbb{Z}K$ and a group $\langle t\rangle$. Similarly, the polynomial ring

$$\mathbb{Z}G \subset \mathcal{D}K[t^\pm] \subset \mathcal{U}G$$

is generated by the subring $\mathcal{D}K \subset \mathcal{U}K \subset \mathcal{U}G$ and $t \in \mathbb{Z}G \subset \mathcal{U}G$. The proof proceeds as follows:

- By assumption, $\mathcal{D}K$ is a field. Therefore $\mathcal{D}K[t^\pm]$ is a PID.

- Any PID is an Ore domain. Namely, if R is a PID, then $xR \cap yR \neq \{0\}$ for any $x, y \in R \setminus \{0\}$. (Choose $z \in R$ such that $xR + yR = zR$ and write $x = zx_1$, $y = zy_1$. There are $a, b \in R$ with $z = xa + yb$. Right multiplying by x_1 we have hence $x(1 - ax_1) = ybx_1$.)
- Therefore, the Ore localization of $\mathcal{D}K[t^{\pm}]$ exists and is a field.
- Any non-zero element of $\mathcal{D}K[t^{\pm}]$ is invertible in $\mathcal{U}G$.

  Proof. Without loss of generality, we prove that $x = a_0 + a_1 t + \cdots + a_n t^n$ is invertible in $\mathcal{U}G$, where $a_0 \in \mathcal{D}K$ is nonzero. Since $\mathcal{U}K$ is the Ore localization of $\mathcal{N}K$ over non-zero-divisors, we can write

  $$x = s^{-1} \cdot (b_0 + b_1 t + \cdots + b_n t^n), \quad b_i \in \mathcal{N}K \text{ are non-zero-divisors.}$$

  In fact $(b_0 + b_1 t + \cdots + b_n t^n)$ is a non-zero-divisor in $\mathcal{N}G$ by [Lin92, Theorem 4]. Hence x is a non-zero-divisor in $\mathcal{U}G$. □

- The localization of $\mathcal{D}K[t^{\pm}]$ embeds in $\mathcal{U}G$, and exactly equals $\mathcal{D}G$.

In conclusion, $\mathcal{D}G$ is a field and G satisfies the Atiyah Conjecture. □

Corollary 2.19. *The poly-$\mathbb{Z}$ groups satisfy the Atiyah Conjecture.*

We use the special properties of free groups and the Approximation Theorem to show that free groups satisfy the Atiyah Conjecture.

Let F be a finitely generated free group. Let

$$F = F_0 \triangleright F_1 \triangleright \cdots \triangleright F_n \triangleright \cdots$$

be the lower central series, where $F_n = [F, F_{n-1}]$. By the classical Magnus embedding of F into non-commutative power series (c.f. [LS77, I.10]), we know that:

- $\cap_n F_n = \{1\}$, and
- F_n / F_{n+1} is a free abelian group.

Therefore, it is easy to show that F/F_n is poly-$\mathbb{Z}$. Now for any matrix A over $\mathbb{Z}F$. Apply the approximation theorem [Lüc02, Theorem 13.3] to the cofinal tower of quotients:

$$F \to \cdots \to F/F_n \to \cdots \to F/F_2 \to F/F_1,$$

we know that

$$\dim_{\mathcal{N}G}(\ker A) = \lim_{n \to \infty} \dim_{\mathcal{N}(F/F_n)}(\ker A_n)$$

where A_n is the image of A in $\mathbb{Z}[F/F_n]$. Since F/F_n is poly-$\mathbb{Z}$ and satisfies the Atiyah Conjecture, $\dim_{\mathcal{N}G}(\ker A)$ is the limit of a sequence of integers, hence must be an integer. Therefore, we have proved the following:

Theorem 2.20. *Any free group satisfies the Atiyah Conjecture.*

3. Lecture 3

In the following we will assume that G is a torsion-free group satisfying Atiyah Conjecture. We know from Linnell's theorem 2.14 that $\mathcal{D}G$ is a field.

3.1. **Polytope map.** Suppose $H := H_1(G;\mathbb{Z})_f$ is the free abelianization of G, then we have the exact sequence

$$1 \to K \to G \to H \to 1.$$

Choose a set $\{t_1, \dots, t_n\} \subset G$ which projects to a $\mathbb{Z}$-basis of H, here $n = b_1(G)$ is the first Betti number of G. By the argument as in Proposition 2.18, given any element $x \in \mathcal{D}G$, we can find $u, v \in \mathcal{D}K[t_1^{\pm}, \dots, t_n^{\pm}]$ such that $x = uv^{-1}$. Namely,

Proposition 3.1. *The field of fraction of the polynomial ring $\mathcal{D}K[t_1^{\pm}, \dots, t_n^{\pm}]$ is isomorphic to $\mathcal{D}G$.*

Each monomial of $\mathcal{D}K[t_1^{\pm}, \dots, t_n^{\pm}]$, say $t_1^{r_1} \cdots t_n^{r_n}$, defines an integer point

$$(r_1, \dots, r_n) \in \mathbb{R} \otimes_{\mathbb{Z}} H = H_1(G;\mathbb{R}).$$

The *Newton polytope* $\mathbb{P}(u)$ of $u \in \mathcal{D}K[t_1^{\pm}, \dots, t_n^{\pm}]$ is the convex hull in $H_1(G;\mathbb{R})$ spanned by all monomials of u. It is easy to verify that $\mathbb{P}(uv) = \mathbb{P}(u) + \mathbb{P}(v)$ for all $u, v \in \mathcal{D}K[t_1^{\pm}, \dots, t_n^{\pm}]$, where "+" refers to the *Minkowski sum* of polytopes.

Definition 3.2 (Polytope group)**.** The polytope group $\mathcal{P}(G)$ is the Grothendieck group of polytopes in $H_1(G;\mathbb{R})$ under the Minkowski sum. Namely, $\mathcal{P}(G)$ is the abelian group generated by symbols $[P]$ where P is a non-empty polytope in $H_1(G;\mathbb{R})$, subject to the relation

$$[P] + [Q] = [P + Q]$$

for each pair of non-empty polytopes P, Q.

Definition 3.3 (Polytope map)**.** Define the polytope map $\mathbb{P} : \mathcal{D}G^{\times} \to \mathcal{P}(G)$ as follows: for any $x \in \mathcal{D}G^{\times}$, write $x = uv^{-1}$ for some $u, v \in \mathcal{D}K[t_1^{\pm}, \dots, t_n^{\pm}]$ and define

$$\mathbb{P}(x) := [\mathbb{P}(u)] - [\mathbb{P}(v)].$$

This is a well-defined group homomorphism.

3.2. **Universal L^2-torsion.** Suppose X is a cocompact, free G-CW complex with trivial L^2-Betti numbers. Equivalently, the $\mathcal{D}G$-chain complex

$$\mathcal{D}G \otimes_{\mathbb{Z}G} C_*(X)$$

is $\mathcal{D}G$-exact.

After choosing a G-basis for the cells of X, we can identify $\mathcal{D}G \otimes_{\mathbb{Z}G} C_*(X)$ with a free $\mathcal{D}G$-module. This exact $\mathcal{D}G$-chain complex can be decomposed as

$$\begin{tikzcd}
\mathcal{D}G^{r_{n+1}} \arrow[dr, phantom, "\oplus"] \arrow[drr, "A_{n+1}", "\cong"'] & \mathcal{D}G^{r_n} \arrow[dr, phantom, "\oplus"] \arrow[drr, "A_n", "\cong"'] & \mathcal{D}G^{r_{n-1}} \arrow[d, phantom, "\oplus"] \\
\mathcal{D}G^{r_{n+2}} \arrow[d, equal] & \mathcal{D}G^{r_{n+1}} \arrow[d, equal] & \mathcal{D}G^{r_n} \arrow[d, equal] \\
\mathcal{D}G \otimes_{\mathbb{Z}G} C_{n+1}(X) \arrow[r, "\partial_{n+1}"] & \mathcal{D}G \otimes_{\mathbb{Z}G} C_n(X) \arrow[r, "\partial_n"] & \mathcal{D}G \otimes_{\mathbb{Z}G} C_{n-1}(X)
\end{tikzcd}$$

Definition 3.4 (Universal L^2-torsion)**.** The *universal L^2-torsion* of X is defined as

$$\tau_u^{(2)}(X) := \prod_{n \in \mathbb{Z}} \det(A_n)^{(-1)^n}.$$

Here $\det(A)$ means the element of $K_1(\mathcal{D}G)$ represented by the invertible matrix A. Alternatively, it can be understood as the *Dieudonné determinant*

$$\det : \mathrm{GL}(\mathcal{D}G) \to K_1(\mathcal{D}G) \cong (\mathcal{D}G)^{\times}_{\mathrm{ab}}.$$

Finally, the universal L^2-torsion $\tau_u^{(2)}(X)$ is well-defined (i.e. independent of the cellular basis of X) in the *Whitehead group*

$$\mathrm{Wh}(\mathcal{D}G) := K_1(\mathcal{D}G)/\pm G.$$

Definition 3.5 (L^2-torsion polytope)**.** The L^2-torsion polytope of X, denoted by $\mathbb{P}(X)$, is the image of $\tau_u^{(2)}(X)$ in the polytope group $\mathcal{P}(G)$.

Remark 3.6. In most cases, X will be the universal cover of a compact CW-complex Y with the deck transformation. In this case we also write

$$\tau_u^{(2)}(Y) := \tau_u^{(2)}(X), \quad \mathbb{P}(Y) := \mathbb{P}(X).$$

Similarly, if a group G has a finite classifying space BG, we can talk about the universal L^2-torsion $\tau_u^{(2)}(BG)$ and the L^2-torsion polytope $\mathbb{P}(BG)$. However, $\tau_u^{(2)}$ depends on the *simple-homotopy type* of BG and we do not know if $\tau_u^{(2)}(BG)$ is an invariant of G unless the Whitehead group of G is trivial. The L^2-torsion polytope $\mathbb{P}(BG)$ only depends on the *homotopy type* of BG, hence an invariant of G [Kie20, Theorem 5.17].

Example 3.7 (Groups of deficiency one)**.** Suppose G is torsion-free and admits a deficiency one presentation

$$G = \langle g_1, \dots, g_{d+1} : r_1, \dots, r_d \rangle.$$

If $\beta_1^{(2)}(G) = 0$, then any deficiency-one presentation 2-complex X is aspherical [Hil02, Section 2.3], hence a classifying space of G. The $\mathbb{Z}G$-cellular chain complex of $\widehat{X}$ has the form

$$C_*(\widehat{X}) = (0 \to \mathbb{Z}G^d \xrightarrow{(\mathcal{J}'_{ij})} \mathbb{Z}G^{d+1} \to \mathbb{Z}G \to 0)$$

where $\mathcal{J}'$ is the *Fox Jacobian* matrix over $\mathbb{Z}G$. Suppose g_{d+1} is not the trivial element in G and let $\mathcal{J}$ be the first d columns of $\mathcal{J}'$, then

$$\tau_u^{(2)}(G) = \tau_u^{(2)}(\widehat{X}) = \det(\mathcal{J}) \cdot [1 - g_{d+1}]^{-1}.$$

Example 3.8 (Universal L^2-torsion of 5_2-knot group)**.** The fundamental group of the 5_2-knot complement has the following presentation:

$$G = \langle a, b : ab^2a^{-2}b^2ab^{-3} =: w \rangle.$$

The presentation 2-complex X is aspherical, and

$$C_*(\widehat{X}) = (0 \to \mathbb{Z}G \xrightarrow{\left(\frac{\partial w}{\partial a} \quad \frac{\partial w}{\partial b}\right)} \mathbb{Z}G \oplus \mathbb{Z}G \xrightarrow{\begin{pmatrix} 1-a \\ 1-b \end{pmatrix}} \mathbb{Z}G \to 0).$$

Therefore,

$$\begin{aligned} \tau_u^{(2)}(G) &= \det\left(\frac{\partial w}{\partial b}\right) \cdot [1-a]^{-1} \\ &= [a + ab + ab^2a^{-2} + ab^2a^{-2}b - b^2 - b - 1] \cdot [1-a]^{-1}. \end{aligned}$$

Theorem 3.9 ([FL17])**.** *Let M be an admissible 3-manifold. Then the dual Thurston norm ball in $H_1(M;\mathbb{R})$ equals $2 \cdot \mathbb{P}(\tau_u^{(2)}(M))$.*

3.3. **Leading term of universal L^2-torsion.**

Definition 3.10 (Leading term map)**.** Suppose $\phi : G \to \mathbb{Z}$ is a nontrivial homomorphism whose kernel is K. Choose $t \in G$ such that $\phi(t)$ generates the image of ϕ. Identify

$$\mathcal{D}G = \mathcal{D}K[t^{\pm}] \cdot \mathcal{D}K[t^{\pm}]^{-1}.$$

For any $x \in \mathcal{D}G$, write

$$x = p \cdot q^{-1}, \quad p, q \in \mathcal{D}K[t^{\pm}],$$

define $L_\phi : \mathcal{D}G \to \mathcal{D}G$,

$$L_\phi(x) := (\text{lowest } t\text{-degree term of } p) \cdot (\text{lowest } t\text{-degree term of } q)^{-1}.$$

This induces homomorphisms

$$L_\phi : K_1(\mathcal{D}G) \to K_1(\mathcal{D}G) \quad \text{and} \quad L_\phi : \mathrm{Wh}(\mathcal{D}G) \to \mathrm{Wh}(\mathcal{D}G).$$

Theorem 3.11 ([Dua25])**.** *Let M be an admissible 3-manifold which is not a closed graph manifold. Then a cohomology class $\phi \in H^1(M;\mathbb{Z})$ is a fibered class if and only if $L_\phi(\tau_u^{(2)}(M)) = 1 \in \mathrm{Wh}(\mathcal{D}(\pi_1(M)))$.*

Remark 3.12. Suppose $L_\phi(\tau_u^{(2)}(M)) = 1$. Recall that the polytope of $\tau_u^{(2)}(M)$ is given by the dual Thurston norm ball. In particular, the level set of ϕ in $H_1(G;\mathbb{R})$ meets the boundary of the dual Thurston norm ball exactly at one of its vertex. Equivalently, ϕ lies in the top dimensional Thurston cone of $H^1(M;\mathbb{R})$.

3.4. **Restriction map.** Suppose G is a torsion-free group satisfying the Atiyah Conjecture. Let $L < G$ be a finite index normal subgroup. Then L is also torsion-free and satisfying the Atiyah Conjecture.

Lemma 3.13. *Let $d := [G : L]$. Write $G = Lg_1 \sqcup \cdots \sqcup Lg_d$, then*

$$\mathcal{D}G \cong \bigoplus_{i=1}^{d} \mathcal{D}L \cdot g_i,$$

identifying $\mathcal{D}G$ with $(\mathcal{D}L)^d$.

For any element $z \in \mathcal{D}G$, the right multiplication $r_z : \mathcal{D}G \to \mathcal{D}G$ is represented by a $d \times d$ matrix $\Lambda(z)$ over $\mathcal{D}L$. This matrix depends on the choice of coset representatives $\{g_i\}$. However, the determinant is independent of $\{g_i\}$.

Definition 3.14 (Restriction map)**.** Define the restriction map $\mathrm{res}_L^G : \mathcal{D}G \to K_1(\mathcal{D}L)$,

$$\mathrm{res}_L^G(z) := \det(\Lambda(z)), \quad z \in \mathcal{D}G.$$

This passes to the K_1 group and the Whitehead group:

$$\mathrm{res}_L^G : K_1(\mathcal{D}G) \to K_1(\mathcal{D}L), \quad \mathrm{res}_L^G : \mathrm{Wh}(\mathcal{D}G) \to \mathrm{Wh}(\mathcal{D}L).$$

The restriction map characterizes the change of $\tau_u^{(2)}$ under finite coverings.

Theorem 3.15 ([FL17]). *Let M be a finite CW-complex with $\pi_1(M) = G$, and let $\overline{M}$ be a finite regular covering with $\pi_1(\overline{M}) = L < G$. Then*

$$\tau_u^{(2)}(\overline{M}) = \mathrm{res}_L^G(\tau_u^{(2)}(M)).$$

The most technical lemma is the following:

Lemma 3.16 ([Dua25]). *The leading term map commutes with the restriction map:*

$$\begin{tikzcd} \mathrm{Wh}(\mathcal{D}G) \arrow[r, "\mathrm{res}_L^G"] \arrow[d, "L_\phi"'] & \mathrm{Wh}(\mathcal{D}L) \arrow[d, "L_{\phi|_L}"] \\ \mathrm{Wh}(\mathcal{D}G) \arrow[r, "\mathrm{res}_L^G"] & \mathrm{Wh}(\mathcal{D}L) \end{tikzcd}$$

where $\phi \in H^1(G;\mathbb{R})$ and $\phi|_L \in H^1(L;\mathbb{R})$ is the restriction.

Admitting this result, we can prove the following:

Proposition 3.17. *Let M be an admissible 3-manifold which is not a closed graph manifold. Then a cohomology class $\phi \in H^1(M;\mathbb{Z})$ is a fibered class if $L_\phi(\tau_u^{(2)}(M)) = 1$.*

Proof. By virtual fibering theorem [AFW15], there is a finite covering $\overline{M}$ with respect to $L < G$ such that $\phi|_L$ is a quasi-fibered class.

By the commutative diagram, we have

$$L_{\phi|_L}\,\mathrm{res}_L^G(\tau_u^{(2)}(M)) = \mathrm{res}_L^G(1) = 1,$$

namely $L_{\phi|_L}\tau_u^{(2)}(\overline{M}) = 1$. This implies that $\phi|_L$ lies in the top dimensional Thurston cone of $\overline{M}$.

Since $\phi|_L$ is quasi-fibered, it is a fibered class. Therefore ϕ is also fibered. □

The converse of Theorem 3.11 is given by the following result:

Theorem 3.18. *Suppose B_1, B_2 are disjoint subcomplexes of X such that $B_i \hookrightarrow X$ is π_1-injective and induces isomorphism on L^2-homology. If there is a homeomorphism $f : B_1 \to B_2$ and let X_f be the CW-complex obtained by gluing B_1, B_2 via f. Then*

$$L_\phi \tau_u^{(2)}(X_f) = i_* \tau_u^{(2)}(X, B_1),$$
$$L_{-\phi} \tau_u^{(2)}(X_f) = i_* \tau_u^{(2)}(X, B_2),$$

where $i : X \to X_f$ is the quotient map.





References

[AFW15] Matthias Aschenbrenner, Stefan Friedl, and Henry Wilton, *3-manifold groups*, EMS Series of Lectures in Mathematics, European Mathematical Society (EMS), Zürich, 2015. MR 3444187

[Ati76] M. F. Atiyah, *Elliptic operators, discrete groups and von Neumann algebras*, Colloque "Analyse et Topologie" en l'Honneur de Henri Cartan (Orsay, 1974), Astérisque, vol. No. 32-33, Soc. Math. France, Paris, 1976, pp. 43–72. MR 420729

[Bar19] Laurent Bartholdi, *Amenability of groups is characterized by Myhill's theorem*, J. Eur. Math. Soc. (JEMS) **21** (2019), no. 10, 3191–3197, With an appendix by Dawid Kielak. MR 3994103

[CSC06] Tullio Ceccherini-Silberstein and Michel Coornaert, *The Garden of Eden theorem for linear cellular automata*, Ergodic Theory Dynam. Systems **26** (2006), no. 1, 53–68. MR 2201937

[Dua25] Jianru Duan, *Universal L^2-torsion and sutured decomposition for 3-manifolds*, 2025, arXiv preprint arXiv:2512.01305.

[FL17] Stefan Friedl and Wolfgang Lück, *Universal L^2-torsion, polytopes and applications to 3-manifolds*, Proc. Lond. Math. Soc. (3) **114** (2017), no. 6, 1114–1151. MR 3661347

[Gra14] Łukasz Grabowski, *On Turing dynamical systems and the Atiyah problem*, Invent. Math. **198** (2014), no. 1, 27–69. MR 3260857

[Hil02] J. A. Hillman, *Four-manifolds, geometries and knots*, Geometry & Topology Monographs, vol. 5, Geometry & Topology Publications, Coventry, 2002. MR 1943724

[Kie20] Dawid Kielak, *The Bieri-Neumann-Strebel invariants via Newton polytopes*, Invent. Math. **219** (2020), no. 3, 1009–1068. MR 4055183

[KL19] Peter Kropholler and Karl Lorensen, *Group-graded rings satisfying the strong rank condition*, J. Algebra **539** (2019), 326–338. MR 3997173

[KL24] Dawid Kielak and Marco Linton, *Group rings of three-manifold groups*, Proc. Amer. Math. Soc. **152** (2024), no. 5, 1939–1946. MR 4728464

[Lin92] Peter A. Linnell, *Zero divisors and $L^2(G)$*, C. R. Acad. Sci. Paris Sér. I Math. **315** (1992), no. 1, 49–53. MR 1172405

[Lin93] ———, *Division rings and group von Neumann algebras*, Forum Math. **5** (1993), no. 6, 561–576. MR 1242889

[LS77] Roger C. Lyndon and Paul E. Schupp, *Combinatorial group theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas], vol. Band 89, Springer-Verlag, Berlin-New York, 1977. MR 577064

[LS07] Peter Linnell and Thomas Schick, *Finite group extensions and the Atiyah conjecture*, J. Amer. Math. Soc. **20** (2007), no. 4, 1003–1051. MR 2328714

[Lüc98] Wolfgang Lück, *Dimension theory of arbitrary modules over finite von Neumann algebras and L^2-Betti numbers. I. Foundations*, J. Reine Angew. Math. **495** (1998), 135–162. MR 1603853

[Lüc02] ———, *L^2-invariants: theory and applications to geometry and K-theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 44, Springer-Verlag, Berlin, 2002. MR 1926649

[PSZ15] Mikaël Pichot, Thomas Schick, and Andrzej Zuk, *Closed manifolds with transcendental L^2-Betti numbers*, J. Lond. Math. Soc. (2) **92** (2015), no. 2, 371–392. MR 3404029

[Sae90] Sadahiro Saeki, *The L^p-conjecture and Young's inequality*, Illinois J. Math. **34** (1990), no. 3, 614–627. MR 1053566

[Sch85] A. H. Schofield, *Representation of rings over skew fields*, London Mathematical Society Lecture Note Series, vol. 92, Cambridge University Press, Cambridge, 1985. MR 800853

Beijing International Center for Mathematical Research, Peking University, No. 5 Yiheyuan Road, Haidian District, Beijing 100871, China

Email address: duanjr@stu.pku.edu.cn