

Zero-inflated Regression models

Zero-inflated Regression models - Summary

- ▶ Zero-inflated models attempt to account for excess zeros.
- ▶ In other words, two kinds of zeros are thought to exist in the data, "**true zeros**" and "**excess zeros**".

Zero-inflated Regression models

Two Distinct Processes

- ▶ The two parts of the a zero-inflated model are a binary model, usually a logit model to model which of the two processes the zero outcome is associated with and a count model, in this case, a negative binomial model, to model the count process.
- ▶ In other words, the excess zeros are generated by a separate process from the count values and that the excess zeros can be modelled independently.
- ▶ Zero-inflated models estimate two equations simultaneously, one for the count model and one for the excess zeros.
- ▶ The expected count is expressed as a combination of the two processes.

The `zeroinfl()` function

- ▶ In R, zero-inflated count data models can be fitted with the `zeroinfl()` function from the **pscl** package.
- ▶ Both the fitting function interface and the returned model objects of class “`zeroinfl`” are modelled after the corresponding GLM functionality in R.

The arguments of `zeroinfl()` are given by

```
zeroinfl(formula, data, subset, na.action,  
          weights, offset,  
          dist = "poisson", link = "logit",  
          control = zeroinfl.control(...),  
          model = TRUE, y = TRUE, x = FALSE, ...)
```

Zero-inflated Regression models

Fishing Data Set

- ▶ We have data on 250 groups that went to a park.
- ▶ Each group was questioned about how many fish they caught (**count**), how many children were in the group (**child**), how many people were in the group (**persons**), and whether or not they brought a camper to the park (**camper**).
- ▶ In addition to predicting the number of fish caught, there is interest in predicting the existence of excess zeros, i.e., the probability that a group caught zero fish.
- ▶ We will use the variables child, persons, and camper in our model.

Zero-inflated Regression models

Fishing Data Set

- ▶ In addition to predicting the number of fish caught, there is interest in predicting the existence of excess zeros, i.e., the probability that a group caught zero fish.
- ▶ We will use the variables child, persons, and camper in our model.

```
> head(fish)
```

	nofish	livebait	camper	persons	child	xb
1	1	0	0	1	0	-0.8963146
2	0	1	1	1	0	-0.5583450
3	0	1	0	1	0	-0.4017310
4	0	1	1	2	1	-0.9562981
5	0	1	0	1	0	0.4368910
6	0	1	1	4	2	1.3944855

```
zg count
```

1	3.0504048	0
2	1.7461489	0
3	0.2799389	0
4	-0.6015257	0
5	0.5277091	1
6	-0.7075348	0

What is a Zero-Inflated Model?

The Fishing Example

- ▶ A zero-inflated model assumes that zero outcome is due to two different processes.
- ▶ For instance, in the example of fishing presented here, the two processes are that a subject has gone 1. *fishing* vs. 2. *not fishing*.
- ▶ If not gone fishing, the only outcome possible is zero.
- ▶ If gone fishing, it is then a count process.

$$\begin{aligned} &E(\text{no. fish caught} = k) \\ &= P(\text{not fishing}) \times 0 + P(\text{fishing}) \times E(y = k | \text{fishing}) \end{aligned}$$