

## Vuong Testing

- ▶ Note that the model output above does not indicate in any way if our zero-inflated model is an improvement over a standard negative binomial regression.
- ▶ We can determine this by running the corresponding standard negative binomial model and then performing a Vuong test of the two models.
- ▶ We use the MASS package to run the standard negative binomial regression.

# Vuong Testing

```
library(MASS)
summary(m2 <- glm.nb(count ~ child + camper, data = zinb))
.....
```

# Vuong Testing

```
## Coefficients:
```

##	Estimate	Std. Error	z value	Pr(> z )	
## (Intercept)	1.073	0.242	4.42	9.7e-06	***
## child	-1.375	0.196	-7.03	2.1e-12	***
## camper1	0.909	0.284	3.21	0.0013	**

```
## ---
```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1
```

## Vuong Testing

```
vuong(m1, m2)

## Vuong Non-Nested Hypothesis Test-Statistic: 1.702
## (test-statistic is asymptotically distributed N(0,1) under
## null that the models are indistinguishable)
## in this case:
## model1 > model2, with p-value 0.0444
```

- ▶ The log odds of being an excessive zero would decrease by 1.67 for every additional person in the group.
- ▶ In other words, the more people in the group the less likely that the zero would be due to not gone fishing.
- ▶ Put plainly, the larger the group the person was in, the more likely that the person went fishing.
- ▶ The Vuong test suggests that the zero-inflated negative binomial model is a significant improvement over a standard negative binomial model.