**Title**: Qualitative studies of advective competition system with Beddington–DeAngelis functional response

Running title: Beddington-DeAngelis competition system

Authors: Ling Jin, Qi Wang, Zengyan Zhang

Corresponding author: Qi Wang

Email: qwang@swufe.edu.cn

Address: Department of Mathematics

Southwestern University of Finance and Economics

555 Liutai Ave, Wenjiang

Chengdu, China 611130

# Qualitative studies of advective competition system with Beddington–DeAngelis functional response

Ling Jin, Qi Wang \* Zengyan Zhang

Department of Mathematics

Southwestern University of Finance and Economics

Chengdu, China 611130

#### Abstract

This paper investigates a reaction—advection—diffusion system modeling interspecific competition between two species over bounded domains. The kinetic terms are assumed to satisfy the Beddington—DeAngelis functional responses. We consider the situation that first species disperse by a combination of random walk and directed movement along the population density of the second species which disperse randomly within the habitat. For multi—dimensional bounded domains, we prove the global existence and boundedness of time—dependent solutions. For one—dimensional finite domains, we study the effect of diffusion and advection on the existence and stability of nonconstant positive steady states to the strongly coupled elliptic system. In particular, our stability result of these nontrivial steady states provides a selection mechanism for stable wavemodes of the time-dependent system. In the limit of diffusion rates, we show that the steady states of this full elliptic system can be approximated by nonconstant positive solutions of a shadow system and then we construct boundary spike solutions to this shadow system. For the full elliptic system, we also investigate solutions with a single boundary spike or an inverted boundary spike, i.e., the first species concentrate on the boundary point while the second species dominate the whole habitat except the boundary point. These spatial structures

<sup>\*</sup>Corresponding author. Email: qwang@swufe.edu.cn. QW receives supports from NSFC-China (No.11501460), the Project sponsored by SRF for ROCS, SEM and the Project (No.15ZA0382) from Department of Education, Sichuan China.

can be used to model the spatial segregation phenomenon through interspecific competitions. Some numerical studies are performed to illustrate and support our theoretical findings.

2010 Mathematics Subject Classification. Primary: 35B25, 35B35, 35B40, 35A01, 35J47. Secondary: 35B32, 92D25,92D40.

Keywords: Competition system, Beddington–DeAngelis functional response, global solutions, steady–state, stability, boundary spike, boundary layer.

#### 1 Introduction

In this paper, we consider the following reaction-advection-diffusion system

$$\begin{cases}
 u_t = \nabla \cdot (D_1 \nabla u + \chi u \phi(v) \nabla v) + f(u, v), & x \in \Omega, t > 0, \\
 v_t = D_2 \Delta v + g(u, v), & x \in \Omega, t > 0, \\
 \partial_{\mathbf{n}} u = \partial_{\mathbf{n}} v = 0, & x \in \partial \Omega, t > 0, \\
 u(x, 0) = u_0(x), v(x, 0) = v_0(x), & x \in \Omega,
\end{cases}$$
(1.1)

where

$$f(u,v) = \left(-\alpha_1 + \frac{\beta_1}{a_1 + b_1 u + c_1 v}\right) u, \ g(u,v) = \left(-\alpha_2 + \frac{\beta_2}{a_2 + b_2 u + c_2 v}\right) v. \tag{1.2}$$

 $\Omega$  is a bounded smooth domain in  $\mathbb{R}^N$ ,  $N \geq 1$ ,  $\partial \Omega$  is its boundary and  $\mathbf{n}$  denotes the unit outer normal on  $\partial \Omega$ .  $D_i$ ,  $\alpha_i$ ,  $\beta_i$ ,  $a_i$ ,  $b_i$  and  $c_i$ , i = 1, 2, are positive constants.  $\chi$  is a constant and  $\phi$  is a smooth function such that  $\phi(v) > 0$  for all v > 0. The initial conditions  $u_0(x)$  and  $v_0(x)$  are non-negative smooth functions which are assumed to be not identically zero.

System (1.1) describes the evolution of population densities of two mutually interfering species over a bounded habitat  $\Omega$ . u(x,t) and v(x,t) are population densities of the competing species at space—time location  $(x,t) \in \Omega \times \mathbb{R}^+$ . Diffusion rates  $D_1$  and  $D_2$  measure unbiased dispersals of the species. It is assumed that species u senses the population pressure from v and directs its dispersal accordingly. In particular, u moves up or down along the population gradient of interspecies v and  $\chi$  is a constant that measures the intensity of such directed movement.  $\chi < 0$  if u invades the dwelling habitat of v and  $\chi > 0$  if u escapes the habitat of v. Therefore u takes active dispersal strategy to cope with population pressure from v, either to seek or avoid interspecific competition.  $\phi$  reflects the variation of the directed movement strength with respect to population density v.

For almost all mechanistic models that describe population dynamics, functional responses play key roles in the spatial–temporal and qualitative properties of the population distributions. One feature of these kinetics is to take into account the environment's maximal load or carrying capacity. Among the most common types are Lotka–Volterra [45, 7], Hollings [26, 52] and Leslie–Gower [4], etc. To investigate mutual interference among intra–species, Beddington [5] and DeAngelis  $et\ al.$  introduced [14] the following functional response of focal species u preying natural resources

$$f_0(R, u) = \frac{1}{1 + ahR + bu},$$

where R is the resource density, a is the attacking rate and h is the handling time, while b measures the interference rate. To account for both intra– and inter–specific competition, B. de Villemereuil and Lopez-Sepulcre [46] generalized the B-D response into

$$f_0(R, u, v) = \frac{1}{1 + ahR + bu + cv},$$

where b and c represent the intra- and inter–specific competition rates. Moreover, they performed field experiments on guppy-killifish system and collected data that the above models in the absence and presence of inter–species. Functional response  $f_0(R, u, v)$  models a competition relationship between u and v based on the idea that an increase in the population density one species should decreases the growth rate of all individuals since they consume the same resources.

In this paper, we assume that species u and v satisfy the Beddington-DeAngelis functional responses in (1.2). Ecologically,  $\alpha_1$  and  $\alpha_2$  represent intrinsic death rates of species u and v and  $\beta_1$  and  $\beta_2$  interpret their growth rates;  $a_1$  and  $a_2$  represent the compound effects of resource handling time and attack rate, while  $b_1$  and  $c_2$  account for intra-specific competition and  $b_2$  and  $c_1$  are the coefficients of inter-specific competition. We are motivated to investigate the dynamics of population distributions to (1.1) and its stationary system due to the effect of biased movement of species u. To manifest this effect, we assume that the resources are spatially homogeneous and all the parameters are assumed to be positive constants.

To model the coexistence and segregation phenomenon through interspecific competition, various reaction-diffusion systems with advection or cross-diffusion have been proposed and studied. For example, Wang *et al.* investigated in [47] the global existence of (1.1) with Lotka-Volterra kinetics  $f(u,v) = (a_1 - b_1u - c_1v)u$  and  $g(u,v) = (a_2 - b_2u - c_2v)v$ . Existence and

stability of nonconstant positive steady states have also been established through rigour bifurcation analysis. They also studied transition layer steady states to the system. Another example of this type is the SKT model proposed by Shigesada, Kawasaki and Teramoto [42] in 1979 to study the directed dispersals due to mutual interactions. See [30, 31, 36, 39, 42] etc. for works and recent developments on the SKT model. It is also worthwhile to mention that predator—prey models with Beddington—DeAngelis type functional response have been extensively used in [8, 15, 21] etc. Moreover, we refer the reader to [11] and the recent survey paper [10] for detailed discussions on reaction—advection—diffusion systems of population dynamics.

From the viewpoint of mathematical modeling, it is interesting and important to investigate the spatially inhomogeneous distribution of population densities such as the coexistence and segregation of mutually interfering species, in particular, due to the effect of the dispersal strategy and population kinetics. Time-dependent solutions can be used to model the segregation phenomenon in terms of finite or infinite blow-ups, that is, a species population density converges to a  $\delta$ -function or a linear combination of  $\delta$ -functions. This approach has been taken for Keller-Segel chemotaxis system that models the directed cellular movements, along the gradient of chemicals in their environment. See [18, 17, 37]. An alternative approach is to show that the solutions exist globally and converge to bounded steady states. Then positive steady states with concentrating or aggregating structures such as boundary spikes, transition layers, etc. can be used to model the segregation phenomenon. This approach has been taken by [31, 39, 48] etc. The blow-up solution or a  $\delta$ -function is evidently connected to the species segregation phenomenon, however, it is not an optimal choice from the viewpoint of mathematical modeling since it challenges the rationality that population density can not be infinity. On the other hand, it brings challenges to numerical simulations and makes it impossible to analyze the states after blow-up.

The rest of this paper is organized as follows. In Section 2, we obtain the existence and uniform boundedness of positive classical solutions to (1.1) over multi-dimensional bounded domains—See Theorem 2.1. Our proof begins with the local existence and extension theory of Amann [2, 3] for general quasilinear parabolic systems. In Section 3, we study the existence and stability of nonconstant positive steady states of (1.1) over  $\Omega = (0, L)$ . Our method is based on the local theory of Crandall and Rabinowitz [12] and its new version recently developed by Shi and Wang in [41]. Our stability results give a selection mechanism for stable wavemodes of system (1.1)—See Theorem 3.2 and Theorem 3.3. Section 4 is devoted to the existence and

asymptotic behaviors of nonconstant steady states with large amplitude. It is shown that (1.1)–(1.2) admits boundary spike and boundary layer solution if  $D_1$ ,  $\chi$  are comparably large and  $D_2$  is small. In Section 5 we briefly discuss our results and their applications. Some interesting problems are also proposed for future studies.

In the sequel, C and  $C_i$  denote positive constants that may vary from line to line.

# 2 Global existence and boundedness

In this section, we investigate the global existence of positive classical solutions to system (1.1)–(1.2). We will show that the  $L^{\infty}$ -norms of u and v are both uniformly bounded in time. The first set of our main results states as follow.

**Theorem 2.1.** Let  $\Omega \subset \mathbb{R}^N$ ,  $N \geq 1$ , be a bounded domain with smooth boundary  $\partial \Omega$  and assume that  $\phi$  is a smooth function satisfying  $\phi(v) > 0$  for all  $v \geq 0$ . Suppose that  $(u_0, v_0) \in C(\bar{\Omega}) \times W^{1,p}$  for some p > N, and  $u_0, v_0 \geq 0, \not\equiv 0$  on  $\bar{\Omega}$ . Then the IBVP (1.1)-(1.2) admits a unique globally bounded classical solution (u(x,t),v(x,t)); moreover u and v are positive on  $\bar{\Omega}$  for all  $t \in (0,\infty)$ .

Many reaction–diffusion systems of population dynamics have maximum principles, which can be used to prove the global existence and boundedness of their classical solutions. However, the presence of advection term makes system (1.1) non–monotone and it inhibits the application of the comparison principle, at least in proving the boundedness of u. A review of literature suggests that there are two well–established methods to prove the global existence for reaction–advection–diffusion systems. One method is to derive the  $L^{\infty}$  through the  $L^p$ -iterations of some compounded functions of the population densities; another method is to apply standard theory on semigroups generated by  $\{e^{-\Delta t}\}_{t\geq 0}$  and  $L^p-L^q$  estimates on the abstract form of the system. Our proof of global existence to (1.1)–(1.2) involves both techniques.

#### 2.1 Preliminary results and Local existence

First of all, we collect some basic properties of the analytic Neumann semigroup  $\{e^{t\Delta}\}_{t\geq 0}$ . We refer the reader to [16] for classical results and [20, 19, 49] for recent developments. Let  $\Omega \subset \mathbb{R}^N$ ,  $N \geq 1$ , be a bounded domain. It is well known that  $A = -\Delta + 1$  is sectorial in  $L^p(\Omega)$  and it possesses closed fractional powers  $A^{\theta}$ ,  $\theta \in (0, 1)$ . For  $m = 0, 1, p \in [1, \infty]$  and  $q \in (1, \infty)$ , the

domain  $\mathcal{D}(A^{\theta})$  is a Banach space endowed with norm

$$||w||_{\mathcal{D}(A^{\theta})} = ||A^{\theta}w||_{L^{p}(\Omega)};$$

moreover, we have the following embeddings

$$\mathcal{D}(A^{\theta}) \hookrightarrow \begin{cases} W^{m,q}(\Omega), & \text{if } m - N/q < 2\theta - N/p, \\ C^{\delta}(\bar{\Omega}), & \text{if } 2\theta - N/p > \delta \ge 0. \end{cases}$$
 (2.1)

furthermore, for each p > 1,  $\{e^{t\Delta}\}_{t \ge 0}$  maps  $L^p$  into  $\mathcal{D}(A^{\theta})$ : there exists a positive constant C dependent on  $\Omega$ , N and p and  $\nu$  such that, for any  $q \in (p, +\infty]$ 

$$||A^{\theta}e^{-At}w||_{L^{q}(\Omega)} \le Ct^{-\theta - \frac{N}{2}(\frac{1}{p} - \frac{1}{q})}e^{-\nu t}||w||_{L^{p}(\Omega)}, \forall w \in L^{p}(\Omega), \tag{2.2}$$

and

$$||A^{\theta}e^{t\Delta}w||_{L^{q}(\Omega)} \le Ct^{-\theta - \frac{N}{2}(\frac{1}{p} - \frac{1}{q})}e^{-\nu t}||w||_{L^{p}(\Omega)}, \forall w \in L^{p}(\Omega), \int_{\Omega} w = 0,$$
(2.3)

where  $\nu > 0$  is the principal Neumann eigenvalue of  $-\Delta$ .

Next we present the local existence of classical solutions to (1.1)–(1.2) and their extension criterion based on Amann's theory in the following theorem.

Theorem 2.2. Let  $\Omega$  be a bounded domain in  $\mathbb{R}^N$ ,  $N \geq 1$  with smooth boundary  $\partial\Omega$  and assume that  $\phi$  is a continuous function. Then for any initial data  $(u_0, v_0) \in C(\bar{\Omega}) \times W^{1,p}(\Omega)$ , p > N, satisfying  $u_0, v_0 \geq \neq 0$  on  $\bar{\Omega}$ , there exists a constant  $T_{\max} \in (0, \infty]$  and a unique solution (u(x,t),v(x,t)) to (1.1)-(1.2) defined on  $\bar{\Omega} \times [0,T_{\max})$  such that  $(u(\cdot,t),v(\cdot,t)) \in C^0(\bar{\Omega},[0,T_{\max}))$ ,  $u,v \in C^{2,1}(\bar{\Omega},[0,T_{\max}))$ , and u(x,t),v(x,t)>0 on  $\bar{\Omega}$  for all  $t \in (0,T_{\max})$ . Moreover, if  $\sup_{s\in(0,t)}\|(u,v)(\cdot,s)\|_{L^\infty}$  is bounded for  $t \in (0,T_{\max})$ , then  $T_{\max}=\infty$ , i.e., (u,v) is global in time.

*Proof.* Let  $\mathbf{w} = (u, v)$ . (1.1) can be rewritten as

$$\begin{cases}
\mathbf{w}_{t} = \nabla \cdot (A(\mathbf{w})\nabla \mathbf{w}) + F(\mathbf{w}), & x \in \Omega, \ t > 0, \\
\mathbf{w}(x,0) = (u_{0}, v_{0}), & x \in \Omega; \frac{\partial \mathbf{w}}{\partial \mathbf{n}} = 0, \ x \in \partial \Omega, & t > 0,
\end{cases}$$
(2.4)

where

$$A(\mathbf{w}) = \begin{pmatrix} D_1 & \chi u \phi(v) \\ 0 & D_2 \end{pmatrix}, \quad F(\mathbf{w}) = \begin{pmatrix} f(u, v) \\ g(u, v) \end{pmatrix}.$$

(2.4) is a triangular normally parabolic system since the eigenvalues of A are positive, then the existence part follows from Theorem 7.3 and Theorem 9.3 of [2] and the extension criterion follows from Theorem 5.2 in [3]. Moreover, one can apply the standard parabolic boundary  $L^p$  estimates and Schauder estimates to see that  $u_t$ ,  $v_t$  and all the spatial partial derivatives of u and v are bounded in  $\bar{\Omega} \times (0, \infty)$  up to the second order, hence (u, v) has the regularities as stated in the Theorem.

On the other hand, we can use parabolic Strong Maximum Principle and Hopf's boundary point lemma to show that u > 0 and v > 0 on  $\bar{\Omega} \times (0, T_{\text{max}})$ . This completes the proof of Theorem 2.2.

#### 2.2 A prior estimates

We collect some properties of the local classical solutions obtained in Theorem 2.2. First of all, we have the following results.

**Lemma 2.3.** Under the same conditions as in Theorem 2.2, there exists a positive constant  $C_1$  dependent on  $a_1, b_1, \alpha_1, \beta_1, |\Omega|$  and  $||u_0||_{L^1}$  such that

$$||u(\cdot,t)||_{L^1(\Omega)} \le C_1, \forall t \in (0,T_{\text{max}});$$
 (2.5)

moreover, for any  $p \in (1, \infty)$ , there exists a positive constant C(p) dependent on  $a_2, b_2, \alpha_2, \beta_2, |\Omega|$  and  $||v_0||_{L^p}$  such that

$$||v(\cdot,t)||_{L^p(\Omega)} \le C(p), \forall t \in (0, T_{\text{max}}). \tag{2.6}$$

*Proof.* To show (2.5), it is sufficient to show that  $\int_{\Omega} u(x,t)dx$  is uniformly bounded for  $t \in (0,\infty)$  since u(x,t) > 0 according to Theorem 2.2. Integrating the first equation in (1.1) over  $\Omega$  yields

$$\frac{d}{dt} \int_{\Omega} u(x,t) dx = \int_{\Omega} \left( -\alpha_1 + \frac{\beta_1}{a_1 + b_1 u + c_1 v} \right) u dx$$

$$= -\alpha_1 \int_{\Omega} u dx + \int_{\Omega} \frac{\beta_1 u}{a_1 + b_1 u + c_1 v} dx$$

$$\leq -\alpha_1 \int_{\Omega} u dx + \frac{\beta_1 |\Omega|}{b_1}, \tag{2.7}$$

then (2.5) follows from (2.7) thanks to Gronwall's lemma.

To prove (2.6), we first see that  $||v||_{L^1(\Omega)}$  is bounded for  $t \in (0, T_{\text{max}})$  by the same arguments as above. Taking v-equation in (1.1), we have from the integration by parts that

$$\frac{1}{p} \frac{d}{dt} \int_{\Omega} v^{p}(x,t) dx = \int_{\Omega} v^{p-1} v_{t} dx = \int_{\Omega} v^{p-1} \left( D_{2} \Delta v + \left( -\alpha_{2} + \frac{\beta_{2}}{a_{2} + b_{2} u + c_{2} v} \right) v \right) dx 
= -\frac{4D_{2}(p-1)}{p^{2}} \int_{\Omega} |\nabla v^{\frac{p}{2}}|^{2} dx + \int_{\Omega} \left( -\alpha_{2} + \frac{\beta_{2}}{a_{2} + b_{2} u + c_{2} v} \right) v^{p} dx 
\leq -\frac{4D_{2}(p-1)}{p^{2}} \int_{\Omega} |\nabla v^{\frac{p}{2}}|^{2} dx - \alpha_{2} \int_{\Omega} v^{p} dx + \frac{\beta_{2}}{c_{2}} \int_{\Omega} v^{p-1} dx 
\leq -\frac{4D_{2}(p-1)}{p^{2}} \int_{\Omega} |\nabla v^{\frac{p}{2}}|^{2} dx - \frac{\alpha_{2}}{2} \int_{\Omega} v^{p} dx + C_{3}(p), \tag{2.8}$$

therefore,  $||v(\cdot,t)||_{L^p(\Omega)}$  is bounded over  $(0,T_{\max})$  for any  $p \in (1,\infty)$ .

Lemma 2.3 provides the  $L^{\infty}$ -bound of v and  $L^{1}$ -bound of u. To establish the  $L^{\infty}$ -bound on u, we need to estimate  $\|\nabla v\|_{L^{p}}$  for some large p. For this purpose, we convert the v-equation in (1.1) into the following abstract form

$$v(\cdot,t) = e^{-D_2 A t} v_0 + \int_0^t e^{-D_2 A (t-s)} \left( D_2 v(\cdot,s) + g(u(\cdot,s), v(\cdot,s)) \right) ds, \tag{2.9}$$

where g(u, v) is given in (1.2). After applying the estimates (2.1)-(2.3) on (2.9), we have the following result.

**Lemma 2.4.** Let (u,v) be a classical solution of (1.1)–(1.2) over  $\Omega \times (0,T_{\max})$ . For any  $1 \leq p \leq q \leq \infty$ , there exists a positive constant C dependent on  $||v_0||_{L^q(\Omega)}$  and  $\Omega$  such that

$$||v(\cdot,t)||_{W^{1,q}(\Omega)} \le C\left(1 + \int_0^t e^{-\nu(t-s)}(t-s)^{-\frac{1}{2} - \frac{N}{2}(\frac{1}{p} - \frac{1}{q})}||v(\cdot,s) + 1||_{L^p}ds\right),\tag{2.10}$$

for any  $t \in (0, T_{\max})$  provided that  $q \in [1, \frac{Np}{N-p})$  if  $p \in [1, N)$ ,  $q \in [1, \infty)$  if p = N and  $q = \infty$  if p > N, where  $\nu$  is the first Neumann eigenvalue of  $-\Delta$  in  $\Omega$ .

By taking p > N in (2.10), we can quickly have the following result.

**Lemma 2.5.** Assume the same conditions as in Lemma 2.3. Then there exists a positive constant C

$$||v(\cdot,t)||_{W^{1,\infty}(\Omega)} < C, \forall t \in (0,T_{\text{max}}).$$
 (2.11)

#### 2.3 Global existence of bounded classical solutions

Proof of Theorem 2.1. Thanks to (2.11) in Lemma 2.5, we only need to show that  $\sup_{t \in (0,T_{\text{max}})} \|u(\cdot,t)\|_{L^{\infty}} < \infty$ , then we must have that  $T_{\text{max}} = \infty$ ; moreover, the regularities of solutions (u,v) follow from Theorem 2.1.

Without loss of our generality, we assume, in light of Lemma 2.5, that  $\|\phi(v(\cdot,t))\nabla v(\cdot,t)\|_{L^{\infty}(\Omega)} \le 1$  for all  $t \in (0, T_{\text{max}})$ . For any p > 1, we test the first equation of (1.1) by  $pu^{p-1}$  and integrate it over  $\Omega$  by parts to have that

$$\frac{1}{p} \frac{d}{dt} \int_{\Omega} u^{p} dx = \int_{\Omega} u^{p-1} u_{t} dx 
= \int_{\Omega} u^{p-1} \Big( \nabla \cdot (D_{1} \nabla u + \chi u \phi(v) \nabla v) + (-\alpha_{1} + \frac{\beta_{1}}{a_{1} + b_{1} u + c_{1} v}) u \Big) 
\leq -\frac{4D_{1}(p-1)}{p^{2}} \int_{\Omega} |\nabla u^{\frac{p}{2}}|^{2} dx + \frac{2(p-1)\chi}{p} \int_{\Omega} u^{\frac{p}{2}} |\nabla u^{\frac{p}{2}}| dx - \alpha_{1} \int_{\Omega} u^{p} dx + \frac{\beta_{1}}{b_{1}} \int_{\Omega} u^{p-1} dx 
\leq -\frac{2D_{1}(p-1)}{p^{2}} \int_{\Omega} |\nabla u^{\frac{p}{2}}|^{2} dx + \frac{(p-1)\chi^{2}}{2D_{1}} \int_{\Omega} u^{p} dx + C_{1}(p), \tag{2.12}$$

where  $C_1(p) = \frac{1}{q} \left( \frac{\beta_1}{\alpha_1 b_1 p} \right)^{\frac{q}{p}} |\Omega|$  with  $q = \frac{p}{p-1}$  and we have applied the Young's inequality

$$ab \le \epsilon a^p + \frac{b^q}{q(p\epsilon)^{\frac{q}{p}}}, \forall a, b, \epsilon > 0.$$

We recall Corollary 1 in [9] due to the Gagliardo–Nirenberg inequality: for any  $\epsilon > 0$ , there exists a positive constant  $C_0$  dependent on N and  $\Omega$  such that

$$||u^{\frac{p}{2}}||_{L^{2}(\Omega)}^{2} \le \epsilon ||\nabla u^{\frac{p}{2}}||_{L^{2}(\Omega)}^{2} + C_{0}(1 + \epsilon^{-\frac{N}{2}})||u^{\frac{p}{2}}||_{L^{1}(\Omega)}^{2}.$$
(2.13)

Choosing  $\epsilon = \left(\frac{\sqrt{2}D_1}{p\chi}\right)^2$  in (2.13), we see that (2.12) becomes

$$\frac{d}{dt} \int_{\Omega} u^{p} dx \leq -\frac{p(p-1)\chi^{2}}{D_{1}} \int_{\Omega} u^{p} dx + \frac{C_{0}p^{2}\chi^{2}}{2D_{1}^{2}} \left(1 + \left(\frac{p\chi}{\sqrt{D_{1}}}\right)^{N}\right) \left(\int_{\Omega} u^{\frac{p}{2}} dx\right)^{2} dx + C_{1}(p)p \quad (2.14)$$

Finally, by applying the standard Moser–Alikakos iteration [1] to (2.14), we can show that  $||u(\cdot,t)||_{L^{\infty}(\Omega)}$  is uniformly bounded for  $t \in (0,\infty)$ . This completes the proof of Theorem 2.1.  $\square$ 

# 3 Existence of nonconstant positive steady states

In this section, we investigate nonconstant positive steady states of (1.1)–(1.2) over one–dimensional finite domain in the following form

$$\begin{cases}
(D_1 u' + \chi u \phi(v) v')' + \left(-1 + \frac{1}{a_1 + b_1 u + c_1 v}\right) u = 0, & x \in (0, L), \\
D_2 v'' + \left(-1 + \frac{1}{a_2 + b_2 u + c_2 v}\right) v = 0, & x \in (0, L), \\
u'(x) = v'(x) = 0, & x = 0, L,
\end{cases}$$
(3.1)

where ' denotes the derivative taken with respect to x. We have assumed in (3.1) that  $\alpha_i = \beta_i = 1$ , i = 1, 2 without loss of our generality. Indeed, through the scalings

$$\tilde{t} = \alpha_1 t, \tilde{x} = \sqrt{\alpha_1} x, \tilde{c}_1 = \frac{\alpha_2 c_1}{\beta_1}, \tilde{c}_2 = \frac{\alpha_2^2 c_2}{\alpha_1 \beta_2}, \tilde{\chi} = \frac{\chi}{\alpha_1}$$
$$\tilde{D}_i = \frac{D_1}{\alpha_i}, \tilde{a}_i = \frac{a_i \alpha_i}{\beta_i}, i = 1, 2.$$

and  $\tilde{v} = \frac{\alpha_1}{\alpha_2}v$ , after dropping the tildes (1.1)–(1.2) becomes

$$\begin{cases}
 u_t = \nabla \cdot \left(D_1 \nabla u + \chi u \phi(v) \nabla v\right) + \left(-1 + \frac{1}{a_1 + b_1 u + c_1 v}\right) u, & x \in \Omega, t > 0, \\
 v_t = D_2 \Delta v + \left(-1 + \frac{1}{a_2 + b_2 u + c_2 v}\right) v, & x \in \Omega, t > 0, \\
 \partial_{\mathbf{n}} u = \partial_{\mathbf{n}} v = 0, & x \in \partial \Omega, t > 0, \\
 u(x, 0) = u_0(x), v(x, 0) = v_0(x), & x \in \Omega,
\end{cases}$$
(3.2)

Then the one-dimensional steady state of (3.2) over  $\Omega = (0, L)$  leads us to (3.1).

We will see that large advection rate  $\chi$  drives the emergence of nonconstant positive solutions to (3.1). There are four constant solutions to system (3.1): (0,0),  $(0,\frac{1-a_2}{c_2})$ ,  $(\frac{1-a_1}{b_1},0)$  and  $(\bar{u},\bar{v})$ , where

$$\bar{u} = \frac{(1 - a_2)c_1 - (1 - a_1)c_2}{b_2c_1 - b_1c_2}, \bar{v} = \frac{(1 - a_1)b_2 - (1 - a_2)b_1}{b_2c_1 - b_1c_2},$$
(3.3)

is the unique positive solution provided that

$$\frac{c_1}{c_2} < \frac{1 - a_1}{1 - a_2} < \frac{b_1}{b_2}, \quad \text{or} \quad \frac{b_1}{b_2} < \frac{1 - a_1}{1 - a_2} < \frac{c_1}{c_2}.$$
 (3.4)

We assume this condition throughout the rest of our paper. The first inequality in (3.4) implies that  $b_2c_1 < b_1c_2$ , therefore the inter–specific competition is weak compared to the intra-specific

competition. We call this condition the weak competition case. Similarly, we call the latter condition in (3.4) the strong competition case. The same weak and strong competition cases are proposed in the studies of SKT competition models in [42].

#### 3.1 Diffusive system without advection

We shall show that the emergence of nonconstant positive solutions to (3.1) is driven by large advection rate  $\chi$ . To see this, we study the existence of nonconstant positive solutions to (3.2) with  $\chi = 0$ , *i.e.*, the following diffusive system

$$\begin{cases}
 u_t = D_1 u'' + \left(-1 + \frac{1}{a_1 + b_1 u + c_1 v}\right) u, & x \in (0, L), t > 0, \\
 v_t = D_2 v'' + \left(-1 + \frac{1}{a_2 + b_2 u + c_2 v}\right) v, & x \in (0, L), t > 0, \\
 u'(x) = v'(x) = 0, & x = 0, L, t > 0, \\
 u(x, 0) = u_0(x), v(x, 0) = v_0(x), & x \in (0, L).
\end{cases}$$
(3.5)

Our main result states as follows.

**Theorem 3.1.** Suppose that (3.4) holds. Then  $(\bar{u}, \bar{v})$  in (3.3) is asymptotically stable with respect to (3.5) and it is the only positive steady state of (3.5) provided with either (i).  $\frac{b_1}{b_2} > \frac{c_1}{c_2}$  or (ii).  $\frac{b_1}{b_2} \le \frac{c_1}{c_2}$  and  $\max\{D_1, D_2\}$  is large.

Proof. Denote

$$f_0(u,v) = -1 + \frac{1}{a_1 + b_1 u + c_1 v}, g_0(u,v) = -1 + \frac{1}{a_2 + b_2 u + c_2 v},$$

and we rewrite (3.5) as

$$\begin{cases} u_t = D_1 u'' + f_0(u, v)u, & x \in (0, L), t > 0 \\ v_t = D_2 v'' + g_0(u, v)v, & x \in (0, L), t > 0 \\ u'(x) = v'(x) = 0, & x = 0, L, t > 0, \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), & x \in (0, L). \end{cases}$$

$$(3.6)$$

Then the matrix

$$\mathcal{M} = \left(\frac{\partial (f_0, g_0)}{\partial (u, v)}\right)\Big|_{(\bar{u}, \bar{v})} = \begin{pmatrix} \frac{\partial f_0}{\partial u} & \frac{\partial f_0}{\partial v} \\ \frac{\partial g_0}{\partial u} & \frac{\partial g_0}{\partial v} \end{pmatrix}\Big|_{(\bar{u}, \bar{v})} = \begin{pmatrix} -b_1 & -c_1 \\ -b_2 & -c_2 \end{pmatrix},$$

has determinant  $|\mathcal{M}| = (b_1c_2 - b_2c_1)$ . According to Theorem 3.1 of [30],  $(\bar{u}, \bar{v})$  is the only steady state of (3.5) if either (i) or (ii) holds.

On the other hand, it follows from straightforward calculations that the linearized stability matrix corresponding to system (3.5) at  $(\bar{u}, \bar{v})$  is

$$\begin{pmatrix} -D_1 \left(\frac{k\pi}{L}\right)^2 - b_1 \bar{u} & -c_1 \bar{u} \\ -b_2 \bar{v} & -D_2 \left(\frac{k\pi}{L}\right)^2 - c_2 \bar{v} \end{pmatrix},$$

which has two negative eigenvalues if either (i) or (ii) occurs, then by the same analysis in Theorem 2.5 in [32] or [43], one can show that system (3.5) generates a strongly monotone semi-flow on  $C([0,L],\mathbb{R}^2)$  with respect to  $\{(u,v)\in C([0,L],\mathbb{R}^2)|u>0,v>0\}$ , hence  $(\bar{u},\bar{v})$  is globally asymptotically stable. This completes the proof of Theorem 3.1.

Our results indicate that the global dynamics of the diffusion system (3.5) is dominated by the ODEs in the weak competition case and also in the strong case if one of the diffusion rates is large. However, if both  $D_1$  and  $D_2$  are small,  $(\bar{u}, \bar{v})$  is unstable in the strong competition case. We surmise that positive solutions with nontrivial patterns may arise when the system parameters and the domain geometry are properly balanced. Nonconstant positive solutions with spikes are investigated for the diffusion system (3.1) with Lotka–Volterra dynamics by various authors. See [23, 33, 34, 35].

#### 3.2 Advection–driven instability

Theorem 3.1 states that diffusion does not change the dynamics of the spatially homogeneous solution and no Turing's instability occurs for the diffusive system (3.5) in most cases. We proceed to investigate the effect of advection  $\chi$  on the emergence of nonconstant positive solutions to (3.1). We shall show that this equilibrium loses its stability as the advection rate  $\chi$  crosses a threshold value. To this end, we first study the linearized stability of the equilibrium  $(\bar{u}, \bar{v})$ . Let  $(u, v) = (\bar{u}, \bar{v}) + (U, V)$ , where U and V are small perturbations from  $(\bar{u}, \bar{v})$ , then

$$\begin{cases}
U_{t} \approx D_{1}U'' + \chi \bar{u}\phi(\bar{v})V'' - b_{1}\bar{u}U - c_{1}\bar{u}V, & x \in (0, L), t > 0, \\
V_{t} \approx D_{2}V'' - b_{2}\bar{v}U - c_{2}\bar{v}V, & x \in (0, L), t > 0, \\
U'(x) = V'(x) = 0, & x = 0, L, t > 0.
\end{cases}$$
(3.7)

We have the following result on the linearized instability of  $(\bar{u}, \bar{v})$  to (3.1).

**Proposition 1.** The constant solution  $(\bar{u}, \bar{v})$  of (3.1) is unstable if

$$\chi > \chi_0 = \min_{k \in \mathbb{N}^+} \frac{\left(D_1(\frac{k\pi}{L})^2 + b_1 \bar{u}\right) \left(D_2(\frac{k\pi}{L})^2 + c_2 \bar{v}\right) - b_2 c_1 \bar{u}\bar{v}}{b_2(\frac{k\pi}{L})^2 \phi(\bar{v}) \bar{u}\bar{v}}.$$
 (3.8)

*Proof.* According to the standard linearized stability analysis, the stability of  $(\bar{u}, \bar{v})$  is determined by the eigenvalues of the following matrix

$$\mathcal{H}_{k} = \begin{pmatrix} -D_{1}(\frac{k\pi}{L})^{2} - b_{1}\bar{u} & -\chi\bar{u}\phi(\bar{v})(\frac{k\pi}{L})^{2} - c_{1}\bar{u} \\ -b_{2}\bar{v} & -D_{2}(\frac{k\pi}{L})^{2} - c_{2}\bar{v} \end{pmatrix}, k \in \mathbb{N}^{+}.$$
(3.9)

In particular,  $(\bar{u}, \bar{v})$  is unstable if  $\mathcal{H}_k$  has an eigenvalue with positive real part for some  $k \in \mathbb{N}^+$ . It is easy to see that the characteristic polynomial of (3.9) takes the form

$$p_k(\lambda) = \lambda^2 + T_k \lambda + D_k,$$

where

$$T_k = (D_1 + D_2)(\frac{k\pi}{L})^2 + b_1\bar{u} + c_2\bar{v} > 0,$$

and

$$D_k = \left(D_1(\frac{k\pi}{L})^2 + b_1\bar{u}\right)\left(D_2(\frac{k\pi}{L})^2 + c_2\bar{v}\right) - \left(\chi\bar{u}\phi(\bar{v})(\frac{k\pi}{L})^2 + c_1\bar{u}\right)b_2\bar{v}.$$

 $p_k(\lambda)$  has a positive root if and only if  $p_k(0) = D_k < 0$ , then (3.8) follows from simple calculations and the proof completes.

We have from Proposition 1 that  $(\bar{u}, \bar{v})$  loses its stability when  $\chi$  surpasses  $\chi_0$ . In the weak competition case  $\frac{b_1}{b_2} > \frac{1-a_1}{1-a_2} > \frac{c_1}{c_2}$ , we see that  $\chi_0$  is always positive, hence  $(\bar{u}, \bar{v})$  remains locally stable for  $\chi$  being small. In the strong competition case  $\frac{b_1}{b_2} < \frac{1-a_1}{1-a_2} < \frac{c_1}{c_2}$ ,  $\chi_0 < 0$  if both  $D_1$  and  $D_2$  are sufficiently small. This corresponds to the fact that  $(\bar{u}, \bar{v})$  is unstable for  $\chi = 0$  in (3.1) in this case. By the same stability analysis above, we can show that the appearance of advection rate  $\chi$  does not change the stability of the rest equilibrium points. It is also worthwhile to point out that Proposition 1 carries over to higher dimensions with  $(\frac{k\pi}{L})^2$  replaced by the k-eigenvalue of Neumann Laplacian.

#### 3.3 Steady state bifurcation

To establish the existence of nonconstant positive solutions to (3.1), we shall use the bifurcation theory due to Crandall–Rabinowitz [12] by taking  $\chi$  as the bifurcation parameter. To this end, we rewrite (3.1) into the following abstract form

$$\mathcal{F}(u, v, \chi) = 0, \ (u, v, \chi) \in \mathcal{X} \times \mathcal{X} \times \mathbb{R},$$

where

$$\mathcal{F}(u,v,\chi) = \begin{pmatrix} (D_1 u' + \chi u \phi(v)v')' + (-1 + \frac{1}{a_1 + b_1 u + c_1 v})u \\ D_2 v'' + (-1 + \frac{1}{a_2 + b_2 u + c_2 v})v \end{pmatrix}, \tag{3.10}$$

and  $\mathcal{X} = \{H^2(0,L)|u'(0) = u'(L) = 0\}$ . We collect some facts about  $\mathcal{F}$ . First of all,  $\mathcal{F}(\bar{u},\bar{v},\chi) = 0$  for any  $\chi \in \mathbb{R}$  and  $\mathcal{F} : \mathcal{X} \times \mathbb{R} \times \mathbb{R} \to \mathcal{Y} \times \mathcal{Y}$  is analytic for  $\mathcal{Y} = L^2(0,L)$ . For any fixed  $(u_0, v_0) \in \mathcal{X} \times \mathcal{X}$ , the Fréchet derivative of  $\mathcal{F}$  is

$$D_{(u,v)}\mathcal{F}(u_0, v_0, \chi)(u, v) = \begin{pmatrix} D_1 u'' + \chi \left( (\phi(v_0)u + u_0 \phi'(v_0)v)v'_0 + u_0 \phi(v_0)v' \right)' + R \\ D_2 v'' - \frac{b_2 v_0}{(a_2 + b_2 u_0 + c_2 v_0)^2} u + \left( -1 + \frac{a_2 + b_2 u_0}{(a_2 + b_2 u_0 + c_2 v_0)^2} \right) v \end{pmatrix}, \quad (3.11)$$

where  $R = \left(-1 + \frac{a_1 + c_1 v_0}{(a_1 + b_1 u_0 + c_1 v_0)^2}\right) u - \frac{c_1 u_0}{(a_1 + b_1 u_0 + c_1 v_0)^2} v$ . By the same arguments that lead to (iv) of Lemma 5.1 in [9] or Lemma 2.3 of [48], one can show that  $D_{(u,v)} \mathcal{F}(u_0, v_0, \chi)$  is Fredholm with zero index.

For bifurcation to occur at  $(\bar{u}, \bar{v}, \chi)$ , we need the Implicit Function Theorem to fail on  $\mathcal{F}$  at this point, hence we require the following necessary condition  $\mathcal{N}(D_{(u,v)}\mathcal{F}(\bar{u}, \bar{v}, \chi)) \neq 0$ . Let (u, v) be a nontrivial solution in this null–space, then it satisfies the following system

$$\begin{cases}
D_1 u'' + \chi \bar{u}\phi(\bar{v})v'' - b_1 \bar{u}u - c_1 \bar{u}v = 0, & x \in (0, L), \\
D_2 v'' - b_2 \bar{v}u - c_2 \bar{v}v = 0, & x \in (0, L), \\
u'(x) = v'(x) = 0, & x = 0, L.
\end{cases}$$
(3.12)

Expanding u and v into the following series

$$u(x) = \sum_{k=0}^{\infty} t_k \cos \frac{k\pi x}{L}, v(x) = \sum_{k=0}^{\infty} s_k \cos \frac{k\pi x}{L},$$

and substituting them into (3.12) yield

$$\begin{pmatrix}
-D_1(\frac{k\pi}{L})^2 - b_1\bar{u} & -\chi\bar{u}\phi(\bar{v})(\frac{k\pi}{L})^2 - c_1\bar{u} \\
-b_2\bar{v} & -D_2(\frac{k\pi}{L})^2 - c_2\bar{v}
\end{pmatrix}
\begin{pmatrix}
t_k \\
s_k
\end{pmatrix} = \begin{pmatrix}
0 \\
0
\end{pmatrix}.$$
(3.13)

k=0 is ruled out in (3.12) thanks to (3.4). For  $k \in \mathbb{N}^+$ , (3.13) has nonzero solutions if and only if its coefficient matrix is singular which implies that local bifurcation might occur at

$$\chi = \chi_k = \frac{\left(D_1(\frac{k\pi}{L})^2 + b_1\bar{u}\right)\left(D_2(\frac{k\pi}{L})^2 + c_2\bar{v}\right) - b_2c_1\bar{u}\bar{v}}{b_2(\frac{k\pi}{L})^2\phi(\bar{v})\bar{u}\bar{v}}, k \in \mathbb{N}^+.$$
(3.14)

Moreover, the null space is one-dimensional and has a span

$$\mathcal{N}(D_{(u,v)}\mathcal{F}(\bar{u},\bar{v},\chi_k)) = \operatorname{span}\{(\bar{u}_k,\bar{v}_k)\},\$$

where

$$(\bar{u}_k, \bar{v}_k) = (Q_k, 1)\cos\frac{k\pi x}{L}, Q_k = -\frac{D_2(\frac{k\pi}{L})^2 + c_2\bar{v}}{b_2\bar{v}}.$$
 (3.15)

Having the candidates for bifurcation values  $\chi_k$ , we now show that local bifurcation does occur at  $(\bar{u}, \bar{v}, \chi_k)$  in the following theorem, which establishes nonconstant positive solutions to (3.1).

**Theorem 3.2.** Suppose that  $\phi \in C^2(\mathbb{R}, \mathbb{R})$  and  $\phi(v) > 0$  for all v > 0. Assume that (3.4) holds, and for all positive different integers  $k, j \in \mathbb{N}^+$ ,

$$(b_1c_2 - b_2c_1)\bar{u}\bar{v} \neq k^2j^2D_1D_2(\frac{\pi}{L})^4, k \neq j,$$
(3.16)

where  $(\bar{u}, \bar{v})$  is the positive equilibrium of (3.1) given in (3.3). Then for each  $k \in \mathbb{N}^+$ , there exists  $\delta > 0$  small such that (3.1) admits nonconstant solutions  $(u_k(s, x), v_k(s, x), \chi_k(s)) \in \mathcal{X} \times \mathcal{X} \times \mathbb{R}^+$  with  $(u_k(0, x), v_k(0, x), \chi_k(0)) = (\bar{u}, \bar{v}, \chi_k)$ . The solutions are continuous functions of s in the topology of  $\mathcal{X} \times \mathcal{X} \times \mathbb{R}$  and have the following expansions for |s| being small,

$$(u_k(s,x), v_k(s,x)) = (\bar{u}, \bar{v}) + s(Q_k, 1)\cos\frac{k\pi x}{L} + o(s),$$
(3.17)

and  $(u_k(s,x), v_k(s,x)) - (\bar{u},\bar{v}) - s(Q_k,1)\cos\frac{k\pi x}{L} \in \mathcal{Z}$  where

$$\mathcal{Z} = \{(u, v) \in \mathcal{X} \times \mathcal{X} | \int_0^L u\bar{u}_k + v\bar{v}_k dx = 0 \}.$$
 (3.18)

with  $(\bar{u}_k, \bar{v}_k)$  and  $Q_k$  defined in (3.15); moreover, all nontrivial solutions of (3.1) around  $(\bar{u}, \bar{v}, \chi_k)$  must stay on the curve  $\Gamma_k(s) = (u_k(s, x), v_k(s, x), \chi_k(s)), s \in (-\delta, \delta)$ .

*Proof.* Our results follow from Theorem 1.7 of Crandall and Rabinowitz [12] once we prove the following transversality condition,

$$\frac{d}{d\chi} \left( D_{(u,v)} \mathcal{F}(\bar{u}, \bar{v}, \chi) \right) (\bar{u}_k, \bar{v}_k)|_{\chi = \chi_k} \notin \mathcal{R}(D_{(u,v)} \mathcal{F}(\bar{u}, \bar{v}, \chi_k)), \tag{3.19}$$

where  $(\bar{u}_k, \bar{v}_k)$  is given in (3.15) and  $\mathcal{R}(\cdot)$  denotes range of the operator. We argue by contradiction and assume that (3.19) fails. Therefore there exist a nontrivial pair (u, v) to the following problem

$$\begin{cases}
D_1 u'' + \chi_k \bar{u}\phi(\bar{v})v'' - b_1 \bar{u}u - c_1 \bar{u}v = -(\frac{k\pi}{L})^2 \bar{u}\phi(\bar{v})\cos\frac{k\pi x}{L}, & x \in (0, L), \\
D_2 v'' - b_2 \bar{v}u - c_2 \bar{v}v = 0, & x \in (0, L), \\
u'(x) = v'(x) = 0, & x = 0, L.
\end{cases}$$
(3.20)

Testing the first two equations in (3.20) by  $\cos \frac{k\pi x}{L}$  over (0, L) yields

$$\begin{pmatrix} -D_{1}(\frac{k\pi}{L})^{2} - b_{1}\bar{u} & -\chi_{k}\bar{u}\phi(\bar{v})(\frac{k\pi}{L})^{2} - c_{1}\bar{u} \\ -b_{2}\bar{v} & -D_{2}(\frac{k\pi}{L})^{2} - c_{2}\bar{v} \end{pmatrix} \begin{pmatrix} \int_{0}^{L}u\cos\frac{k\pi x}{L}dx \\ \int_{0}^{L}v\cos\frac{k\pi x}{L}dx \end{pmatrix} = \begin{pmatrix} -\frac{(k\pi)^{2}\bar{u}\phi(\bar{v})}{2L} \\ 0 \end{pmatrix}$$
(3.21)

The coefficient matrix is singular due to (3.14), then we reach a contradiction in (3.21) and this proves (3.19). On the other hand, we need condition (3.16) such that  $\chi_k \neq \chi_j$  for all integers  $k \neq j$ , which is also required for the application of the local theory in [12]. The proof of Theorem 3.2 is complete.

## 3.4 Stability analysis of the nonconstant steady states

We proceed to study stabilities of the nontrivial bifurcating solutions  $(u_k(s, x), v_k(s, x), \chi_k(s))$  obtained in Theorem 3.2. Here the stability or instability means that of the nonconstant solution

viewed as an equilibrium of the time-dependent system of (3.1).  $\mathcal{F}$  is  $C^4$ -smooth in s if  $\phi$  is  $C^4$ , therefore, according to Theorem 1.18 in [12],  $(u_k(s,x), v_k(s,x), \chi_k(s))$  is  $C^3$ -smooth, we have the following expansions,

$$\begin{cases} u_k(s,x) = \bar{u} + sQ_k \cos\frac{k\pi x}{L} + s^2\psi_1(x) + s^3\psi_2(x) + o(s^3), \\ v_k(s,x) = \bar{v} + s\cos\frac{k\pi x}{L} + s^2\varphi_1(x) + s^3\varphi_2(x) + o(s^3), \\ \chi_k(s) = \chi_k + sK_1 + s^2K_2 + o(s^2). \end{cases}$$
(3.22)

where  $(\psi_i, \varphi_i) \in \mathcal{Z}$  as defined in (3.18) and  $K_i$  is a constant for i = 1, 2. Moreover, we have from Taylor's Theorem that

$$\phi(v_k(s,x)) = \phi(\bar{v}) + s\phi'(\bar{v})\cos\frac{k\pi x}{L} + s^2(\phi'(\bar{v})\varphi_1 + \frac{1}{2}\phi''(\bar{v})\cos^2\frac{k\pi x}{L}) + o(s^3).$$
 (3.23)

 $o(s^3)$ -terms in (3.22) and (3.23) are taken in  $H^2$ -topology. For the sake of simplicity, we introduce the following notations

$$\begin{split} P_1 &= \frac{1}{2} \Big( f_{uu}(\bar{u}, \bar{v}) Q_k^2 + 2 f_{uv}(\bar{u}, \bar{v}) Q_k + f_{vv}(\bar{u}, \bar{v}) \Big) \\ &= b_1 (b_1 \bar{u} - 1) Q_k^2 + c_1 (2b_1 \bar{u} - 1) Q_k + c_1^2 \bar{u} \\ P_2 &= \frac{1}{2} \Big( g_{uu}(\bar{u}, \bar{v}) Q_k^2 + 2 g_{uv}(\bar{u}, \bar{v}) Q_k + g_{vv}(\bar{u}, \bar{v}) \Big) \\ &= b_2^2 \bar{v} Q_k^2 + b_2 (2c_2 \bar{v} - 1) Q_k + c_2 (c_2 \bar{v} - 1) \\ P_3 &= \frac{1}{6} \Big( f_{uuu}(\bar{u}, \bar{v}) Q_k^3 + 3 f_{uuv}(\bar{u}, \bar{v}) Q_k^2 + 3 f_{uvv}(\bar{u}, \bar{v}) Q_k + f_{vvv}(\bar{u}, \bar{v}) \Big) \\ &= b_1^2 (1 - b_1 \bar{u}) Q_k^3 + b_1 c_1 (2 - 3b_1 \bar{u}) Q_k^2 + c_1^2 (1 - 3b_1 \bar{u}) Q_k - c_1^3 \bar{u} \\ P_4 &= \frac{1}{6} \Big( g_{uuu}(\bar{u}, \bar{v}) Q_k^3 + 3 g_{uuv}(\bar{u}, \bar{v}) Q_k^2 + 3 g_{uvv}(\bar{u}, \bar{v}) Q_k + g_{vvv}(\bar{u}, \bar{v}) \Big) \\ &= - b_2^3 \bar{v} Q_k^3 + b_2^2 (1 - 3c_2 \bar{v}) Q_k^2 + b_2 c_2 (2 - 3c_2 \bar{v}) Q_k + c_2^2 (1 - c_2 \bar{v}) \\ P_5 &= f_{uu}(\bar{u}, \bar{v}) Q_k + f_{uv}(\bar{u}, \bar{v}) = 2b_1 (b_1 \bar{u} - 1) Q_k + c_1 (2b_1 \bar{u} - 1) \\ P_6 &= f_{uv}(\bar{u}, \bar{v}) Q_k + f_{vv}(\bar{u}, \bar{v}) = c_1 (2b_1 \bar{u} - 1) Q_k + 2c_1^2 \bar{u} \\ P_7 &= g_{uu}(\bar{u}, \bar{v}) Q_k + g_{uv}(\bar{u}, \bar{v}) = 2b_2^2 \bar{v} Q_k + b_2 (2c_2 \bar{v} - 1) \\ P_8 &= g_{uv}(\bar{u}, \bar{v}) Q_k + g_{vv}(\bar{u}, \bar{v}) = b_2 (2c_2 \bar{v} - 1) Q_k + 2c_2 (c_2 \bar{v} - 1). \end{aligned}$$

Substituting (3.22) and (3.23) into (3.1) and equating the  $s^2$ -terms, we collect

$$\begin{cases}
D_{1}\psi_{1}^{"} + \chi_{k}\bar{u}\phi(\bar{v})\varphi_{1}^{"} - b_{1}\bar{u}\psi_{1} - c_{1}\bar{u}\varphi_{1} - K_{1}\bar{u}\phi(\bar{v})(\frac{k\pi}{L})^{2}\cos\frac{k\pi x}{L} \\
= \chi_{k}(\frac{k\pi}{L})^{2}(\phi(\bar{v})Q_{k} + \bar{u}\phi'(\bar{v}))\cos\frac{2k\pi x}{L} - P_{1}\cos^{2}\frac{k\pi x}{L}, \\
D_{2}\varphi_{1}^{"} - b_{2}\bar{v}\psi_{1} - c_{2}\bar{v}\varphi_{1} = -P_{2}\cos^{2}\frac{k\pi x}{L}, \\
\psi_{1}^{\prime}(0) = \phi_{1}^{\prime}(0) = 0, \psi_{1}^{\prime}(L) = \phi_{1}^{\prime}(L) = 0.
\end{cases} (3.25)$$

Multiplying the first equation of (3.25) by  $\cos \frac{k\pi x}{L}$  and integrating it over (0,L) by parts yield

$$\frac{(k\pi)^2 \bar{u}\phi(\bar{v})K_1}{2L} = -\left(\chi_k \bar{u}\phi(\bar{v})(\frac{k\pi}{L})^2 + c_1\bar{u}\right) \int_0^L \varphi_1 \cos\frac{k\pi x}{L} dx 
-\left(D_1(\frac{k\pi}{L})^2 + b_1\bar{u}\right) \int_0^L \psi_1 \cos\frac{k\pi x}{L} dx.$$
(3.26)

Multiplying the second equation by  $\cos \frac{k\pi x}{L}$  and integrating it over (0,L) by parts yield

$$b_2 \bar{v} \int_0^L \psi_1 \cos \frac{k\pi x}{L} dx + \left( D_2 (\frac{k\pi}{L})^2 + c_2 \bar{v} \right) \int_0^L \varphi_1 \cos \frac{k\pi x}{L} dx = 0.$$
 (3.27)

On the other hand, (3.18) and the fact that  $(\psi_1, \varphi_1) \in \mathcal{Z}$  give us

$$Q_k \int_0^L \psi_1 \cos \frac{k\pi x}{L} dx + \int_0^L \varphi_1 \cos \frac{k\pi x}{L} dx = 0, \tag{3.28}$$

where  $Q_k = -\frac{D_2(\frac{k\pi}{L})^2 + c_2\bar{v}}{b_2\bar{v}}$ . Solving (3.27) and (3.28) leads us to

$$(1 + Q_k^2) \int_0^L \psi_1 \cos \frac{k\pi x}{L} dx = 0,$$

which implies that

$$\int_0^L \psi_1 \cos \frac{k\pi x}{L} dx = \int_0^L \varphi_1 \cos \frac{k\pi x}{L} dx = 0, \ \forall k \in \mathbb{N}^+.$$

It follows from (3.26) that  $K_1 = 0$ , hence the bifurcation branch  $\Gamma_k(s)$  around  $\chi_k$  is of pitch-fork type, i.e., one-sided. We proceed to evaluate  $K_2$  which determines branch direction hence the stability of  $(u_k(s, x), v_k(s, x), \chi_k(s))$  as we shall see in the coming analysis.

Equating the  $s^3$ -terms in (3.1), we collect

$$\begin{cases}
D_{1}\psi_{2}^{"} + \chi_{k}\bar{u}\phi(\bar{v})\varphi_{2}^{"} - b_{1}\bar{u}\psi_{2} - c_{1}\bar{u}\varphi_{2} - K_{2}\bar{u}\phi(\bar{v})(\frac{k\pi}{L})^{2}\cos\frac{k\pi x}{L} \\
= \chi_{k}A_{0} - P_{5}\psi_{1}\cos\frac{k\pi x}{L} - P_{6}\varphi_{1}\cos\frac{k\pi x}{L} - P_{3}\cos^{3}\frac{k\pi x}{L}, \\
D_{2}\varphi_{2}^{"} - b_{2}\bar{v}\psi_{2} - c_{2}\bar{v}\varphi_{2} = -P_{7}\psi_{1}\cos\frac{k\pi x}{L} - P_{8}\varphi_{1}\cos\frac{k\pi x}{L} - P_{4}\cos^{3}\frac{k\pi x}{L}, \\
\psi_{2}^{\prime}(0) = \varphi_{2}^{\prime}(0) = 0, \psi_{2}^{\prime}(L) = \varphi_{2}^{\prime}(L) = 0,
\end{cases} (3.29)$$

where

$$A_{0} = -\left(\phi(\bar{v})Q_{k} + \bar{u}\phi^{'}(\bar{v})\right)\varphi_{1}^{"}\cos\frac{k\pi x}{L} + \left(\frac{k\pi}{L}\right)^{2}(\phi(\bar{v})\psi_{1} + \bar{u}\phi^{'}(\bar{v})\varphi_{1})\cos\frac{k\pi x}{L}$$

$$+\left(\frac{k\pi}{L}\right)\left(\phi(\bar{v})\psi_{1}^{'} + (\phi(\bar{v})Q_{k} + 2\bar{u}\phi^{'}(\bar{v}))\varphi_{1}^{'}\right)\sin\frac{k\pi x}{L}$$

$$-\left(\frac{k\pi}{L}\right)^{2}\left(2\phi^{'}(\bar{v})Q_{k} + \bar{u}\phi^{''}(\bar{v})\right)\cos\frac{k\pi x}{L}$$

$$+3\left(\frac{k\pi}{L}\right)^{2}\left(\phi^{'}(\bar{v})Q_{k} + \frac{1}{2}\bar{u}\phi^{''}(\bar{v})\right)\cos^{3}\frac{k\pi x}{L}.$$

Testing the first equation in (3.29) by  $\cos \frac{k\pi x}{L}$  over (0, L), we conclude from straightforward calculations that

$$\frac{\bar{u}\phi(\bar{v})(k\pi)^{2}}{L}K_{2} = -2\left(D_{1}\left(\frac{k\pi}{L}\right)^{2} + b_{1}\bar{u}\right) \int_{0}^{L} \psi_{2}\cos\frac{k\pi x}{L}dx 
-2\left(\chi_{k}\bar{u}\phi(\bar{v})\left(\frac{k\pi}{L}\right)^{2} + c_{1}\bar{u}\right) \int_{0}^{L} \varphi_{2}\cos\frac{k\pi x}{L}dx 
+ \left(P_{5} - \chi_{k}\phi(\bar{v})\left(\frac{k\pi}{L}\right)^{2}\right) \int_{0}^{L} \psi_{1}dx 
+ \left(P_{6} - \chi_{k}\bar{u}\phi'(\bar{v})\left(\frac{k\pi}{L}\right)^{2}\right) \int_{0}^{L} \varphi_{1}dx 
+ \left(P_{5} + \chi_{k}\phi(\bar{v})\left(\frac{k\pi}{L}\right)^{2}\right) \int_{0}^{L} \psi_{1}\cos\frac{2k\pi x}{L}dx 
+ \left(P_{6} - \chi_{k}\left(2\phi(\bar{v})Q_{k} + \bar{u}\phi'(\bar{v})\right)\left(\frac{k\pi}{L}\right)^{2}\right) \int_{0}^{L} \varphi_{1}\cos\frac{2k\pi x}{L}dx 
+ \frac{3LP_{3}}{4} - \left(\phi'(\bar{v})Q_{k} + \frac{1}{2}\bar{u}\phi''(\bar{v})\right)\frac{k^{2}\pi^{2}\chi_{k}}{4L},$$
(3.30)

where  $P_3$ ,  $P_5$  and  $P_6$  are given in (3.24), and  $\chi_k$  is defined by (3.14).

On the other hand, testing the second equation of (3.29) by  $\cos \frac{k\pi x}{L}$  over (0, L) gives rise to

$$b_{2}\bar{v}\int_{0}^{L}\psi_{2}\cos\frac{k\pi x}{L}dx + \left(D_{2}(\frac{k\pi}{L})^{2} + c_{2}\bar{v}\right)\int_{0}^{L}\varphi_{2}\cos\frac{k\pi x}{L}dx$$

$$= \frac{P_{7}}{2}\left(\int_{0}^{L}\psi_{1}dx + \int_{0}^{L}\psi_{1}\cos\frac{k\pi x}{L}dx\right) + \frac{P_{8}}{2}\left(\int_{0}^{L}\varphi_{1}dx + \int_{0}^{L}\varphi_{1}\cos\frac{k\pi x}{L}dx\right) - \frac{3P_{4}L}{8},$$
(3.31)

Since  $(\psi_2, \varphi_2) \in \mathcal{Z}$ , we can solve (3.31) to find that

$$\int_{0}^{L} \psi_{2} \cos \frac{k\pi x}{L} dx = \frac{\left(\int_{0}^{L} \psi_{1} \cos \frac{2k\pi x}{L} dx + \int_{0}^{L} \psi_{1} dx\right) P_{7}}{2b_{2} \bar{v}(Q_{k}^{2} + 1)} + \frac{\left(\int_{0}^{L} \varphi_{1} \cos \frac{2k\pi x}{L} dx + \int_{0}^{L} \varphi_{1} dx\right) P_{8}}{2b_{2} \bar{v}(Q_{k}^{2} + 1)} + \frac{3P_{4}L}{8b_{2} \bar{v}(Q_{k}^{2} + 1)},$$
(3.32)

$$\int_{0}^{L} \varphi_{2} \cos \frac{k\pi x}{L} dx = -\frac{Q_{k} \left(\int_{0}^{L} \psi_{1} \cos \frac{2k\pi x}{L} dx + \int_{0}^{L} \psi_{1} dx\right) P_{7}}{2b_{2} \bar{v}(Q_{k}^{2} + 1)} - \frac{Q_{k} \left(\int_{0}^{L} \varphi_{1} \cos \frac{2k\pi x}{L} dx + \int_{0}^{L} \varphi_{1} dx\right) P_{8}}{2b_{2} \bar{v}(Q_{k}^{2} + 1)} - \frac{3Q_{k} P_{4} L}{8b_{2} \bar{v}(Q_{k}^{2} + 1)}.$$
(3.33)

In order to find  $K_2$  in (3.30), we see from (3.32) and (3.33) that it is necessary to evaluate the following integrals

$$\int_0^L \psi_1 dx, \quad \int_0^L \varphi_1 dx, \quad \int_0^L \psi_1 \cos \frac{2k\pi x}{L} dx, \text{ and } \int_0^L \varphi_1 \cos \frac{2k\pi x}{L} dx.$$

Integrating both equations in (3.25) over (0, L), we obtain from straightforward calculations by taking the fact  $K_1 = 0$  that

$$\int_{0}^{L} \psi_{1} dx = \frac{L(c_{2}\bar{v}P_{1} - c_{1}\bar{u}P_{2})}{2(b_{1}c_{2} - b_{2}c_{1})\bar{u}\bar{v}}, \int_{0}^{L} \varphi_{1} dx = \frac{L(b_{1}\bar{u}P_{2} - b_{2}\bar{v}P_{1})}{2(b_{1}c_{2} - b_{2}c_{1})\bar{u}\bar{v}}$$
(3.34)

On the other hand, we multiply both two equations in (3.25) by  $\cos \frac{2k\pi x}{L}$  and integrate them over (0, L) by parts. Then again thanks to  $K_1 = 0$ , we have from straightforward calculations that

$$\int_{0}^{L} \psi_{1} \cos \frac{2k\pi x}{L} dx = \frac{|\mathcal{A}_{1}|}{|\mathcal{A}|}, \quad \int_{0}^{L} \psi_{1} \cos \frac{2k\pi x}{L} dx = \frac{|\mathcal{A}_{2}|}{|\mathcal{A}|}.$$
 (3.35)

where

$$|\mathcal{A}| = 12D_1 D_2 (\frac{k\pi}{L})^4 - 3(b_1 c_2 - b_2 c_1) \bar{u}\bar{v},$$

$$|\mathcal{A}_1| = -2\chi_k D_2 L (\frac{k\pi}{L})^4 (\phi(\bar{v})Q_k + \bar{u}\phi'(\bar{v})) + \frac{L}{4} (c_2 \bar{v}P_1 - c_1 \bar{u}P_2)$$

$$+ \frac{k^2 \pi^2}{L} \Big( D_2 P_1 - \chi_k \bar{u}\phi(\bar{v})P_2 - \frac{1}{2} \chi_k c_2 \bar{v} \big(\phi(\bar{v})Q_k + \bar{u}\phi'(\bar{v}) \big) \Big),$$

and

$$|\mathcal{A}_{2}| = \frac{k^{2}\pi^{2}}{L} \Big( D_{1}P_{2} + \chi_{k}b_{2}\bar{v} \big( \phi(\bar{v})Q_{k} + \bar{u}\phi'(\bar{v}) \big) \Big) + \frac{L}{4} (b_{1}\bar{u}P_{2} - b_{2}\bar{v}P_{1}).$$

Finally, we are able to evaluate  $K_2$  in terms of system parameters, thanks to (3.30), (3.34) and (3.35). The rest calculations are straightforward but tedious and we skip them here. We present the stability of the bifurcating solutions in the following Theorem.

**Theorem 3.3.** Assume the same conditions as in Theorem 3.2. Let  $K_2$  be given in (3.30). Suppose that  $\chi_{k_0} = \min_{k \in \mathbb{N}^+} \chi_k$  of (3.11). Then for all positive integers  $k \neq k_0$  and small  $\delta > 0$ ,  $(u_k(s,x), v_k(s,x))$  is unstable for  $s \in (-\delta, \delta)$ ; moreover,  $(u_{k_0}(s,x), v_{k_0}(s,x))$  is stable for  $s \in (-\delta, \delta)$  if  $K_2 > 0$  and it is unstable for  $s \in (-\delta, \delta)$  if  $K_2 < 0$ .

**Remark 1.** Theorem 3.3 provides a rigourous selection mechanism for stable nonconstant positive solutions to system (3.1). If  $(u_k(s,x),v_k(s,x))$  is stable, then k must be the integer that minimizes the bifurcation value  $\chi_k$  over  $\mathbb{N}^+$ , that being said, if a bifurcation branch is stable, then it must be the first branch counting from the left to the right.

*Proof.* For each  $k \in \mathbb{N}^+$ , we linearize (3.1) around  $(u_k(s,x), v_k(s,x), \chi_k(s))$  and obtain the following eigenvalue problem

$$D_{(u,v)}\mathcal{F}(u_k(s,x), v_k(s,x), \chi_k(s))(u,v) = \lambda(s)(u,v), \ (u,v) \in \mathcal{X} \times \mathcal{X}.$$
(3.36)

Then solution  $(u_k(s, x), v_k(s, x), \chi_k(s))$  will be asymptotically stable if and only if the real part of any eigenvalue  $\lambda(s)$  to (3.36) is negative for  $s \in (-\delta, \delta)$ .

Sending s to 0, we know that  $\lambda = \lambda(0) = 0$  is a simple eigenvalue of  $D_{(u,v)}\mathcal{F}(\bar{u},\bar{v},\chi_k)(u,v) = \lambda(u,v)$  or equivalently, the following eigenvalue problem

$$\begin{cases}
D_1 u'' + \chi_k \bar{u}\phi(\bar{v})v'' - b_1 \bar{u}u - c_1 \bar{u}v = \lambda u, & x \in (0, L), \\
D_2 v'' - b_2 \bar{v}u - c_2 \bar{v}v = \lambda v, & x \in (0, L), \\
u'(x) = v'(x) = 0, & x = 0, L;
\end{cases}$$
(3.37)

moreover, it has a one-dimensional eigen-space  $\mathcal{N}\left(D_{(u,v)}\mathcal{F}(\bar{u},\bar{v},\chi_k)\right) = \{(Q_k,1)\cos\frac{k\pi x}{L}\}$  and one can also prove that  $(Q_k,1)\cos\frac{k\pi x}{L} \notin \mathcal{R}\left(D_{(u,v)}\mathcal{F}(\bar{u},\bar{v},\chi_k)\right)$  following the same analysis that leads to (3.19).

Multiplying (3.37) by  $\cos \frac{k\pi x}{L}$  and integrating it over (0, L) by parts give rise to

$$\begin{pmatrix} -D_1 \left(\frac{k\pi}{L}\right)^2 - b_1 \bar{u} - \lambda & -\chi_k \left(\frac{k\pi}{L}\right)^2 \bar{u}\phi(\bar{v}) - c_1 \bar{u} \\ -b_2 \bar{v} & -D_2 \left(\frac{k\pi}{L}\right)^2 - c_2 \bar{v} - \lambda \end{pmatrix} \begin{pmatrix} \int_0^L u \cos\frac{k\pi x}{L} dx \\ \int_0^L v \cos\frac{k\pi x}{L} dx \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

where the eigenvalue  $\lambda$  satisfies

$$\bar{p}_k(\lambda) = \lambda^2 + \bar{T}_k \lambda + \bar{D}_k = 0.$$

with

$$\bar{T}_k = (D_1 + D_2) \left(\frac{k\pi}{L}\right)^2 + b_1 \bar{u} + c_2 \bar{v} > 0,$$

and

$$\bar{D}_k = \left(D_1 \left(\frac{k\pi}{L}\right)^2 + b_1 \bar{u}\right) \left(D_2 \left(\frac{k\pi}{L}\right)^2 + c_2 \bar{v}\right) - \left(\chi_k \bar{u}\phi(\bar{v})\left(\frac{k\pi}{L}\right)^2 + c_1 \bar{u}\right) b_2 \bar{v}.$$

If  $\chi_k \neq \chi_{k_0} = \min_{k \in \mathbb{N}^+} \chi_k$ ,  $\bar{D}_k < 0$  thanks to (3.14), therefore  $\bar{p}_k(\lambda)$  hence (3.37) always have a positive root  $\lambda(0)$  for all  $k \neq k_0$ . From the standard eigenvalue perturbation theory in [24], (3.36) always has a positive root  $\lambda(s)$  for small s if  $k \neq k_0$ . This finishes the proof of the instability part.

To study the stability of  $(u_{k_0}(s,x), v_{k_0}(s,x), \chi_{k_0}(s))$ , we first note that  $\bar{p}_{k_0}(\lambda)$  or (3.37) with  $k = k_0$  has two eigenvalues, one being negative and the other being zero. Hence we need to investigate the asymptotic behavior of the zero eigenvalue as  $s \approx \neq 0$ . According to Corollary 1.13 in [13], there exists an interval I with  $\chi_{k_0} \in I$  and  $C^1$ -smooth functions  $(\chi, s) : I \times (-\delta, \delta) \to (\mu(\chi), \lambda(s))$  such that  $(\mu(\chi_{k_0}), \lambda(0)) = (0, 0)$ ; moreover,  $\lambda(s)$  is the only eigenvalue in any fixed neighbourhood of the complex plane origin and  $\mu(\chi)$  is the only eigenvalue of the following eigenvalue problem around  $\chi_{k_0}$ 

$$D_{(u,v)}\mathcal{F}(\bar{u},\bar{v},\chi)(u,v) = \mu(u,v), \ (u,v) \in \mathcal{X} \times \mathcal{X}, \tag{3.38}$$

or equivalently

$$\begin{cases}
D_1 u'' + \chi \bar{u}\phi(\bar{v})v'' - b_1 \bar{u}u - c_1 \bar{u}v = \mu u, & x \in (0, L), \\
D_2 v'' - b_2 \bar{v}u - c_2 \bar{v}v = \mu v, & x \in (0, L), \\
u'(x) = v'(x) = 0, & x = 0, L;
\end{cases}$$
(3.39)

furthermore, the eigenfunction of (3.38) can be represented by  $(u(\chi, x), v(\chi, x))$ , which depends on  $\chi$  smoothly and is uniquely determined by  $(u(\chi_{k_0}, x), v(\chi_{k_0}, x)) = (Q_{k_0} \cos \frac{k_0 \pi x}{L}, \cos \frac{k_0 \pi x}{L})$  and  $(u(\chi, x), v(\chi, x)) - (Q_{k_0} \cos \frac{k_0 \pi x}{L}, \cos \frac{k_0 \pi x}{L}) \in \mathcal{Z}$ , with  $Q_{k_0}$  and  $\mathcal{Z}$  being given by (3.15) and (3.18) respectively.

Differentiating (3.39) with respect to  $\chi$  and then taking  $\chi = \chi_{k_0}$ , we have that

$$\begin{cases}
D_1 \dot{u}'' - \bar{u}\phi(\bar{v}) \left(\cos\frac{k_0\pi x}{L}\right)'' + \chi_{k_0} \bar{u}\phi(\bar{v})\dot{v}'' - b_1 \bar{u}\dot{u} - c_1 \bar{u}\dot{v} = \dot{\mu}(\chi_{k_0})Q_{k_0}\cos\frac{k_0\pi x}{L}, \\
D_2 \dot{v}'' - b_2 \bar{v}\dot{u} - c_2 \bar{v}\dot{v} = \dot{\mu}(\chi_{k_0})\cos\frac{k_0\pi x}{L}, \\
\dot{u}'(x) = \dot{v}'(x) = 0, \ x = 0, L,
\end{cases}$$
(3.40)

where in (3.40) denotes the derivative taken with respect to  $\chi$  and evaluated at  $\chi = \chi_{k_0}$ , i.e.,  $\dot{u} = \frac{\partial u(\chi,x)}{\partial \chi}\Big|_{\chi=\chi_{k_0}}$ ,  $\dot{v} = \frac{\partial v(\chi,x)}{\partial \chi}\Big|_{\chi=\chi_{k_0}}$ .

Testing (3.40) by  $\cos \frac{k\pi x}{L}$  over (0, L) yields

$$\begin{pmatrix} -D_1 \left(\frac{k\pi}{L}\right)^2 - b_1 \bar{u} & -\chi_{k_0} \left(\frac{k\pi}{L}\right)^2 \bar{u}\phi(\bar{v}) - c_1 \bar{u} \\ -b_2 \bar{v} & -D_2 \left(\frac{k\pi}{L}\right)^2 - c_2 \bar{v} \end{pmatrix} \begin{pmatrix} \int_0^L \dot{u} \cos\frac{k_0 \pi x}{L} dx \\ \int_0^L \dot{v} \cos\frac{k_0 \pi x}{L} dx \end{pmatrix} = \begin{pmatrix} \left(\dot{\mu}(\chi_{k_0}) Q_{k_0} - \bar{u}\phi(\bar{v}) \left(\frac{k_0 \pi}{L}\right)^2\right) \frac{L}{2} \\ \dot{\mu}(\chi_{k_0}) \frac{L}{2} \end{pmatrix}.$$

The coefficient matrix is singular thanks to (3.14), then we must have that  $\frac{D_1(\frac{k_0\pi}{L})^2 + b_1\bar{u}}{b_2\bar{v}} = \frac{\dot{\mu}(\chi_{k_0})Q_{k_0} - \bar{u}\phi(\bar{v})(\frac{k_0\pi}{L})^2}{\dot{\mu}(\chi_{k_0})}$  which, in light of (3.15), implies that

$$\dot{\mu}(\chi_{k_0}) = -\frac{b_2 \bar{u}\phi(\bar{v})\bar{v}\left(\frac{k_0\pi}{L}\right)^2}{(D_1 + D_2)\left(\frac{k_0\pi}{L}\right)^2 + b_1\bar{u} + c_2\bar{v}} < 0.$$

According to Theorem 1.16 in [13], the functions  $\lambda(s)$  and  $-s\chi'_{k_0}(s)\dot{\mu}(\chi_{k_0})$  have the same zeros and signs near s=0, and

$$\lim_{s \to 0} \frac{-s\chi'_{k_0}(s)\dot{\mu}(\chi_{k_0})}{\lambda(s)} = 1, \text{for } \lambda(s) \neq 0,$$

then we conclude that  $\operatorname{sgn}(\lambda(0)) = \operatorname{sgn}(K_2)$  in light of  $K_1 = 0$ . Following the standard perturbation theory, one can show that  $\operatorname{sgn}(\lambda(s)) = \operatorname{sgn}(K_2)$  for  $s \in (-\delta, \delta)$ ,  $\delta$  being small. This finishes the proof of Theorem 3.3.

According to Theorem 3.3, if  $k_0$  is the positive integer that minimizes  $\chi_k$  over  $\mathbb{N}^+$ , the bifurcation branch  $\Gamma_{k_0}(s)$  around  $(\bar{u}, \bar{v}, \chi_{k_0})$  is stable if it turns to the left and it is unstable if

it turns to the right. However, for all  $k \neq k_0$ ,  $\Gamma_k(s)$  is always unstable for s being small. It is unknown about the global behavior of the continuum of  $\Gamma_k(s)$ . We discuss this in details in Section 5.

Proposition 1 indicates that  $(\bar{u}, \bar{v})$  loses its stability as  $\chi$  surpasses  $\chi_{k_0}$ . Theorem 3.3 shows that the stability is lost to the nonconstant steady state  $(u_{k_0}(s, x), v_{k_0}(s, x))$  in the form  $(Q_{k_0}, 1) \cos \frac{k_0 \pi x}{L}$  and we call this the stable wavemode of (1.1). If the initial value  $(u_0, v_0)$  being a small perturbation from  $(\bar{u}, \bar{v})$ , then spatially inhomogeneous patterns can emerge through this mode, at least when  $\chi$  is around  $\chi_{k_0}$ , therefore stable patterns with interesting structures, such as interior spikes, transition layers, etc. can develop as time evolves. Rigorous analysis of the boundary spike solutions will be analyzed in the coming section.

If the interval length L is small, we see that

$$\chi_k = \frac{\left(D_1(\frac{k\pi}{L})^2 + b_1 \bar{u}\right) \left(D_2(\frac{k\pi}{L})^2 + c_2 \bar{v}\right) - b_2 c_1 \bar{u}\bar{v}}{b_2(\frac{k\pi}{L})^2 \bar{u}\phi(\bar{v})\bar{v}} \approx \frac{D_1 D_2(\frac{k\pi}{L})^2}{b_2\phi(\bar{v})\bar{u}\bar{v}},$$

therefore  $\chi_1 = \chi_{k_0} = \min_{k \in \mathbb{N}^+} \chi_k$  and Theorem 3.3 shows that the only stable pattern is  $(u_1(s,x), v_1(s,x))$  which is spatially monotone. It is easy to see that  $k_0$  increases if L increases. This indicates that small domain only supports monotone stable solutions, while large domain supports non-monotone stable solutions. Indeed, one can construct non-monotone solutions to (3.1) by reflecting and periodically extending the monotone ones at the boundary points  $0, \pm L, \pm 2L, \dots$ 

# 4 Boundary spikes with limiting diffusion rates

This section is devoted to investigate positive solutions to (3.1) with large amplitude compared to the small amplitude bifurcating solutions. In particular, we study the solution profiles as  $D_1$  and  $\chi$  approach infinity with  $\chi/D_1 \in (0, \infty)$  being fixed. Since we will show that small  $D_2$  gives rise to boundary spikes for system (3.1), we put  $\epsilon = \sqrt{D_2}$ , and we put  $\phi(v) \equiv 1$  in (3.1) without loss of our generality. It is the goal of this section to investigate positive solutions with large amplitude to the following system

$$\begin{cases}
(D_1 u' + \chi u v')' + \left(-1 + \frac{1}{a_1 + b_1 u + c_1 v}\right) u = 0, & x \in (0, L), \\
\epsilon^2 v'' + \left(-1 + \frac{1}{a_2 + b_2 u + c_2 v}\right) v = 0, & x \in (0, L), \\
u'(x) = v'(x) = 0, & x = 0, L,
\end{cases}$$
(4.1)

Our main results are the followings.

**Theorem 4.1.** Let  $r = \frac{\chi}{D_1} \in (0, \infty)$  and  $a_2 \in (0, 1)$  be fixed. For any  $\epsilon > 0$  being small, we can find  $\bar{D} > 0$  large such that if  $D_1 > \bar{D}$ , there always exists a nonconstant positive solution (u, v) to (4.1). Moreover, as  $D_1 \to \infty$ , (u(x), v(x)) converges to  $(\lambda_{\epsilon} e^{-rv_{\epsilon}(x)}, v_{\epsilon}(x))$  uniformly in [0, L], where  $\lambda_{\epsilon}$  is a positive constant and  $\lambda_{\epsilon} \to 0$  as  $\epsilon \to 0$ ;  $v_{\epsilon}(x)$  is a positive function of x and  $v_{\epsilon}(x) \to \frac{1-a_2}{c_2}$  compact uniformly on (0, L] and  $v_{\epsilon}(0) \to \frac{1-a_2}{2c_2}$ .

According to Theorem 4.1, u has a boundary spike and v has a boundary layer at x = 0 if  $D_1$  and  $\chi$  are sufficiently large and  $D_2$  is sufficiently small. The elliptic spiky solutions can be used to model segregation phenomenon through inter–specific competitions.

#### 4.1 Convergence to shadow system

Theorem 4.1 is an immediate consequence of several preliminary results. We first study (4.1) by passing  $D_1$  to infinity. To this end, we need the following a prior estimates.

**Lemma 4.2.** Let (u, v) be any positive solution to (4.1). Then there exists a positive constant  $C_0$  independent of  $D_1$  and  $\chi$  such that

$$0 < \max_{x \in [0,L]} v(x), \|v(x)\|_{C^2([0,L])} \le C_0; \tag{4.2}$$

moreover, if  $\frac{\chi}{D_1}$  is bounded, there exists  $C_1 > 0$  such that

$$||u'(x)||_{L^2} \le C_1. \tag{4.3}$$

*Proof.* We have from the Maximum Principles that

$$\max_{x \in [0,L]} v(x) \le \frac{1 - a_2}{c_2};$$

therefore v is bounded in  $C^2([0,L])$  thanks to the standard elliptic Schauder estimates.

On the other hand, we integrate the u-equation over (0, L) to see that

$$\int_{0}^{L} u(x)dx = \int_{0}^{L} \frac{u(x)}{a_1 + b_1 u(x) + c_1 v(x)} dx \le \frac{L}{b_1}.$$

Testing the u-equation by u and integrating it over (0, L) give rise to

$$D_1 \int_0^L (u')^2 dx = -\chi \int_0^L uu'v' dx + \int_0^L f(u,v)u dx$$

$$\leq \frac{D_1}{2} \int_0^L (u')^2 dx + \left(\frac{\chi^2 \|v'\|_{L^\infty}}{2D_1} + \frac{1}{a_1} - 1\right) \int_0^L u^2 dx,$$

therefore  $\|u'(x)\|_{L^2}$  is bounded for finite  $\frac{\chi}{D_1}$  thanks to the Gagliardo-Nirenberg inequality:  $\|u\|_{L^2(0,L)}^2 \leq \delta \|u'\|_{L^2(0,L)}^2 + C(\delta) \|u\|_{L^1(0,L)}^2$ , where  $\delta > 0$  is an arbitrary constant and  $C(\delta) > 0$  only depends on  $\delta$  and L.

We now study the asymptotic behaviors of positive solutions (u, v) to (4.1) by passing advection rate  $D_1$  and advection rate  $\chi$  to infinity. We assume that  $\frac{\chi}{D_1}$  remains bounded in this process, therefore both u and v are bounded as in Lemma 4.2.

**Proposition 2.** Let  $(u_i, v_i)$  be positive solutions of (4.1) with  $(D_{1,i}, \epsilon_i, \chi_i) = (D_1, \epsilon, \chi)$ . Denote  $\frac{\chi_i}{D_{1,i}} = r_i$ . Assume that  $\chi_i \to \infty$ ,  $\epsilon_i \to \epsilon \in (0, \infty)$  and  $r_i \to r \in (0, \infty)$  as  $i \to \infty$ , then there exists a nonnegative constant  $\lambda_{\epsilon}$  such that  $u_i e^{r_i v_i} \to \lambda_{\epsilon}$  uniformly on [0, L] as  $i \to \infty$ ; moreover,  $(u_i, v_i) \to (\lambda_{\epsilon} e^{-rv_{\epsilon}}, v_{\epsilon})$  in  $C^1([0, L]) \times C^1([0, L])$  after passing to a subsequence if necessary, where  $v_{\epsilon} = v_{\epsilon}(x)$  satisfies the following shadow system

$$\begin{cases}
\epsilon^{2}v_{\epsilon}'' + \left(-1 + \frac{1}{a_{2} + b_{2}\lambda_{\epsilon}e^{-rv_{\epsilon}} + c_{2}v_{\epsilon}}\right)v_{\epsilon} = 0, & x \in (0, L), \\
\int_{0}^{L} \left(-1 + \frac{1}{a_{1} + b_{1}\lambda_{\epsilon}e^{-rv_{\epsilon}} + c_{1}v_{\epsilon}}\right)\lambda_{\epsilon}e^{-rv_{\epsilon}}dx = 0, \\
v_{\epsilon}'(0) = v_{\epsilon}'(L) = 0.
\end{cases} (4.4)$$

*Proof.* Since  $v_i$  is bounded in  $C^2([0,L])$  uniformly for all  $\chi_i$  and  $D_{1,i} > 0$ . By the compact embedding,  $v_i$  converges to some  $v_{\infty}$  in  $C^1([0,L])$  as  $i \to \infty$ , after passing to a subsequence if necessary. On the other hand, we integrate the u-equation in (4.1) over (0,x) to have

$$u_i' + r_i u_i v_i' = \frac{1}{D_{1,i}} \int_0^x (1 - \frac{1}{a_1 + b_1 u_i + c_1 v_i}) u_i dx.$$

Denoting  $w_i = u_i e^{r_i v_i}$ , we have that

$$\left| e^{-r_i v_i} w_i' \right| \le \frac{1 + \frac{1}{a_1}}{D_{1,i}} \int_0^x u_i dx.$$
 (4.5)

Sending i to  $\infty$  in (4.5), we conclude that  $w'_i \to 0$  uniformly on [0, L]. Therefore,  $w_i = u_i e^{rv_i}$  converges to a nonnegative constant  $\lambda_{\epsilon}$ . Moreover we can use standard elliptic regularity theory

to show that  $v_{\epsilon}$  is  $C^{\infty}$ -smooth and it satisfies the shadow system (4.4).

Proposition 2 implies that when both  $D_1$  and  $\chi$  are sufficiently large, the steady state (u(x), v(x)) can be approximated by the structures of the shadow system (4.4). We want to remark that the arguments in this Proposition carry over to the case of higher-dimensional bounded domains. To find boundary spikes of (1.1)–(1.2), we study the asymptotic behavior of the shadow system with small diffusion rate  $\epsilon$  and then investigate the original system with large diffusion and advection rates. This approach is due to the idea from [25], applied by [38, 44] in reaction–diffusion systems and developed by [30, 31] for reaction–diffusion system with cross–diffusions.

#### 4.2 Boundary spike layers of the shadow system

We proceed to construct boundary–spike solutions to (4.4). Our results state as follows.

**Proposition 3.** Assume that  $r \in (0, \infty)$  and  $a_2 \in (0, 1)$ . Then there exists a  $\epsilon_0 > 0$  such that (4.4) has a nonconstant positive solution  $(\lambda_{\epsilon}, v_{\epsilon}(\lambda_{\epsilon}, x))$  for all  $\epsilon \in (0, \epsilon_0)$ ; moreover,  $\lambda_{\epsilon} \to 0$  as  $\epsilon \to 0$  and

$$v_{\epsilon}(\lambda_{\epsilon}, x) \to \begin{cases} v^* = \frac{1 - a_2}{c_2}, & compact \ uniformly \ for \ x \in (0, L], \\ \frac{1 - a_2}{2c_2}, & x = 0. \end{cases}$$

$$(4.6)$$

For  $\epsilon$  being sufficiently small,  $v_{\epsilon}(\lambda_{\epsilon}; x)$  has a single inverted boundary spike at x = 0, then one can construct solutions to (4.4) with multi- boundary and interior spikes by periodically reflecting and extending  $v_{\epsilon}(x)$  at  $x = \pm L$ ,  $\pm 2L$ ,...

To prove Proposition 3, we first choose  $\lambda > 0$  to be a predetermined fixed constant and establish nonconstant positive solutions  $v_{\epsilon}(\lambda; x)$  to the following problem

$$\begin{cases}
\epsilon^{2}v_{\epsilon}'' + \left(-1 + \frac{1}{a_{2} + b_{2}\lambda e^{-rv_{\epsilon}} + c_{2}v_{\epsilon}}\right)v_{\epsilon} = 0, & x \in (0, L), \\
v_{\epsilon}(x) > 0, x \in (0, L); v_{\epsilon}'(0) = v_{\epsilon}'(L) = 0
\end{cases}$$
(4.7)

then we find  $\lambda = \lambda_{\epsilon} > 0$  such that  $(\lambda_{\epsilon}, v_{\epsilon}(\lambda_{\epsilon}; x))$  satisfies the integral constraint

$$\int_0^L \left( -1 + \frac{1}{a_1 + b_1 \lambda_{\epsilon} e^{-rv_{\epsilon}} + c_1 v_{\epsilon}} \right) \lambda_{\epsilon} e^{-rv_{\epsilon}} dx = 0.$$
 (4.8)

Denote

$$w_{\epsilon}(x) = v^* - v_{\epsilon}(x) = \frac{1 - a_2}{c_2} - v_{\epsilon}(x)$$
 (4.9)

then (4.7) becomes

$$\begin{cases}
\epsilon^2 w_{\epsilon}'' + f(\lambda; w_{\epsilon}) = 0, & x \in (0, L), \\
w_{\epsilon}'(0) = w_{\epsilon}'(L) = 0,
\end{cases}$$
(4.10)

where

$$f(\lambda; w_{\epsilon}) = -\left(-1 + \frac{1}{1 + b_2 \lambda e^{-r(v^* - w_{\epsilon})} - c_2 w_{\epsilon}}\right) \left(v^* - w_{\epsilon}\right). \tag{4.11}$$

It is equivalent to study the existence and asymptotic behaviors of (4.10) in order to prove Proposition 3. We first collect some facts about  $f(\lambda; s)$  introduced in (4.11). We denote

$$f(\lambda; s) = \frac{g(\lambda; s)(v^* - s)}{1 + g(\lambda; s)}, s \in (0, \infty),$$

where

$$g(\lambda; s) = b_2 \lambda e^{-r(v^* - s)} - c_2 s.$$
 (4.12)

**Lemma 4.3.** Let  $r \in (0, \infty)$ . For each  $\lambda \in (0, \frac{c_2}{b_2 r}e^{rv^* - \frac{r}{c_2} - 1})$ ,  $g(\lambda; s)$  has two positive roots  $s_1 < s_2$  and  $1 + g(\lambda; s)$  has two positive roots  $s_3 < s_4$  such that  $s_1 < s_3 < s_4 < s_2$ ; moreover as  $\lambda \to 0$ ,  $(s_1, s_3) \to (0, \frac{1}{c_2})$  and  $s_2, s_4 \to \infty$ .

*Proof.* It is easy to see that the roots of g(s) or 1+g(s) must be positive if they exist; moreover, if 1+g(s) has positive roots, so does g(s). Hence we only need to show that 1+g(s) has positive roots and it is equivalent to show that its minimum over  $\mathbb{R}^+$  is negative. Let  $s^*$  be the critical point of 1+g(s), we have from straightforward calculations that

$$\begin{cases} g(0) = b_2 \lambda e^{-rv^*} > 0, \\ g'(s^*) = 0, \ s^* = \frac{1}{r} \ln(\frac{c_2 e^{rv^*}}{b_2 \lambda r}) > \frac{1}{c_2} + \frac{1}{r}, \\ g''(s) > 0, \ s \in (-\infty, \infty), \end{cases}$$

therefore 1 + g(s) is convex and its minimum value is  $1 + g(s^*) = 1 + \frac{c_2}{r} - \frac{c_2}{r} \ln(\frac{c_2 e^{rv^*}}{b_2 \lambda r}) < 0$  as desired. Since  $s_i$  is continuous in  $\lambda$ , putting  $\lambda = 0$  in 1 + g(s) gives  $s_1 = 0$  and  $s_3 = \frac{1}{c_2}$ . Moreover,  $s_2 > s_4 > s^* \to \infty$  as  $\lambda \to 0$ . This finishes the proof.

Graphes of g and 1+g are presented in Figure 1.

**Lemma 4.4.** Let  $r \in (0, \infty)$ . Then  $s_1 < v^* < s_3$  if

$$\lambda < \min\left\{\frac{c_2}{b_2 r} e^{rv^* - \frac{r}{c_2} - 1}, \frac{1 - a_2}{b_2}\right\}. \tag{4.13}$$




Figure 1: Graphes of g(s) and 1 + g(s).

*Proof.* We only need to show that  $-1 = g(s_3) < g(v^*) < g(s_1) = 0$  and  $g(s^*) < -1$ , where we already have from above that  $g(v^*) = b_2 \lambda - c_2 v^*$ ; moreover,  $g(s^*) = \frac{c_2}{r} - \frac{c_2}{r} \ln(\frac{c_2 e^{rv^*}}{b_2 \lambda r}) < -1$ .  $\square$ 

We shall assume condition (4.13) from now on. Let us introduce the notations

$$\tilde{w}(x) = w(x) - s_1$$

and

$$\tilde{f}(\lambda;s) = f(\lambda;s+s_1) = -\left(-1 + \frac{1}{1 + b_2 \lambda e^{-r(v^*-s_1-s)} - c_2(s+s_1)}\right) \left(v^* - s_1 - s\right), \quad (4.14)$$

then  $\tilde{w}$  satisfies

$$\begin{cases} \epsilon^2 \tilde{w}'' + \tilde{f}(\lambda; \tilde{w}) = 0, & x \in (0, L), \\ \tilde{w}'_{\epsilon}(0) = \tilde{w}'_{\epsilon}(L) = 0. \end{cases}$$

$$(4.15)$$

Moreover, if  $\tilde{w}_{\epsilon}(x)$  is a positive solution to (4.15) then

$$v_{\epsilon}(x) = v^* - s_1 - \tilde{w}_{\epsilon} \tag{4.16}$$

is a solution to (4.7).

**Lemma 4.5.** For each  $\lambda$  satisfying (4.13), the following problem has a unique solution  $W_0 =$ 

 $W_0(\lambda;z)$ 

$$\begin{cases} W_0'' + \tilde{f}(\lambda; W_0) = 0, z \in \mathbb{R}, \\ W_0(\lambda; 0) = \frac{v^* - s_1}{2} > 0, z \in (0, \infty); W_0'(\lambda; 0) = 0, \\ \lim_{z \to \pm \infty} W_0(\lambda; z) = 0, \end{cases}$$
(4.17)

such that  $W_0 \in C^2(\mathbb{R})$ ,  $W_0(\lambda; z) > 0$  in  $\mathbb{R}$  and  $W_0'(\lambda; z)z < 0$  for  $z \in \mathbb{R} \setminus \{0\}$ . Moreover,  $W_0(\lambda; z)$  is radially symmetric and  $W_0$ ,  $W_0'$ ,  $W_0''$  decays exponentially at  $\infty$  uniformly in  $\lambda$ , i.e., there exists  $C_0$ ,  $\eta > 0$  independent of  $\lambda$  such that

$$0 \le W_0(\lambda; z), |W_0'(\lambda; z)|, |W_0''(\lambda; z)| \le C_0 e^{-\eta|z|}, z \in \mathbb{R}.$$

*Proof.* We introduce the transformation

$$\tilde{F}(\lambda; s) = \int_0^s \tilde{f}(\lambda; t) dt = \int_{s_1}^{s+s_1} f(\lambda; t) dt.$$

therefore  $\tilde{F}(\lambda;s) < 0$  for  $s \in (0, v^* - s_1)$ —see the graph of  $\tilde{f}$  for example; on the other hand, we can show that  $\lim_{s \to (s_3 - s_1)^-} \tilde{F}(\lambda;s) = +\infty$  and according to Intermediate value theorem, there exists  $s_0 \in (v^* - s_1, s_3 - s_1)$  such that  $\tilde{F}(\lambda;s_0) = 0$ ; moreover,  $s_0 = \inf\{s > 0 \mid \tilde{F}(\lambda;s) = 0\}$  and  $\tilde{f}(\lambda;s_0) > 0$ . Therefore  $\tilde{f}$  satisfies condition (6.2) in [6] and Theorem 5 there implies our existence results. Moreover, the exponential decay follows from Remark 6.3 in [6], since  $\tilde{f}'(\lambda;0) < 0$  for all  $\lambda$  satisfying (4.13).

By the unique solution  $W_0$  to (4.17), we now construct a boundary spike to (4.14) hence a boundary layer to (4.7). To this end, we choose a smooth cut-off function  $\rho(x)$  such that  $\rho(x) \equiv 1$  for  $|x| \leq \frac{L}{3}$ ,  $\rho(x) \equiv 0$  for  $|x| \geq \frac{2L}{3}$  and  $\rho(x) \in [0,1]$  for  $x \in \mathbb{R}$ . Denoting

$$W_{\epsilon,\lambda}(x) = \rho(x)W_0(\lambda; x/\epsilon),$$

we want to prove that (4.15) has a solution in the form  $\tilde{w}_{\epsilon}(\lambda, x) = W_{\epsilon, \lambda}(x) + \epsilon \psi(\lambda, x)$ . Then  $\psi$  satisfies

$$\mathcal{L}_{\epsilon}\psi + \mathcal{P}_{\epsilon} + \mathcal{Q}_{\epsilon} = 0, \tag{4.18}$$

where

$$\mathcal{L}_{\epsilon} = \epsilon^2 \frac{d^2}{dx^2} + \tilde{f}_w(\lambda; W_{\epsilon, \lambda}(x)), \tag{4.19}$$

$$\mathcal{P}_{\epsilon} = \epsilon^{-1} \left( \epsilon^2 \frac{d^2}{dx^2} W_{\epsilon,\lambda} + \tilde{f}(\lambda; W_{\epsilon,\lambda}) \right), \tag{4.20}$$

and

$$Q_{\epsilon} = \epsilon^{-1} \Big( \tilde{f}(\lambda; W_{\epsilon, \lambda} + \epsilon \psi) - \tilde{f}(\lambda; W_{\epsilon, \lambda}) - \epsilon \psi \tilde{f}_w(\lambda; W_{\epsilon, \lambda}) \Big). \tag{4.21}$$

According to (4.18)–(4.21),  $\mathcal{P}_{\epsilon}$  and  $\mathcal{Q}_{\epsilon}$  measure the accuracy that  $W_{\epsilon,\lambda}(x)$  approximates solution  $w_{\epsilon}(\lambda,x)$ . Our existence result is a consequence of several lemmas. Set

$$C_{\mathbf{n}}^{2}([0,L]) = \{u \in C^{2}([0,L]) : u'(0) = u'(L) = 0\},\$$

then we first present the following set of results.

**Lemma 4.6.** Let  $r \in (0, \infty)$ . Suppose that  $a_2 \in (0, 1)$  and  $\lambda$  satisfies (4.13). Then there exists a small  $\epsilon_0$  such that if  $\epsilon \in (0, \epsilon_0)$ ,  $\mathcal{L}_{\epsilon}$  with domain  $C_n^2([0, L])$  has an inverse  $\mathcal{L}_{\epsilon}^{-1}$ ; moreover,  $\mathcal{L}_{\epsilon}^{-1} : C([0, L]) \to C([0, L])$  is uniformly bounded in  $\epsilon$ , i.e., there exists  $C_0$  independent of  $\epsilon$  such that

$$\sup_{x \in [0,L]} |\mathcal{L}_{\epsilon}^{-1} f| \le C_0 \sup_{x \in [0,L]} |f|, \forall f \in C([0,L]). \tag{4.22}$$

**Lemma 4.7.** Suppose that the conditions in Lemma 4.6 hold. Then there exist  $C_1 > 0$  and small  $\epsilon_1 > 0$  such that for all  $\epsilon \in (0, \epsilon_1)$ 

$$\sup_{x \in [0,L]} |\mathcal{P}_{\epsilon}(x)| \le C_1.$$

**Lemma 4.8.** Suppose that the conditions in Lemma 4.6 hold. For each  $R_0 > 0$ , denote  $\mathcal{B}(R_0) = \{f | \sup_{x \in [0,L]} |f| < R_0\}$ . Then there exist  $C_2 = C_2(R_0) > 0$  and small  $\epsilon_2 = \epsilon_2(R_0) > 0$  such that for all  $\epsilon \in (0, \epsilon_2)$ ,

$$\sup_{x \in [0,L]} |\mathcal{Q}_{\epsilon}[\psi_i]| \le C_2 \epsilon \sup_{x \in [0,L]} |\psi_i|, \ \forall \psi_i \in \mathcal{B}(R_0), \tag{4.23}$$

$$\sup_{x \in [0,L]} |\mathcal{Q}_{\epsilon}[\psi_1] - \mathcal{Q}_{\epsilon}[\psi_2]| \le C_2 \epsilon \sup_{x \in [0,L]} |\psi_1 - \psi_2|, \ \forall \psi_1, \psi_2 \in \mathcal{B}(R_0). \tag{4.24}$$

We want to point out that Lemma 4.6 generalizes Lemma 5.3 in [47] which holds for  $L^p$ , p > 1. Assuming Lemmas 4.6–4.8, we prove the following results of positive solutions to (4.15).

**Proposition 4.** Let  $r \in (0, \infty)$  and  $a_2 \in (0, 1)$ . Suppose that (4.13) is satisfied. There exists a small  $\epsilon_4 > 0$  such that for all  $\epsilon \in (0, \epsilon_4)$ , (4.15) has a positive solution  $\tilde{w}_{\epsilon}(\lambda; x) \in C([0, L])$  such

that

$$\sup_{x \in (0,L)} |\tilde{w}_{\epsilon}(\lambda;x) - W_{\epsilon}(\lambda;x)| \le C_4 \epsilon,$$

where  $C_4$  is a positive constant independent of  $\epsilon$ . In particular,

$$\lim_{\epsilon \to 0^{+}} \tilde{w}_{\epsilon}(\lambda; x) = \begin{cases} 0, \text{ compact uniformly on } (0, L], \\ v^{*} - s_{1} = \frac{1 - a_{2}}{c_{2}} - s_{1} > 0, \ x = 0, \end{cases}$$

$$(4.25)$$

where  $s_1$  is the positive root of  $g(\lambda; s) = 0$  obtained in Lemma 4.3.

*Proof.* We shall apply the Fixed point theorem on (4.18) to show that  $\tilde{w}_{\epsilon} \in C([0, L])$  takes the form of  $\tilde{w}_{\epsilon} = W_{\epsilon, \lambda} + \epsilon \psi$  for a smooth function  $\psi$ . We define

$$S_{\epsilon}[\psi] = -\mathcal{L}_{\epsilon}^{-1}(\mathcal{P}_{\epsilon} + \mathcal{Q}[\psi]). \tag{4.26}$$

Then  $S_{\epsilon}$  is a bounded linear operator from C([0,L]) to C([0,L]) uniform in  $\lambda$ . Moreover, choosing  $R_0 \geq 2C_0C_1$ , we have that  $\sup_{x \in (0,L)} |\mathcal{L}^{-1}\mathcal{P}_{\epsilon}| \leq C_0C_1$  thanks to Lemma 4.6 and 4.7. Therefore, it follows from Lemma 4.8 that if  $\epsilon$  is small,

$$\sup_{x \in (0,L)} |\mathcal{S}_{\epsilon}[\psi]| \le C_0 C_1 + C_0 C_2 \epsilon R_0 \le R_0, \ \forall \psi \in \mathcal{B}(R_0).$$

and

$$\sup_{x \in (0,L)} |\mathcal{S}_{\epsilon}[\psi_1] - \mathcal{S}_{\epsilon}[\psi_2]| \leq \frac{1}{2} \sup_{x \in (0,L)} |\psi_1 - \psi_2|, \ \forall \psi_1, \psi_2 \in \mathcal{B},$$

hence  $S_{\epsilon}$  is a contraction mapping on  $\mathcal{B}(R_0)$  for small positive  $\epsilon$ . We conclude from the Banach Fixed Point Theorem that  $S_{\epsilon}$  has a fixed point  $\psi_{\epsilon}$  in  $\mathcal{B}$ , which is a smooth solution of (4.15). It is easy to show that  $\tilde{w}_{\epsilon}$  satisfies (4.25) and this finishes the proof of Proposition 4.

Thanks to (4.13)–(4.15),  $v_{\epsilon}(x) = v^* - s_1 - \tilde{w}_{\epsilon}(x)$  is a smooth solution of (4.11). We have the following results.

Corollary 1. Under the same condition as in Proposition 4. There exists  $\epsilon_4 > 0$  such that for all  $\epsilon \in (0, \epsilon_4(\delta))$ , (4.7) has a positive solution  $\tilde{v}_{\epsilon}(\lambda; x) \in C([0, L])$  satisfying

$$\lim_{\epsilon \to 0^{+}} \tilde{v}_{\epsilon}(\lambda; x) = \begin{cases} v^{*} - s_{1} = \frac{1 - a_{2}}{c_{2}} - s_{1}, & compact \ uniformly \ on \ (0, L], \\ (v^{*} - s_{1})/2, & x = 0, \end{cases}$$

$$(4.27)$$

Now we prove Lemmas 4.6-4.8.

Proof of Lemma 4.6. We argue by contradiction. Choose a positive sequence  $\{\epsilon_i\}_{i=1}^{\infty}$  with  $\epsilon_i \searrow 0$  as  $i \to \infty$ . Suppose that there exist  $\Psi_i(x) \in C^2([0,L])$  and  $h_i(x) \in C([0,L])$  satisfying

$$\begin{cases}
\mathcal{L}_{\epsilon_i} \Psi_i = h_i, & x \in (0, L), \\
\Psi_i'(0) = \Psi_i'(L) = 0,
\end{cases} (4.28)$$

such that  $\sup_{x\in[0,L]}|\Psi_i(x)|=1$  and  $\sup_{x\in[0,L]}|h(x)|\to 0$  as  $i\to\infty$ . Define

$$\tilde{\Psi}_i(z) = \Psi_i(\epsilon x), \tilde{h}_i(z) = h_i(\epsilon_i x), \text{ and } \tilde{W}_{\epsilon_i,\lambda}(z) = W_{\epsilon_i,\lambda}(\epsilon_i x),$$

then

$$\frac{d^2\tilde{\Psi}_i}{dz^2} + \tilde{f}_w(\lambda; \tilde{W}_{\epsilon_i, \lambda}(z))\tilde{\Psi}_i = \tilde{h}_i, z \in \left(0, \frac{L}{\epsilon_i}\right).$$

Let  $R_0 < \frac{L}{\epsilon_i}$  be an arbitrarily chosen but fixed constant. Without loss of our generality, we assume  $\sup_{[0,L]} \Psi_i(x) = 1$ . Then we infer from the boundedness of  $\{\tilde{h}_i(z)\}_{i=1}^{\infty}$  and the elliptic Schauder estimate that  $\{\tilde{\Psi}_i(z)\}_{i=1}^{\infty}$  is bounded in  $C^2([0,R_0])$ , therefore there exists a subsequence as  $i \to \infty$  such that  $\tilde{\Psi}_i(z) \to \tilde{\Psi}_0(z)$  in  $C^{1,\theta}([0,R_0])$  thanks to the compact embedding  $C^2([0,R_0]) \subset C^{1,\theta}([0,R_0])$ ,  $\theta \in (0,1)$ . On the other hand, since  $\tilde{f}_w(\lambda;\tilde{W}_{\epsilon_i,\lambda}(z)) \to \tilde{f}_w(\lambda;W_0(\lambda;z))$  and  $\tilde{h}_i(z) \to 0$  uniformly on  $[0,R_0]$ ,  $\frac{d^2\tilde{\Psi}_i}{dz^2}$  converges in  $C([0,R_0])$  hence  $\tilde{\Psi}_i \to \tilde{\Psi}_0$  in  $C^2([0,R_0])$ . Applying standard diagonal argument and elliptic regularity theory, we can show that  $\tilde{\Psi}_0$  is in  $C^{\infty}(0,\infty)$  and it satisfies

$$\begin{cases} \frac{d^2 \tilde{\Psi}_0}{dz^2} + \tilde{f}_w(\lambda; W_0(\lambda; z)) \tilde{\Psi}_0 = 0, & z \in (0, \infty), \\ \tilde{\Psi}'_0(0) = 0, & \end{cases}$$
(4.29)

where  $W_0$  is the unique solution to (4.17).

Now we show that  $\tilde{\Psi}_0(0) = 1$ . To this end, let  $Q_i \in [0, L]$  be such that  $\tilde{\Psi}_i(Q_i) = 1$ . We claim that  $Q_i \leq C_0 \epsilon$  for some bounded  $C_0 > 0$  independent of  $\epsilon$ . In order to prove this claim, we argue by contradiction and assume that  $\frac{Q_i}{\epsilon} \to \infty$  as  $i \to \infty$ . Define

$$\hat{\Psi}_i(z) = \Psi_i(\epsilon z + Q_i)$$
 and  $\hat{h}_i(z) = h_i(\epsilon_i z + Q_i)$ 

for  $\epsilon_i z + Q_i \in [0, L]$  or  $z \in [-\frac{Q_i}{\epsilon_i}, \frac{L - Q_i}{\epsilon_i}]$ , therefore

$$\begin{cases}
\frac{d^2\hat{\Psi}_i}{dz^2} + \tilde{f}_w(\lambda; \hat{W}_{\epsilon_i, \lambda}(z))\hat{\Psi}_i = \hat{h}_i, & z \in \left[-\frac{Q_i}{\epsilon_i}, \frac{L - Q_i}{\epsilon_i}\right], \\
\hat{\Psi}_i(0) = \sup_{z \in \left[-\frac{Q_i}{\epsilon_i}, \frac{L - Q_i}{\epsilon_i}\right]} \hat{\Psi}_i(z) = 1, \frac{d\hat{\Psi}_i(0)}{dz} = 0.
\end{cases} (4.30)$$

By the same arguments as above, we can show that  $\hat{\Psi}_i$  converges to some  $\hat{\Psi}_0$  (at least) in  $C^2((-\infty,0])$  and  $\hat{\Psi}_0$  is in  $C^\infty((-\infty,0])$ ; moreover, since  $\frac{Q_i}{\epsilon_i} \to \infty$ , we have from the exponential decaying property of  $W_0$  that  $\hat{W}_{\epsilon_i,\lambda}(z) = \rho(\epsilon_i z + Q_i)W_0(\lambda; \frac{\epsilon_i z + Q_i}{\epsilon}) \to 0$  and

$$\frac{d^2\hat{\Psi}_0}{dz^2} + \tilde{f}_w(\lambda;0)\hat{\Psi}_0 = 0, \frac{d\hat{\Psi}_0(0)}{dz} = 0, \sup_{z \in (-\infty,0)} \hat{\Psi}_0(0) = 1,$$

then we have from Maximum Principle that  $\hat{\Psi}_0''(0) = -\tilde{f}_w(\lambda;0)\hat{\Psi}_0(0) = -\tilde{f}_w(\lambda;0) \le 0$ , however, this is a contradiction to the fact that  $\tilde{f}_w(\lambda;0) < 0$ . This proves our claim and we must have that  $\tilde{\Psi}_0(0) = \sup_{z \in [0,\infty)} \tilde{\Psi}_0(z) = 1$  in (4.29).

Differentiate (4.17) with respect to z and we have that

$$\begin{cases}
\frac{d^3 W_0}{dz^3} + \tilde{f}_w(\lambda; W_0) \frac{dW_0}{dz} = 0, z \in (-\infty, \infty), \\
\frac{dW_0(0)}{dz} = 0, W_0(0) = \sup_{z \in (-\infty, \infty)} W_0(z) > 0,
\end{cases}$$
(4.31)

Multiplying (4.29) by  $\frac{dW_0}{dz}$  and integrating it over  $(0,\infty)$  lead us to

$$\int_{0}^{\infty} \frac{d^{2}\tilde{\Psi}_{0}}{dz^{2}} \frac{dW_{0}}{dz} + \tilde{f}_{w}(\lambda; W_{0})\tilde{\Psi}_{0} \frac{dW_{0}}{dz} = 0.$$
(4.32)

Multiplying (4.33) by  $\tilde{\Psi}_0$  and integrating it over  $(0,\infty)$  lead us to

$$\int_{0}^{\infty} \frac{d^{3}W_{0}}{dz^{3}} \tilde{\Psi}_{0} + \tilde{f}_{w}(\lambda; W_{0}) \frac{dW_{0}}{dz} \tilde{\Psi}_{0} = 0.$$
(4.33)

We infer from (4.32) and (4.33) and the integrations by parts that

$$\begin{split} 0 &= \int_0^\infty \tilde{\Psi}_0'' W_0' - \tilde{\Psi}_0 W_0''' \\ &= \int_0^\infty (\tilde{\Psi}_0' W_0')' - \tilde{\Psi}_0' W_0'' - \left( (\tilde{\Psi}_0 W_0'')' - \tilde{\Psi}_0' W_0'' \right) \\ &= \int_0^\infty (\tilde{\Psi}_0' W_0')' - (\tilde{\Psi}_0 W_0'')' \\ &= \tilde{\Psi}_0' W_0' \Big|_0^\infty - \tilde{\Psi}_0 W_0'' \Big|_0^\infty, \end{split}$$

where ' denotes the derivative against z. In light of the exponential decay of  $W'_0$ ,  $W''_0$  at infinity and the fact  $W''_0(0) \neq 0$ , we have that  $\tilde{\Psi}_0(0) = 0$ , and this is a contradiction. The proof of this lemma is finished.

*Proof of Lemma* 4.7. Substituting  $W_{\epsilon,\lambda}(x) = \rho(x)W_0(\lambda;\frac{x}{\epsilon})$  into (4.20) gives rise to

$$\mathcal{P}_{\epsilon} = \epsilon^{-1} \left( \epsilon^{2} \rho''(x) W_{0}(\lambda; x/\epsilon) + 2\epsilon \rho'(x) (W_{0})_{x}(\lambda; x/\epsilon) \right.$$

$$\left. + \rho(x) (W_{0})_{xx}(\lambda; x/\epsilon) + \tilde{f}(\lambda; \rho(x) W_{0}(\lambda; x/\epsilon)) \right)$$

$$= \epsilon^{-1} \left( \epsilon^{2} \rho''(x) W_{0}(\lambda; x/\epsilon) + 2\epsilon \rho'(x) (W_{0})_{x}(\lambda; x/\epsilon) \right.$$

$$\left. - \rho(x) \tilde{f}(\lambda; W_{0}(\lambda; x/\epsilon)) + \tilde{f}(\lambda; \rho(x) W_{0}(\lambda; x/\epsilon)) \right).$$

$$(4.34)$$

If  $x \in [0, \frac{L}{3}] \cup [\frac{2L}{3}, L]$ ,  $\rho(x) \equiv 1$  hence  $\mathcal{P}_{\epsilon}$  is bounded for all  $\lambda$ . If  $x \in (\frac{L}{3}, \frac{2L}{3})$ , since  $W_0$  decays exponentially at  $\infty$ , we can also see that  $\mathcal{P}_{\epsilon}$  is bounded. Therefore  $\sup_{x \in [0,L]} |\mathcal{P}_{\epsilon}|$  is bounded.  $\square$ 

*Proof of Lemma* 4.8. We have from the Intermediate Value Theorem that

$$\sup_{x \in [0,L]} |\mathcal{Q}_{\epsilon}[\psi](x)| = \epsilon^{-1} \sup_{x \in [0,L]} \left[ \tilde{f}(\lambda, W_{\epsilon,\lambda} + \epsilon \psi) - \tilde{f}(\lambda, W_{\epsilon,\lambda}) - \epsilon \psi \tilde{f}_{w}(\lambda, W_{\epsilon,\lambda}) \right] 
= \epsilon^{-1} \sup_{x \in [0,L]} \left| \int_{0}^{1} \left[ \tilde{f}_{w}(\lambda, W_{\epsilon,\lambda} + t\epsilon \psi) - \tilde{f}_{w}(\lambda, W_{\epsilon,\lambda}) \right] (\epsilon |\psi|) dt \right| 
= \epsilon \sup_{x \in [0,L]} |\psi^{2}| \int_{0}^{1} \int_{0}^{t} \sup_{x \in [0,L]} \left| \tilde{f}_{ww}(\lambda, W_{\epsilon,\lambda} + s\epsilon \psi) \right| ds dt 
\leq C \epsilon \sup_{x \in [0,L]} |\psi|^{2},$$
(4.35)

and

$$\begin{split} \sup_{x \in [0,L]} |\mathcal{Q}_{\epsilon}[\psi_{1}] - \mathcal{Q}_{\epsilon}[\psi_{2}]| \\ = &\epsilon \sup_{x \in [0,L]} \Big| \int_{0}^{1} \int_{0}^{t} \tilde{f}_{ww}(\lambda, W_{\epsilon,\lambda} + s\epsilon\psi_{1}) \psi_{1}^{2} - \tilde{f}_{ww}(\lambda, W_{\epsilon,\lambda} + s\epsilon\psi_{2}) \psi_{2}^{2} ds dt \Big| \\ = &\epsilon \sup_{x \in [0,L]} \Big| \int_{0}^{1} \int_{0}^{t} \tilde{f}_{ww}(\lambda, W_{\epsilon,\lambda} + s\epsilon\psi_{1}) \psi_{1}^{2} - \int_{0}^{1} \int_{0}^{t} \tilde{f}_{ww}(\lambda, W_{\epsilon,\lambda} + s\epsilon\psi_{2}) \psi_{1}^{2} \\ &+ \int_{0}^{1} \int_{0}^{t} \tilde{f}_{ww}(\lambda, W_{\epsilon,\lambda} + s\epsilon\psi_{2}) \psi_{1}^{2} - \int_{0}^{1} \int_{0}^{t} \tilde{f}_{ww}(\lambda, W_{\epsilon,\lambda} + s\epsilon\psi_{2}) \psi_{2}^{2} ds dt \Big| \\ = &\epsilon \sup_{x \in [0,L]} \Big| \int_{0}^{1} \int_{0}^{t} \tilde{f}_{ww}(\lambda, W_{\epsilon,\lambda} + s\epsilon\psi_{1}) - \tilde{f}_{ww}(\lambda, W_{\epsilon,\lambda} + s\epsilon\psi_{2}) ds dt \Big| |\psi_{1}|^{2} \\ &+ \epsilon \sup_{x \in [0,L]} \Big| \int_{0}^{1} \int_{0}^{t} \tilde{f}_{ww}(\lambda, W_{\epsilon,\lambda} + s\epsilon\psi_{1}) ds dt \Big| |\psi_{1}^{2} - \psi_{2}^{2}| \\ \leq &\epsilon C_{0} \sup_{x \in [0,L]} |\psi_{1}|^{2} \sup_{x \in [0,L]} |\psi_{1} - \psi_{2}| + \epsilon C_{1} \sup_{x \in [0,L]} |\psi_{1} + \psi_{2}| \sup_{x \in [0,L]} |\psi_{1} - \psi_{2}| \\ \leq &(C_{0}R^{2} + 2C_{1}R) \epsilon \sup_{x \in [0,L]} |\psi_{1} - \psi_{2}| \\ \leq &C_{2} \epsilon \sup_{x \in [0,L]} |\psi_{1} - \psi_{2}|, \end{split}$$

where we have used the fact that  $|\tilde{f}_{ww}(\Psi_i)|$ ,  $|\tilde{f}_{www}(\Psi_i)|$  are bounded for  $\Psi_i \in \mathcal{B}(R_0)$ .

We proceed to prove Proposition 3. Therefore we need to find  $\lambda = \lambda_{\epsilon}$  satisfying (4.8) and (4.13). In particular, we shall show that  $\lambda_{\epsilon} \to 0$  as  $\epsilon \to 0$ . We define

$$G(\epsilon, \lambda) = \int_0^L \left( -1 + \frac{1}{a_1 + b_1 \lambda e^{-rv_{\epsilon}} + c_1 v_{\epsilon}} \right) \lambda e^{-rv_{\epsilon}} dx, \tag{4.37}$$

over  $(\epsilon, \lambda) \in (-\delta, \delta) \times (-\delta, \delta)$ , where  $\delta > 0$  be a small constant. We put  $v_{\epsilon}(\lambda; x) \equiv v^* = \frac{1-a_2}{c_2}$  if  $\epsilon < 0$  or  $\lambda < 0$ .

*Proof of Proposition* 3. It is easy to see that

$$G(0,0) = \lim_{\epsilon,\lambda \to 0} G(\epsilon,\lambda) = \int_0^L \left( -1 + \frac{1}{a_1 + b_1 \lambda e^{-rv_{\epsilon}} + c_1 v_{\epsilon}} \right) \lambda e^{-rv_{\epsilon}} = 0.$$

For  $\epsilon, \lambda > 0$ , we have that

$$\frac{\partial G(\epsilon,\lambda)}{\partial \lambda} = \int_0^L \left( r - \frac{(a_1 + c_1 v_{\epsilon}(\lambda;x))r + c_1}{(a_1 + b_1 \lambda e^{-rv_{\epsilon}(\lambda;x)} + c_1 v_{\epsilon}(\lambda;x))^2} \right) \lambda e^{-rv_{\epsilon}(\lambda;x)} \frac{\partial v_{\epsilon}(\lambda;x)}{\partial \lambda} dx 
+ \int_0^L \left( \frac{a_1 + c_1 v_{\epsilon}(\lambda;x)}{(a_1 + b_1 \lambda e^{-rv_{\epsilon}(\lambda;x)} + c_1 v_{\epsilon}(\lambda;x))^2} - 1 \right) e^{-rv_{\epsilon}(\lambda;x)} dx,$$
(4.38)

where

$$\frac{\partial v_{\epsilon}(\lambda;x)}{\partial \lambda} = -\frac{\partial \tilde{w}_{\epsilon}(\lambda;x)}{\partial \lambda} - \frac{\partial s_{1}}{\partial \lambda} = -\frac{\partial \tilde{w}_{\epsilon}(\lambda;x)}{\partial \lambda} - \frac{b_{2}e^{-r(v^{*}-s_{1})}}{c_{2}}$$

since  $v_{\epsilon} = v^* - \tilde{w} - s_1$ . We claim that  $\sup_{x \in [0,L]} |\frac{\partial w_{\epsilon}(\lambda;x)}{\partial \lambda}|$  is uniformly bounded in  $\epsilon$  and  $\lambda$ . According to (4.15), we have

$$\epsilon^2 \frac{d^2 \frac{\partial \tilde{w}_{\epsilon}}{\partial \lambda}}{dx^2} + \tilde{f}_w(\lambda, \tilde{w}_{\epsilon}(\lambda; x)) \frac{\partial \tilde{w}_{\epsilon}}{\partial \lambda} + \tilde{f}_{\lambda}(\lambda; \tilde{w}_{\epsilon}(\lambda; x)) = 0,$$

where

$$\tilde{f}_{\lambda} = -\frac{b_2 e^{r(\tilde{w}_{\epsilon}(\lambda;x) - v^*)} (v^* - \tilde{w}_{\epsilon}(\lambda;x))}{(1 + b_2 \lambda e^{r(\tilde{w}_{\epsilon}(\lambda;x) - v^*)} - c_2 \tilde{w}_{\epsilon}(\lambda;x))^2}$$

and it is uniformly bounded in  $\epsilon$  and  $\lambda$ . Write  $\tilde{\mathcal{L}}_{\epsilon} = \epsilon^2 \frac{d^2}{dx^2} + \tilde{f}_w(\lambda; \tilde{w}_{\epsilon}(\lambda; x))$ . By the standard perturbation arguments, we can show that  $\tilde{\mathcal{L}}_{\epsilon}$  has an inverse  $\tilde{\mathcal{L}}_{\epsilon}^{-1}$ :  $C([0, L]) \to C([0, L])$  which is uniformly bounded in  $\epsilon$ . Therefore  $\frac{\partial \tilde{w}_{\epsilon}}{\partial \lambda}$  is bounded in C([0, L]). Now we conclude from the Lebesgue Dominated Convergence Theorem  $\frac{\partial G(\epsilon, \lambda)}{\partial \lambda}$  is continuous around  $(\epsilon, \lambda) = (0, 0)$ ; moreover, since  $v^* = \frac{1-a_2}{c_2} \neq \frac{1-a_1}{c_1}$ , we have

$$\frac{\partial G(0,0)}{\partial \lambda} = \int_0^L \left( \frac{a_1 + c_1 v^*}{(a_1 + c_1 v^*)^2} - 1 \right) e^{-rv^*} dx \neq 0, \tag{4.39}$$

then Proposition 4 follows from the Implicit Function Theorem.

#### 4.3 Boundary spike and boundary layer to the full system

Now we prove Theorem 4.1 by showing that the solutions to (4.1) perturb from its shadow system (4.7) when  $D_1$  is sufficiently large. First of all, we let  $s = \frac{1}{D_1}$ ,  $\psi = v$ ,  $\phi = ue^{rv} - \tau$ , and  $u = (\phi + \tau)e^{-rv}$ , where  $\tau$  is a constant. Since  $\frac{\partial(\phi,\psi)}{\partial(u,v)} = \begin{pmatrix} e^{rv} & rue^{rv} \\ 0 & 1 \end{pmatrix}$  is invertible, (4.1) is

equivalent as  $F(s, \phi, \tau, \psi) = 0$ , where

$$F(s,\phi,\tau,\psi) = \begin{pmatrix} \phi_{xx} - r\phi_x\psi_x + sP\{(-1 + \frac{1}{a_1 + b_1(\phi + \tau)e^{-r\psi} + c_1\psi})(\phi + \tau)\} \\ \epsilon^2\psi_{xx} + (-1 + \frac{1}{a_2 + b_2(\phi + \tau)e^{-r\psi} + c_2\psi})\psi \\ \int_0^L (-1 + \frac{1}{a_1 + b_1(\phi + \tau)e^{-r\psi} + c_1\psi})(\phi + \tau)e^{-r\psi} dx \end{pmatrix}$$
(4.40)

We decompose C([0,L]) into  $Y_0 \oplus Y_1$ , where  $Y_0 = \{y \in C([0,L]) | \int_0^L u dx = 0\}$ . Define the projection operator  $P: C^2([0,L]) \to Y_0$  by

$$P(u) = u - \frac{1}{L} \int_0^L u(x) dx.$$

Proof of Theorem 4.1. Let  $(\lambda_{\epsilon}, v_{\epsilon}(\lambda; x))$  be a solution of (4.7), then we see that  $F(0, 0, \lambda_{\epsilon}, v_{\epsilon}(x)) = 0$ . Moreover,  $F(s, \phi, \tau, \psi)$  is an analytic from  $\mathbb{R}^+ \times C^2([0, L]) \times \mathbb{R} \times C^2([0, L])$  to  $Y_0 \times C([0, L]) \times \mathbb{R}$  and its Frechét derivative with respect to  $(\phi, \tau, \psi)$  at  $(s, \phi, \tau, \psi) = (0, 0, \lambda_{\epsilon}, v_{\epsilon})$  is given by

$$D_{(\phi,\tau,\psi)}F|_{(0,0,\lambda_{\epsilon},v_{\epsilon})} = \begin{pmatrix} \frac{d^{2}}{dx^{2}} - rv_{\epsilon}'\frac{d}{dx} & 0 & 0\\ \tilde{f}_{\lambda}(\lambda_{\epsilon};\tilde{w}_{\epsilon}) & \tilde{f}_{\lambda}(\lambda_{\epsilon};\tilde{w}_{\epsilon}) & \epsilon^{2}\frac{d^{2}}{dx^{2}} + \tilde{f}_{w}(\lambda_{\epsilon};\tilde{w}_{\epsilon})\\ E_{1} & E_{1} & E_{2} \end{pmatrix}$$
(4.41)

where

$$E_1 = \int_0^L \left( \frac{a_1 + c_1 v_{\epsilon}}{(a_1 + b_1 \lambda_{\epsilon} e^{-rv_{\epsilon}} + c_1 v_{\epsilon})^2} - 1 \right) e^{-rv_{\epsilon}}$$

and

$$E_2 = \int_0^L \left( \frac{b_1 r \lambda_{\epsilon} e^{-rv_{\epsilon}} - c_1}{(a_1 + b_1 \lambda_{\epsilon} e^{-rv_{\epsilon}} + c_1 v_{\epsilon})^2} + r \right) \lambda_{\epsilon} e^{-rv_{\epsilon}}$$

The operator  $\frac{d^2}{dx^2} - rv'_{\epsilon}(\lambda; x) \frac{d}{dx}$  is an isomorphism from  $C^2([0, L]) \cap Y_0$  onto  $Y_0$ . Indeed, choosing u in the kernel of this operator and writing  $u = \sum_{k=0}^{\infty} S_k \cos \frac{k\pi x}{L}$ , we have that  $-S_k \frac{k\pi}{L} \cos \frac{k\pi x}{L} + rS_k v'_{\epsilon}(\lambda; x) \sin \frac{k\pi x}{L} = 0$ , which is impossible unless  $S_k = 0$ . Therefore  $D_{(\phi, \tau, \psi)} F|_{(0, 0, \lambda_{\epsilon}, v_{\epsilon})}$  is bounded invertible if and only if the following problem

$$\begin{cases}
\epsilon^{2} \frac{d^{2} \eta_{\epsilon}}{dx^{2}} + \tilde{f}_{w}(\lambda_{\epsilon}; \tilde{w}_{\epsilon}(\lambda, x)) \eta_{\epsilon} + \tilde{f}_{\lambda}(\lambda_{\epsilon}; \tilde{w}_{\epsilon}(\lambda, x)) \xi_{\epsilon} = 0, \\
\xi_{\epsilon} \int_{0}^{L} \left( \frac{a_{1} + c_{1} v_{\epsilon}}{(a_{1} + b_{1} \lambda_{\epsilon} e^{-rv_{\epsilon}} + c_{1} v_{\epsilon})^{2}} - 1 \right) e^{-rv_{\epsilon}} dx \\
+ \int_{0}^{L} \left( \frac{b_{1} r \lambda_{\epsilon} e^{-rv_{\epsilon}} - c_{1}}{(a_{1} + b_{1} \lambda_{\epsilon} e^{-rv_{\epsilon}} + c_{1} v_{\epsilon})^{2}} + r \right) \lambda_{\epsilon} e^{-rv_{\epsilon}} \eta_{\epsilon} dx = 0.
\end{cases} (4.42)$$

has only the trivial solution  $\xi_{\epsilon} = 0$  and  $\eta_{\epsilon}(x) \equiv 0$ . We argue by contradiction and without loss of generality, we assume that  $|\xi_{\epsilon}| + \sup_{[0,L]} |\eta_{\epsilon}(x)| = 1$ . Since  $\lambda_{\epsilon} \to 0$  as  $\epsilon \to 0$ , we infer from the second equation in (4.42) that  $\xi_{\epsilon} \to 0$  as  $\epsilon \to 0$ . On the other hand, since  $\tilde{f}_{\lambda}(\lambda_{\epsilon}; \tilde{w}_{\epsilon}(\lambda, x))$  is bounded and  $\tilde{\mathcal{L}}_{\epsilon}$  has a bounded inverse uniform in  $\epsilon$ , we conclude from the first equation that  $\sup_{[0,L]} |\eta_{\epsilon}| \to 0$  as  $\epsilon \to 0$ . However, this contradicts our assumption. Therefore  $D_{(\phi,\tau,\psi)}F|_{(0,0,\lambda_{\epsilon},v_{\epsilon})}$  is nonsingular at  $(0,0,\lambda_{\epsilon},v_{\epsilon}(\lambda;x))$  and Theorem 4.1 follows from the Implicit Function Theorem.

It is an interesting and also important mathematical question to study the stability of the single boundary spike solutions to the shadow system and the full system. To this end, one needs more detailed asymptotic expansions on the perturbed solutions than we obtained and this is postponed to future study. We refer to [22, 29, 27, 28, 50, 51] and the references therein for stability analysis of spiky solutions of some closely related reaction—diffusion systems and/or the shadows systems.

## 5 Conclusions and Discussions

In this paper, we investigate the interspecific competition through a reaction-advection-diffusion system with Beddington–DeAngelis response functionals. Species segregation phenomenon is modeled by the global existence of time-dependent solutions and the formation of stable boundary spike/layers over one-dimensional finite domains. There are several findings in our theoretical results. First of all, we prove that system (1.1)-(1.2) admits global-in-time classical positive solutions which are uniformly bounded globally for any space dimensions. For  $\Omega = (0, L)$ , we show that large directed dispersal rate  $\chi$  of the species destabilizes the homogeneous solution  $(\bar{u},\bar{v})$  and nonconstant positive steady states bifurcate from the homogeneous solution. Moreover, we study the linearized stability of the bifurcating solutions which shows that the only stable bifurcating solutions must have a wavemode number  $k_0$ , which is a positive integer that minimizes bifurcating value  $\chi_k$  in (3.14).—see Theorem 3.2 and Theorem 3.3. That being said, the constant solution loses its stability to the wavemode  $\cos \frac{k_0 \pi x}{L}$  over (0, L). Our result indicates that if the domain size L is sufficiently small, then only the first wavemode  $\cos \frac{\pi x}{L}$  can be stable, which is spatially monotone; moreover, large domain supports stable patterns with more aggregates than small domains. We prove that the steady states of (1.1)–(1.2) over (0, L)admits boundary spike in u and boundary layer in v if  $D_1$  and  $\chi$  are comparably large and  $D_2$ 




Figure 2: The sensitivity function  $\phi(v) \equiv 1$  is selected. Stable wave mode in the form of  $\cos \frac{k_0 \pi x}{L}$ , where  $k_0$  is given in Table 1.  $\chi$  is chosen to be slightly larger than  $\chi_0$  and the rest system parameters are chosen to be the same as in Table 1. Initial data are small perturbations of  $(\bar{u}, \bar{v})$ .

is small. Therefore, we can construct multi-spike or multi-layer solutions to the system by reflecting and periodically extending the single spike over  $\pm L$ ,  $\pm 2L$ ,... These nontrivial structures can be used to the segregation phenomenon through inter–specific competition. Compared to the transition layer in [47], our results suggests the boundary spike solutions to (1.1) is due to the Beddington–DeAngelis dynamics since a similar approach was adapted in [47].

In Table 1, we list the wavemode numbers  $k_0$  and the corresponding minimum bifurcation values for different lengths. Figure 2 gives numerical simulations on the evolutions of stable patterns over different intervals. In Figure 3, we plot the spatial-temporal solutions to illustrate the formation of single boundary spike to u and boundary layer to v. The boundary spike and layer here correspond to those obtained in Theorem 3.1. We propose some problems for future studies. System (1.1) describes a situation that u directs its disperse strategy over the habitat to deal with the population pressures from v, which moves randomly over the domain. From the viewpoint of mathematical modeling, it is interesting to study the situation when both species takes active dispersals over the habitat. Then system (1.1) becomes a double-advection system with nonlinear kinetics. Analysis of the new system becomes much more complicated now since no maximum principle is available for the second equation. We also want to point out that

Domain size $L$	3	5	7	9	11
$k_0$	1	2	2	3	3
$\chi_k$	9.9418	10.392	9.9120	9.9418	9.9647
Domain size $L$	13	15	17	19	21
$k_0$	4	5	5	6	6
$\chi_k$	9.8872	9.9418	9.8937	9.8956	9.9120

Table 1: Stable wavemode number  $k_0$  and the corresponding bifurcation value  $\chi_{k_0}$  for different interval length. The system parameters are  $D_1 = 1, D_2 = 0.1, a_1 = a_2 = 0.5, b_1 = 2, b_2 = 1$  and  $c_1 = 0.5, c_2 = 1$ . According to Theorem 3.2, the stable wavemode is  $\cos \frac{k_0 \pi x}{L}$ . Therefore, stable and nontrivial patterns must develop in the form of  $\cos \frac{k_0 \pi x}{L}$  if  $\chi$  is chosen to be slightly larger than  $\chi_{k_0}$ . We can also see that larger domains support bigger wavemode number. Figure 2 is given to illustrate the wavemode selection mechanism.




Figure 3: Formation of stable single boundary spike of u and boundary layer of v. Diffusion and advection rates are chosen to be  $D_1=5,~\chi=30,~D_2=5\times10^{-3}$ . The rest system parameters are  $a_1=0.2,~b_1=0.8,~c_1=0.1$  and  $a_2=0.6,~b_2=0.2,~c_2=0.4$ . Initial data are  $u_0=\bar{u}+0.5\cos\frac{2\pi x}{5}$  and  $v_0=\bar{v}+0.5\cos\frac{2\pi x}{5}$ , where  $(\bar{u},\bar{v})=(0.933...,0.533...)$ .

it is also interesting to study the effect of environment heterogeneity on the spatial–temporal behaviors of the solutions.

Our bifurcation analysis is carried out around bifurcation point  $(\bar{u}, \bar{v}, \chi_k)$ . It is interesting to investigate the global behavior of the continuum of  $\Gamma_k(s)$ , denoted by  $\mathcal{C}$ . According to the global bifurcation theory in [40] and its developed version in [41], there are three possible happenings

for C: (i) it is either not compact in  $\mathcal{X} \times \mathcal{X} \times \mathbb{R}^+$ , or it contains a point  $(\bar{u}, \bar{v}, \chi_j)$ ,  $j \neq k$ . Actually, by the standard elliptic embeddings, we can show that C is bounded in the axis of  $\mathcal{X} \times \mathcal{X}$  if  $\chi$  is finite. Therefore, it either extends to infinity in the  $\chi$ -axis or it stops at  $(\bar{u}, \bar{v}, \chi_j)$ . It is unclear to us which is the case for  $\chi_k$ . Considering Keller–Segel chemotaxis models over (0, L) without cellular growth, some topology arguments are applied in [9, 48] etc. to show that all solutions on the first branch must either be monotone increasing or deceasing. Therefore  $\chi_1$  can not intersect at  $(\bar{u}, \bar{v}, \chi_j)$  for  $j \neq 1$  since all the solutions around it must be non-monotone. This shows that  $\Gamma_1$  extends to infinity in the  $\chi$  direction. However, the appearance of complex structures of the kinetics in our model inhabits this methodology.

Our numerical simulations suggest that these boundary spikes and layers are globally stable for a wide range of system parameters. Rigorous stability analysis of these spikes is needed to verify our theoretical results. It is also interesting to investigate the large—time behavior of (1.1)—(1.2), which requires totally different approached from we apply in our paper.

# References

- N.D. Alikakos, L<sup>p</sup> bounds of solutions of reaction-diffusion equations, Comm. Partial Differential Equations, 4 (1979), 827–868.
- [2] H. Amann, Dynamic theory of quasilinear parabolic equations. II. Reaction-diffusion systems, Differential Integral Equations, 3 (1990), 13–75.
- [3] H. Amann, Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems, Function Spaces, differential operators and nonlinear Analysis. Teubner, Stuttgart, Leipzig, (1993), 9–126.
- [4] M.A. Aziz-Alaoui and M. Daher Okiye, Boundedness and global stability for a predatorprey model with modified Leslie-Gower and Holling-type II schemes, Appl. Math. Lett., 16 no. 7, October 2003, 1069–1075
- [5] J.R. Beddington, Mutual interference between parasites or predators and its effect on searching efficiency, J. Animal Ecol., 44 (1975), pp. 331–340
- [6] H. Berestycki and P. -L. Lions Nonlinear scalar field equations, I existence of a ground state Arch. Ration. Mech. Anal., 82 (1983), 313–345.
- [7] E. Beretta and Y. Takeuchi, Global asymptotic stability of Lotka-Volterra diffusion models with continuous time delay, SIAM J. Appl. Math., 48 (1988), 627–651

- [8] R. Cantrell, C. Cosner, On the dynamics of predator-prey models with the Beddington-Deangelis functional response, J. Math. Anal. Appl., 257 no. 1, 1 May 2001, 206-222
- [9] A. Chertock, A. Kurganov, X. Wang and Y. Wu, On a chemotaxis model with saturated chemotactic flux, Kinet. Relat. Models, 5 (2012), 51–95.
- [10] C. Cosner, Reaction-diffusion-advection models for the effects and evolution of dispersal, Discrete Contin. Dyn. Syst., 34 (2014), 1701–1745.
- [11] C. Cosnera, D. DeAngelisb, J. Aultc and D. Olsonc Effects of spatial grouping on the functional response of predators, Theoretical Population Biology, 56 (1999), 65–75
- [12] M.G. Crandall and P.H. Rabinowitz, Bifurcation from simple eigenvalues, J. Functional Analysis, 8 (1971) 321–340.
- [13] M.G. Crandalland P.H. Rabinowitz, Bifurcation, perturbation of simple eigenvalues, and linearized stability, Arch. Ration. Mech. Anal., 52 (1973) 161–180.
- [14] D.L. DeAngelis, R.A. Goldstein and R.V. O'Neill, A model for trophic interaction, Ecology, 56 (1975), 881–892
- [15] M. Fan and Y. Kuang, Dynamics of a nonautonomous predator-prey system with the Beddington-DeAngelis functional response, J. Math. Anal. Appl., 295 (2004), 15–39
- [16] D. Henry, Geometric Theory of Semilinear Parabolic Equations, Springer-Verlag, Berlin-New York, 1981.
- [17] M.A. Herrero and J.J.L. Velazquez, Chemotactic collapse for the Keller-Segel model, J. Math. Biol., 35 (1996), 583-623.
- [18] T. Hillen and K.J. Painter, A user's guidence to PDE models for chemotaxis, J. Math. Biol., 58 (2009), 183-217.
- [19] D. Horstmann and M. Winkler, Boundedness vs. blow-up in a chemotaxis system, J. Differential Equations, 215 (2005), 52–107.
- [20] D. Horstmann, From 1970 until present: the Keller-Segel model in chemotaxis and its consequences, Nonlinearity, 105(2003),103-165.
- [21] T.-W. Hwang, Global analysis of the predator-prey system with Beddington-DeAngelis functional response, J. Math. Anal. Appl., 281 (2003), 395–401.
- [22] D. Iron, M.-J. Ward and J. Wei, The stability of spike solutions to the one-dimensional Gierer-Meinhardt model, Phys. D, 150 (2001), 25-62.

- [23] Y. Kan-on and E. Yanagida, Existence of nonconstant stable equilibria in competition-diffusion equations, *Hiroshima Math. J.*, **23** (1993), 193–221.
- [24] T. Kato, Study of partial differential equations by means of functional analysis, Springer Classics in Mathematics, (1996).
- [25] J.P. Keener, Activators and inhibitors in pattern formation, Stud. Appl. Math. 59 (1978), 1–23.
- [26] W. Ko and K. Ryu, Qualitative analysis of a predator-prey model with Holling type II functional response incorporating a prey refuge, J. Differential Equations, 231 (2006), 534– 550.
- [27] T. Kolokolnikov, M.-J. Ward and J. Wei, The existence and stability of spike equilibria in the one-dimensional Gray-Scott model: the pulse-splitting regime, Phys. D, 202 (2005), 258–293.
- [28] T. Kolokolnikov, M.-J. Ward and J. Wei, The existence and stability of spike equilibria in the one-dimensional Gray-Scott model: the low feed-rate regime., Stud. Appl. Math. 115 (2005), 21–71.
- [29] T. Kolokolnikov and J. Wei, Stability of spiky solutions in a competition model with cross-diffusion, SIAM J. Appl. Math. 71 (2011), 1428–1457.
- [30] Y. Lou and W.-M. Ni, Diffusion, self-diffusion and cross-diffusion, J. Differential Equations, 131 (1996), 79–131.
- [31] Y. Lou and W.-M. Ni, Diffusion vs cross-diffusion: An elliptic approach, J. Differential Equations, 154 (1999), 157–190.
- [32] M. Ma, C. Ou and Z.-A. Wang, Stationary solutions of a volume filling chemotaxis model with logistic growth, SIAM Journal on Applied Mathematics 72 (2012), 740–766.
- [33] H. Matano and M. Mimura, Pattern formation in competition-diffusion systems in non-convex domains, Publ. Res. Inst. Math. Sci., 19 (1983), 1049–1079.
- [34] M. Mimura, Stationary patterns of some density-dependent diffusion system with competitive dynamics, Hiroshima Math. J., 11 (1981), 621-635.
- [35] M. Mimura, S.-I. Ei and Q. Fang, Effect of domain-shape on coexistence problems in a competition-diffusion system, J. Math. Biol., 29 (1991), 219–237.
- [36] M. Mimura and K. Kawasaki, Spatial segregation in competitive interaction-diffusion equations, J. Math. Biol., 9 (1980), 49–64.

- [37] V. Nanjundiah, Chemotaxis, signal relaying and aggregation morphology, Journal. Theor. Biol., 42 (1973), 63–105.
- [38] I. Takagi and W.-M. Ni, Point condensation generated by a reaction-diffusion system in axially symmetric domains., Japan J. Indust. Appl. Math. 12 (1995), 327–365.
- [39] W.-M. Ni, Y. Wu and Q. Xu, The existence and stability of nontrivial steady states for SKT competition model with cross-diffusion, Discret Cotin Dyn. Syst., 34 (2014), 5271–5298.
- [40] P. Rabinowitz, Some global results for nonlinear eigenvalue problems, J. Functional Analysis, 7 (1971), 487–513.
- [41] J. Shi and X. Wang, On global bifurcation for quasilinear elliptic systems on bounded domains, J. Differential Equations, 246 (2009), 2788–2812.
- [42] N. Shigesada, K. Kawasaki and E. Teramoto, Spatial segregation of interacting species, J. Theoret. Biol., 79 (1979), 83–99.
- [43] H.L. Smith, Monotone Dynamical Systems: An Introduction to the Theory of Competitive and Cooperative Systems, Math. Surveys Monogr. 41, American Mathematical Society, Providence, RI, 1995.
- [44] I. Takagi, Point-condensation for a reaction-diffusion system., J. Differential Equations 61 (1986), no. 2, 208–249.
- [45] Y. Takeuchi, Global stability in generalized Lotka-Volterra diffusion systems, J. Math. Anal. Appl., 116 (1986), 209-221
- [46] Pierre B. de Villemereuil and A. Lopez-Sepulcre, Consumer functional responses under intra- and inter-specific interference competition, Ecological Modelling, 222 (2011), 419–426
- [47] Q. Wang, C. Gai and J. Yan, Qualitative analysis of a Lotka-Volterra competition system with advection, Discrete Contin. Dvn. Syst. 35 (2015), 1239–1284.
- [48] X. Wang and Q. Xu, Spiky and transition layer steady states of chemotaxis systems via global bifurcation and Helly's compactness theorem, J. Math. Biol., 66 (2013), 1241–1266.
- [49] M. Winkler, Aggregation vs. global diffusive behavior in the higher-dimensional Keller-Segel model, J. Differential Equations, 248 (2010), 2889–2905.
- [50] M. Winter and J. Wei, Stability of monotone solutions for the shadow Gierer-Meinhardt system with finite diffusivity, Differential Integral Equations, 16 (2003), 1153–1180.

- [51] M. Winter and J. Wei, Mathematical aspects of pattern formation in biological systems, Applied Mathematical Sciences, 189. Springer, London, 2014. xii+319 pp. ISBN: 978-1-4471-5525-6; 978-1-4471-5526-3
- [52] F. Yi, J. Wei and J. Shi, Bifurcation and spatiotemporal patterns in a homogeneous diffusive predator-prey system, J. Differential Equations, 246 (2009), 1944–1977