Colours of minor bodies in the outer solar system

II: A statistical analysis revisited

O. R. Hainaut¹, H. Boehnhardt², and S. Protopapa³

- ¹ European Southern Observatory (ESO), Karl Schwarzschild Straße, 85 748 Garching bei München, Germany e-mail: ohainaut@eso.org
- Max-Planck-Institut f
 ür Sonnensystemforschung, Max-Planck Stra
 ße 2, 37 191 Katlenburg-Lindau, Germany
- ³ Department of Astronomy, University of Maryland, College Park, MD 20742-2421, USA

Received May 2012; accepted July 2012

ABSTRACT

We present an update of the visible and near-infrared colour database of Minor Bodies in the Outer Solar System (MBOSSes), which now includes over 2000 measurement epochs of 555 objects, extracted from over 100 articles. The list is fairly complete as of December 2011. The database is now large enough to enable any dataset with a large dispersion to be safely identified and rejected from the analysis. The selection method used is quite insensitive to individual outliers. Most of the rejected datasets were observed during the early days of MBOSS photometry. The individual measurements are combined in a way that avoids possible rotational artifacts. The spectral gradient over the visible range is derived from the colours, as well as the R absolute magnitude M(1,1). The average colours, absolute magnitude, and spectral gradient are listed for each object, as well as the physico-dynamical classes using a classification adapted from Gladman and collaborators. Colour-colour diagrams, histograms, and various other plots are presented to illustrate and investigate class characteristics and trends with other parameters, whose significances are evaluated using standard statistical tests. Except for a small discrepancy for the J-H colour, the largest objects, with M(1,1) < 5, are indistinguishable from the smaller ones. The larger ones are slightly bluer than the smaller ones in J - H. Short-period comets, Plutinos and other resonant objects, hot classical disk objects, scattered disk objects and detached disk objects have similar properties in the visible, while the cold classical disk objects and the Jupiter Trojans form two separate groups of their spectral properties in the visible wavelength range. The well-known colour bimodality of Centaurs is confirmed. The hot classical disk objects with large inclinations, or large orbital excitations are found to be bluer than the others, confirming a previously known result. Additionally, the hot classical disk objects with a smaller perihelion distance are bluer than those that do not come as close to the Sun. The bluer hot classical disk objects and resonant objects have fainter absolute magnitudes than the redder ones of the same class. Finally, we discuss possible scenarios for the origin of the colour diversity observed in MBOSSes, i.e. colouration caused by evolutionary or formation processes.

The colour tables * and all plots are also available on the MBOSS colour web page**, which will be updated when new measurements are published

Key words. Kuiper Belt: general – Comets: general – Technique: photometric – Methods: statistical – Astronomical databases: miscellaneous.

1. Introduction

Minor bodies in the outer solar system (MBOSSes) comprise objects in the Kuiper Belt (KB) and more generally in the Transneptunian (TN) region, as well as small bodies in the giant planets region that came from either the KB or TN regions, but are now no longer immediate members of these environments. Centaurs are considered as escapees (Gladman et al. 2008; Kavelaars et al. 2008), scattered from the KB and TN environments towards the Sun, possibly also representing a major source of short-period comets (SPCs). Some MBOSSes may also be found among the satellites of the giant planets since they might have been stranded there by the gravitational capturing from the Centaur population (Duncan & Levison 1997). Another large population of MBOSSes can be found as long-period (LPCs) or Oort Cloud comets. These objects are believed to have formed in the region of the giant planets and been scattered afterwards into the very distant domains of the solar system (Dones et al. 2004), although it cannot presently be excluded that contamination by extra-solar comets (Levison et al. 2010) may exist among the Oort Cloud population.

The MBOSSes found in the planetary system environment formed and have evolved in the region of the giant planets. They are considered as remnants of the planetesimal population, but their current distance range is not necessarily that of their formation. Nonetheless, MBOSSes may contain valuable information about the environment and the physical conditions at the time of their formation. On the other hand, their prolonged presence in the region of the distant planets and beyond may have caused alterations to their physical properties. For instance, high energy and particle radiation is suspected to modify the colours and albedos of materials in space (de Bergh et al. 2008) and collisions can affect the body either as a whole (fragmentation or growth) or in parts (resurfacing

^{*} Tables 2 and 3 are only available in electronic form at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/.

^{**} http://www.eso.org/~ohainaut/MBOSS

with excavated material). In addition, it cannot be excluded that intrinsic activity may have changed the surface constitution of the MBOSSes, since a number of ices, believed or known to be present in these objects, can sublimate at very large distances from the Sun (for instance CH_4 , CO, and N_2 up to about 45, 65, and 80AU, respectively; Delsemme 1982; Meech & Svoren 2004). Last, but not least, larger MBOSSes may have experienced some alteration to their internal structure (McKinnon et al. 2008). It is thus of interest to characterize the population properties and explore possible connections with the origin of the bodies and/or their evolutionary pathways in the solar system.

With this study, we focus on the photometric properties of MBOSSes and the characterization of the different dynamical populations among them. The photometric properties comprise published measurements of their photometric brightnesses, filter colours, and spectral slopes of reflected continuum light in the visible and near-infrared (NIR) wavelength regions (Doressoundiram et al. 2008). The brightness of an object is a first indicator on its size and albedo; filter colours allow a coarse characterization of the spectral energy distribution of its reflected light and may provide constraints on its surface properties, i.e. its wavelength-dependent reflectivity.

The visible and NIR wavelength region up to 5μ m contains surface-reflected sunlight, i.e. essentially either a bluish, neutral, or a reddened solar spectrum, occasionally with imprinted absorption features from specific surface materials. Continuum colours and gradients are well defined, perhaps with the exception of the HKLM bands for objects with very strong ice absorption features (Trujillo et al. 2011). The photometric parameters of MBOSSes and their correlations with then dynamical and other properties of the bodies were analysed in the past using various different data-sets and statistical methods (for a review, see Doressoundiram et al. 2008, and references therein). The key findings were that the TNOs cover a wide range of colours and spectral slopes in the visible from slightly bluish (-5 to -10%/100 nm) to very red (40 to 55%/100 nm), while in the NIR they display a fairly narrow dispersion around solar colours. Differences in the wavelength-dependent surface reflectivity were noted for dynamically hot and cold Classical Disk Objects (CDOs), with the latter representing a very red population in the solar system. Among the hot CDOs, an anti-correlation with high significance between the surface reddening on one hand and inclination and eccentricity on the other was interpreted as an indicator of evolutionary changes due to impacts and cosmic radiation (Trujillo & Brown 2002; Peixinho et al. 2008). Centaurs may have a bimodal colour distribution with a neutral to slightly red and a very red sub-population (Tegler et al. 2008). A number of weaker correlations between photometric and dynamical properties of KBOs are addressed in the literature. However, no clear convincing picture with quantitative modelling results has evolved, although various attempts to obtain a qualitative understanding of how the surface colours were produced as the result of evolutionary processes at the surface in the outer space environment have been published (for a brief review, see Doressoundiram et al. 2008).

A completely independent approach to MBOSS colour statistics was introduced by Barucci et al. (2005a), where the objects were grouped based on their B, V, R, I, J colours to form taxonomic classes. The colour study was performed using a principal component (PC) analysis (Reyment & Joreskog 1993) and with the G-mode multivariate statistics (Coradini 1977; Fulchignoni et al. 2000). Fulchignoni et al. (2008) describe in detail the method and its main results. In the PC analysis, the original n colours are replaced by the minimum possible number of linear combinations of colours (the m principal components) required to reproduce the original distribution, the n-m additional ones containing only noise. They found that most of the MBOSS colour range is accounted for by a single variable, closely related to the overall slope of the reflectivity spectrum over the visible range (which we define quantitatively in Sect. 2.1). The G-mode analysis reveals two main classes, labeled BB and RR, containing the neutral-grey and very red objects, respectively. The intermediate objects sort themselves into two additional classes, BR and IR. Perna et al. (2010) published an updated list of the taxonomic classes for 151 objects, and studied their distributions over the dynamical classes and other parameters. In the 1970s-1980s, the G-mode-based taxonomic classification of main belt asteroids was very successful (Tholen 1984; Tholen & Barucci 1989; Barucci & Fulchignoni 1990), the resulting classes becoming the basis for the physico-dynamical understanding of these objects, which is also the ultimate goal of the ongoing MBOSS taxonomic classification.

In many cases, the published colour and spectral gradient analyses rely on different data-sets collected at different telescopes and instruments and different groups. The sample sizes have grown with time reaching more than 100 TNOs and Centaurs. A summarizing database of MBOSS colours comprising data from many different papers were first compiled and analysed by Hainaut & Delsanti (2002, hereafter Paper I) and was publicly accessible through the Internet; it is still available from the MBOSS web site, but is superseded by this work.

In this paper, we provide a version of this MBOSS colour database that has been updated both in terms of populations and collected data. It can be used either for population-wide analysis, as done in this paper, or to support other works on specific objects or group of objects either relying on the averaged data presented here (e.g. to get the colour of a given object) or by going back to the original publications listed for each object. We present new statistical analyses of the MBOSS populations based on an enhanced database of photometric brightnesses and colour measurements of the objects published in the literature and applying qualified selection criteria to the data. The qualifying criterion of the data are described in Section 2. Section 3 introduces the statistical methods and their application goals for the MBOSS analysis and outlines the results achieved. Possible interpretations of the findings related to their formation and evolutionary pathways are discussed in Section 4. The paper ends with a brief summary of both the major findings and open and controversial issues, as well as prospects for their clarification in Section 5.

2. Description of the database

The database collects photometric information about MBOSSes, namely of objects in the Kuiper Belt (KBOs) and in the immediate Transneptunian region, of Centaurs, of short- and long-period comets (SPCs and LPCs), and of Jupiter Trojans. It does not contain information on satellites of the giant planets as far as they are considered to be captured MBOSSes, nor of Trojans of other planets. The objects are listed under their current official designation, i.e. number and name if available, or number and provisional designations, or provisional designation only.

For objects presenting cometary activity, the measurements collected here refer to the nucleus only, not to the dust and gas component. In the case of objects with satellites, measurements for the whole system are listed under the main designation (e.g. 26308 = 1998SM165), while an individual member of the system is indicated by a suffix (+B for the first satellite, for instance 134340-Pluto+B refers to Charon). In the statistical analysis, the satellites are not given a special status: they just count as individual objects.

The photometric information in the database comprises magnitudes or filter colours in the *UBVRIJHKL* broadband filter system for the visible and the NIR. In rare cases also spectral gradients and/or colours measured from spectroscopic data of MBOSSes were entered, relying on the spectrum to colours conversion presented in the original paper.

All flavours of the main filters are considered together without any conversion. For instance, in this database, Bessell R, Kron-Cousin R and r' filters are directly listed as R. Most authors calibrate their system using Landolt (1992), thereby naturally unifying these subtly different systems. More exotic filters are included in the database, but not used in this study.

The literature is searched for relevant papers using the SAO/NASA ADS¹, the distant EKO newsletter², the Neese (2011) compilation of TNO and Centaurs at the Planetary Data System³, and the astro-ph preprints from arXiv⁴.

Only magnitude measurements that are explicitly presented as simultaneous are considered as a single epoch. This means that the individual filters must have been observed in a sequence, so that no more than ~1h elapsed during a colour measurement. The only exception was for average colours obtained from full light-curves. By default, spectral gradients and colours obtained from spectroscopy fulfill the "simultaneity" criterion. Photometric measurements not obtained simultaneously are listed as separate data, and are not used for the average colour estimates. The method used to carefully combine measurements from different epochs without introducing colour artifacts from a possible lightcurve is the same as in paper I.

The original database, described in paper I, contained information for only 104 objects. We had neither the means to evaluate the quality of, nor the luxury to reject some of the measurements. The online version continuously grew, and the current MBOSS database includes over 550 objects, with over 2000 measurements extracted from over 100 papers. We can now afford to reject some measurements based upon a careful and quantifiable approach. The process of cleaning data should be approached with caution, as it could potentially select against special or interesting objects. We developed a method to evaluate the quality of a data-set that is robust against genuine outliers. As first noted by Boehnhardt et al. (2001) and discussed later in the present paper, most objects have a fairly linear reflectivity spectrum over the visible wavelength range; we used this to assess the quality of a set of measurements. For each individual measurement, the distance between the data point in a colour-colour diagram and the reddening line (locus of the objects with a linear reflection spectrum, see Section 2.3) was computed as follows. We defined (Cx, Cy) to be the colours of the object, and (Cx, Fy), (Fx, Cy) the two points on the reddening line that have the same x and y as the data point, respectively. The distance estimator was defined as

$$d = \sqrt{\sum_{i=1}^{n} \frac{(C_i - F_i)^2}{n}},$$
(1)

where the C_i and F_i terms are the coordinates in the B - V/V - R and V - R/R - I colour-colour diagrams where available, and n the number of coordinates available (2 or 4). The overall quality estimator for a data-set is

$$D = \sqrt{\sum_{k=1}^{N} \frac{d_k^2}{N}},\tag{2}$$

where d_k are the individual distances and N the number of measurement sets in the considered paper. D behaves like an error in the photometry, expressed in magnitudes. Each of the papers with D > 0.25 were scrutinized—this threshold was selected visually as a 2σ -like limit in the plot. The faintest objects were flagged out and D re-estimated. If D remained above 0.25, the process was iterated. In some cases, this brought D below 0.25, and the remaining measurements were retained, assuming that those rejected were affected by low signal-to-noise ratio. In other cases, the dispersion in the measurements was uncorrelated with magnitude and the full data-set was rejected; we consider that a problem affected that whole paper. This was the case for some of the early literature on MBOSS photometry. Before rejecting a data-set or a measurements, the colours were compared with other measurements of the same object, if available, so as to preserve objects with intrinsic unusual spectra. In addition, the majority of the objects have linear spectra, so even a few objects with non-linear spectra did not strongly affect the overall D estimator of a paper (thanks to the quadratic average), and these objects were then preserved in the database.

This method obviously worked only for measurements in the visible range. Among the 107 papers considered, 33 did not have suitable data, and 7 were globally rejected. For each data-set (including IR-only papers), outliers were individually considered. Only the points that we have a strong reason to believe were affected by a problem were rejected (for instance where a note in the paper reported a problem). In addition, some papers that concentrated on objects with exceptional spectra were obviously preserved. We believe that this statistical approach to the cleaning process with a careful and conservative *a posteriori* assessment ensures that the global data-set is of higher quality.

The database presented in this paper is available online⁵, together with all the plots related to this static version of the database, as a reference. In parallel, as the database is continually evolving with the addition of new measurements, another up-to-date version⁶

http://adswww.harvard.edu

http://www.boulder.swri.edu/ekonews

http://sbn.psi.edu/pds/resource/tnocencol.html

http://arxiv.org/archive/astro-ph

http://www.eso.org/~ohainaut/MBOSS/MBOSS2.dat

⁶ http://www.eso.org/~ohainaut/MBOSS

is available, with all plots and tables. The former should be used only in direct reference to this paper, while the dynamically updated version can be used for further generic studies of MBOSSes. The updated versions are produced in exactly the same way as the static one presented in this paper, with additional objects and measurements processed as described above.

2.1. Content of the database

Each record internally lists the object designation as in the original publication (i.e. typically using the temporary designation for early papers, and the final number in later papers). These were converted to the current official designation, i.e. either number and name when available, number and provisional designation, or only provisional designation.

The internal database lists the measurements for each epoch. One record is constituted by the name of the target, the epoch of the measurements, the reference to the original paper, and the list of measurements for that epoch as they appear in the paper, i.e. as magnitudes, colours, or a combination of both.

The orbital elements were retrieved from the MPCORB file, which is regularly updated from the Minor Planets Center (MPC) website⁷. These elements were used to compute the position of the object at the epoch of the observations (if available). The orbit semi-major axis a, perihelion q, eccentricity e, inclination i, orbital excitation defined as $\mathcal{E} = \sqrt{\sin^2 i + e^2}$, and the helioand geocentric distances r and Δ as well as the solar phase α at the time of the observations were stored with the individual measurements.

The measurements were averaged as in paper I: For a given epoch, the matrix of all possible colours was populated from the magnitudes and colours available at that epoch – but without mixing different epochs. The average colours for an object were then obtained as a weighted average of the corresponding colours, using $1/\sigma$ as the weight, where σ was either the reported or propagated error in the individual colour measurements.

Additionally, for each epoch, we converted the R magnitude (either the one reported, or obtained from another magnitude together with the corresponding colour index) into an absolute $M(1,1,\alpha)$ magnitude with the computed r and Δ using $M(1,1,\alpha) = R - 5 \log(r\Delta)$. As in paper I, we neither made no assumption about the solar phase function, nor corrected for the solar phase effect. The values of $M(1,1,\alpha)$ obtained for different epochs were then averaged into a final absolute R magnitude, which is reported as M(1,1) in Table 2. The solar phase effects were not considered. The observations were obtained at small solar phase angles because of the distance of the objects, and because most of the observations were acquired close to opposition. The colours of the objects, whose study is at the core of this paper, were even less affected than the absolute magnitude. We also did not make any assumption about the albedo of the object, hence did not convert the absolute magnitudes into diameters.

For each object, the slope of the (very low resolution) reflectivity spectrum obtained from the average colour indexes (Jewitt & Meech 1986) was computed as in paper I, via a linear regression over the B-V-R-I range (assigning a lower weight to the B reflectivity), where most objects display a linear reflectivity spectrum (Boehnhardt et al. 2001). This gradient, S, is expressed in percent of reddening per 100 nm.

2.2. Physico-dynamical classes

Since the publication of paper I, the understanding of the dynamical classes in the outer solar system has greatly progressed. Paper I used fairly arbitrary definitions of the dynamical classes; moreover, the fairly small number of objects in each class did not allow for a fine-grained classification. In this paper, we used the dynamical classes defined by Gladman et al. (2008) (the now so-called SSBN08 classification, from the book in which it is published). The membership of the object was allocated using a combination of integration of the orbit over time together with cut-off in the orbital element space.

For this paper, we relied on the membership list published by Gladman et al. (2008), to which about 200 objects had been added from a list generated by C. Ejeta and H. Boehnhardt (private communication) in the context of the Herschel Key Program "TNOs are Cool" (Müller et al. 2009). This list was completed by the MPC catalogue of Jupiter Trojans⁸.

For the objects not included in these lists, we determined the dynamical class using a simplified method: objects with an obvious designation (LPCs, SPCs) were first flagged as such, then for the remaining few objects (only 26 objects in the current database), we used an algorithm based only on the osculating orbital elements, without integration of the orbit, directly inspired by the flow-chart Fig. 1 in Gladman et al. (2008). Furthermore, we separated the classical TNOs between dynamically hot classical disk objects (with $i \ge 5^{\circ}$) and dynamically cold classical disk objects ($i < 5^{\circ}$). The arguments for this separation of CDOs come from the analysis of their orbital properties, although the first indications of different sub-populations among the classical TNOs came from a photometric study of TNOs by Tegler & Romanishin (2000) and were further evaluated in papers by Trujillo & Brown (2002) and Doressoundiram et al. (2002). The dynamical aspects are addressed in Morbidelli et al. (2008a) and Gladman et al. (2008). We chose an inclination of 5° for the separation of the two populations. A split around orbital excitation $\mathcal{E} = 0.12$ gave similar results (only 3 objects among the 89 cold classical TNOs changed class). Clearly, a sharp cut in i, e, or \mathcal{E} space is a simplification, since it can be expected that both the cold and hot populations actually partially overlap. Following Gladman et al. (2008), we also distinguished TNOs as scattered disk objects (SDOs) or detached disk objects (DDOs). The sharp cuts in orbital element space can again only be seen as an approximation, since the actual boundary between the classes is expected to have a "complicated, fractal structure" (Morbidelli et al. 2008a).

The majority of 'resonant' objects belongs to the class of Plutinos in 3:2 resonance with Neptune. Objects belonging to other resonances are summarized as 'Res. others' objects in the analysis below. Finally, some of the SPCs with Centaur-like or-

⁷ http://www.minorplanetcenter.net/iau/Ephemerides

⁸ http://www.minorplanetcenter.net/iau/lists/JupiterTrojans.html

Table 1. Object physico-dynamical classes (see text for the definitions) and number of objects in each class, and some statistics about the overall database.

Class	Number
Jupiter Trojans	80
Resonant (3:2)	47
Resonant (others)	28
Long-period comets (LPCs)	14
Short-period comets (SPCs)	136
Centaurs	35
Scattered disk objects (SDOs)	30
Detached disk objects (DDOs)	28
Cold classical disk objects (cold CDOs)	89
Hot classical disk objects (hot CDOs)	68
Database	
Objects	555
Epochs	2045
Papers	100

bital elements, as defined in Gladman et al. (2008) are re-assigned to that class, following Jewitt (2009a) and Tegler et al. (2008), namely 29P/Schwassmann-Wachmann 1, 30P/Oterma, 165P/LINEAR, 166P/2001 T4, 167P/2004 PY42 174P aka 60 558 Echeclus, C/2001 M10 (NEAT), P/2004 A1 (LONEOS), and P/2005 T3 (Read).

2.3. The database

Table 1 lists the classes considered and the number of objects in each class. Table 2 provides examples of individual objects, listing the object identification and its dynamical class together with the main average colours, the spectral gradient S, and the absolute R magnitude M(1, 1). The full table is available in the online supplement.

Table 2. Object class and colour data for some example objects. The full table is available in electronic form at the CDS, and maintained up-to-date on the MBOSS colour web site.

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	$Grt^{(3)} \pm \sigma$	B-V $\pm \sigma$	V-R $\pm \sigma$	R-I $\pm \sigma$	V-J ±σ	J-H $\pm\sigma$	H-K $\pm\sigma$
1P/Halley	SP Comet	2	13.558 ± 0.596	4.266 ± 2.015	0.720 ± 0.040	0.410 ± 0.030	0.390 ± 0.060	_	_	_
9P/Tempel 1	SP Comet	14	14.560 ± 0.460	9.959 ± 0.826	_	0.468 ± 0.010	0.469 ± 0.013	_		
1172-Aneas	J.Trojan	2	8.729 ± 0.042	9.261 ± 0.863	0.727 ± 0.030	0.510 ± 0.022	0.400 ± 0.030	1.577 ± 0.036	0.430 ± 0.042	0.135 ± 0.036
1994 ES ₂	Classic Cold	1	7.509 ± 0.130		_	_		_		_
1997 SZ ₁₀	Res.(other)	1	8.148 ± 0.060	28.167 ± 2.909	1.140 ± 0.080	0.650 ± 0.030		_		_
1999 CF ₁₁₉	Detached	3	6.933 ± 0.133	14.704 ± 4.163	_	0.622 ± 0.115	0.365 ± 0.106	_	0.380 ± 0.226	_
1999 CX ₁₃₁	Res.(other)	3	6.910 ± 0.107	19.521 ± 4.293	0.918 ± 0.124	0.664 ± 0.128	0.434 ± 0.105	_	0.370 ± 0.238	_
2003 HX ₅₆	Classic Hot	1	7.030 ± 0.209	-1.963 ±13.676	_	0.350 ± 0.226	0.260 ± 0.459	_		_
2060-Chiron	Centaur	34	6.092 ± 0.069	0.114 ± 0.996	0.700 ± 0.020	0.361 ± 0.017	0.325 ± 0.023	1.199 ± 0.110	0.294 ± 0.079	0.065 ± 0.094
5145-Pholus	Centaur	41	7.165 ± 0.076	48.354 ± 1.930	1.261 ± 0.121	0.788 ± 0.036	0.822 ± 0.054	2.612 ± 0.048	0.391 ± 0.047	-0.037 ± 0.047
12917-1998 TG ₁₆	J.Trojan	2	11.388 ± 0.073	11.839 ± 1.962	0.724 ± 0.042	0.537 ± 0.042	0.410 ± 0.069	1.707 ± 0.066	0.475 ± 0.086	0.205 ± 0.081
15760-1992 QB ₁	Classic Cold	5	6.979 ± 0.094	27.538 ± 4.806	0.869 ± 0.143	0.707 ± 0.093	0.651 ± 0.166	_	_	_
15809-1994 JS	Res.(other)	2	7.479 ± 0.160	28.194 ± 6.000	_	0.760 ± 0.170	0.480 ± 0.149	_	0.460 ± 0.705	_
19308-1996 TO ₆₆	Classic Hot	16	4.520 ± 0.042	2.077 ± 2.168	0.671 ± 0.057	0.389 ± 0.043	0.356 ± 0.053	0.997 ± 0.101	_	_
20000-Varuna	Classic Hot	8	3.455 ± 0.090	26.843 ± 1.917	0.906 ± 0.052	0.637 ± 0.040	0.628 ± 0.040	2.010 ± 0.050	0.564 ± 0.070	-0.038 ± 0.105
20161-1996 TR ₆₆	Res.(other)	1	_	_	_	_	_	_	0.470 ± 0.296	_
24835-1995 SM ₅₅	Classic Hot	10	4.332 ± 0.040	0.272 ± 1.805	0.652 ± 0.032	0.357 ± 0.043	0.356 ± 0.052	1.010 ± 0.050	-0.270 ± 0.163	_
35671-1998 SN ₁₆₅	Classic Cold	6	5.679 ± 0.320	6.857 ± 3.068	0.712 ± 0.095	0.444 ± 0.078	0.437 ± 0.083	1.270 ± 0.050	_	_
42355-Typhon	Scattered	12	7.252 ± 0.054	12.657 ± 1.354	0.758 ± 0.039	0.525 ± 0.022	0.414 ± 0.053	1.560 ± 0.045	0.406 ± 0.087	0.160 ± 0.071
42355-Typhon+B	Scattered	1	_	9.763 ± 0.000	_	_	_	_	_	_
50000-Quaoar	Classic Hot	4	2.220 ± 0.029	29.224 ± 1.706	0.958 ± 0.035	0.650 ± 0.020	0.610 ± 0.028	2.180 ± 0.058	0.360 ± 0.050	0.030 ± 0.057
52747-1998 HM ₁₅₁	Classic Cold	1	7.417 ± 0.100	24.891 ± 4.691	0.930 ± 0.090	0.620 ± 0.050		_		_
90377-Sedna	Detached	3	1.077 ± 0.065	32.954 ± 2.972	1.131 ± 0.079	0.686 ± 0.077	0.657 ± 0.067	2.320 ± 0.060	0.290 ± 0.222	0.050 ± 0.314
90482-Orcus	Res. 3:2	6	1.982 ± 0.099	2.761 ± 1.831	0.664 ± 0.041	0.370 ± 0.039	0.390 ± 0.045	1.070 ± 0.042	0.120 ± 0.051	0.053 ± 0.055
134340-Pluto	Res. 3:2	4	-0.881 ± 0.400	7.607 ± 0.681	0.867 ± 0.016	0.515 ± 0.035	0.400 ± 0.010	_		_
134340-Pluto+B	Res. 3:2	1	_	3.334 ± 0.000	0.710 ± 0.002	_		_		_
134340-Pluto+C	Res. 3:2	1	_	-2.234 ± 0.000	0.644 ± 0.028	_		_		_
134340-Pluto+D	Res. 3:2	1	_	18.074 ± 0.000	0.907 ± 0.031	_	_	_	_	_
136108-Haumea	Classic Hot	3	0.217 ± 0.030	-0.010 ± 0.850	0.631 ± 0.025	0.370 ± 0.020	0.320 ± 0.020	1.051 ± 0.020	-0.044 ± 0.037	-0.111 ± 0.048
136199-Eris	Detached	64	-1.462 ± 0.036	3.866 ± 0.823	0.805 ± 0.015	0.389 ± 0.049	0.363 ± 0.061	0.849 ± 0.108	0.080 ± 0.072	-0.280 ± 0.085
136472-Makemake	Classic Hot	1	_	4.693 ± 0.900	0.828 ± 0.022	_	_	_	_	_

(1) Classes refer to Gladman et al, SSBN08. M11 is the absolute R magnitude; (2) Number of epochs; (3) Grt is the spectral gradient S.

Table 3. For some objects, list of the references used, and number of epochs. The full table is available in electronic form at the CDS, and maintained up up-to-date on the MBOSS colour web site.

Object	References	Nr.
1994 ES ₂	Green et al. (1997)	1
7066-Nessus	Davies et al. (1998)	
	Romanishin & Tegler (1999)	
	Bauer et al. (2003)	
	Davies (2000)	
	Tegler & Romanishin (1998)	17
15760-1992 QB ₁	Jewitt & Luu (2001)	
	Tegler & Romanishin (2000)	
	Boehnhardt et al. (2001)	
	Benecchi et al. (2011)	
	Romanishin et al. (1997)	5
19308-1996 TO ₆₆	Jewitt et al. (2007)	
	Jewitt & Luu (1998)	
	Romanishin & Tegler (1999)	
	Hainaut et al. (2000)	
	Gil-Hutton & Licandro (2001)	
	Davies et al. (2000)	
	Jewitt & Luu (2001)	
	Boehnhardt et al. (2001)	
	Sheppard (2010)	
	Barucci et al. (1999)	
	Davies (2000)	
	Tegler & Romanishin (1998)	16
	-	

Table 3 shows examples of the list of references per object that were included in this study; the information for all objects is given in the online supplement.

Figure 1 presents some of the representative colour-colour diagrams. As a reference, the solar colours are indicated by a red star (see paper I, Table 2 for a list of the solar colours and their references). The red "reddening line", introduced in paper I, marks the locus of objects with a perfectly linear reflectivity spectrum over the considered colours, with a tick mark every 10 units of S. For colour diagrams in the visible range, the distance from a data point to the reddening line therefore indicates a "bend" in the reflectivity spectrum. The symbols used indicate the dynamical class, and are explained in Fig. 2. Figure 3 shows examples of the coarse reflectivity spectra for a set of objects obtained from their filter photometry. All the other plots are available at the MBOSS site.

Figure 7 displays the spectral gradient S as a function of the main orbital elements, the orbital excitation, and the M(1,1) magnitude of the objects. Other similar plots, for all the colours, are available online. The symbols indicate the dynamical class.

Figure 4 shows the histogram and the cumulative distribution for an example colour and the spectral gradient. Other similar plots are available on-line.

3. Results from the statistical analysis of the enhanced MBOSS database

3.1. Statistical tools

Comparing the distribution by comparing their histograms "by eye" is unreliable: the size of the bins can cause artifacts or hide real features, and what appears as a strong difference can actually be of no significance. Similarly, the eye is a very powerful tool to detect alignments and clustering, even when these are insignificant.

The colour, spectral gradient and M(1, 1) distributions of the various MBOSS populations were therefore compared using a set of simple statistical tests that quantifies the significance of the apparent differences, i.e. the t-test, the f-test, and the Kolmogorov-Smirnov test. The tests are described in detail in Appendix B of paper I and references therein and can be briefly summarized as follows:

- The Student t-test indicates whether the mean values of the two distributions are statistically different. The implementation used here deals properly with distributions with different variances.
- The f-test considers whether two distributions have significantly different variances.
- Finally, the Kolmogorov-Smirnov (KS) test uses all of the information contained in the distributions (and not just their means and variances) to estimate whether they are different.

For these tests to give meaningful results, the samples compared must be sufficiently large. We set the threshold at 15 units for *t*- and *f*-tests, and at 20 units for the KS test.




Fig. 1. Selected colour-colour diagrams of the objects. The physico-dynamical class of the objects is identified by their symbols, which are explained in Fig. 2. The red star indicates the solar colours, and the red line is the locus of objects with a flat reflection spectrum, with a small mark every 10%/100nm. The other colour combinations are available from the MBOSS site.

3.2. The case of the brightest objects

Large KBOs have atypical surface properties, e.g high albedo (Stansberry et al. 2008; Brown 2008), flatter spectral gradients, and CH_4 ice-dominated spectra (Brown et al. 2005). For instance, Pluto is known to have a tenuous but captured atmosphere that can be re-deposited on the surface (Stern & Tholen 1998; Protopapa et al. 2008), while (136199) Eris has an extremely high albedo (Sicardy et al. 2011) and a spectrum dominated by CH_4 ice (Brown et al. 2005) and a nitrogen ice signature (Dumas et al. 2007). Schaller (2010) discusses the presence of volatiles (nitrogen, methane, and carbon monoxide) on bodies large enough to retain an atmosphere. The transition from small objects with volatile-free surfaces (such as Charon or Quaoar) to large objects with volatile-rich ones (such as Pluto) can be explained with a simple model of atmospheric escape formulated by Schaller & Brown (2007).

We therefore decided to remove the largest objects (using their intrinsic brightness as a proxy) for some studies, so that their potentially different characteristics do not pollute the colour distributions of the other, "normal" MBOSSes. To select a cut-off value,

- A Resonant 3:2 (Plutinos)
- △ Resonant (others)
- Cold classic disk objects
- Hot classic disk objects
- Scattered disk objects
- Detached disk objects
- _ Jupiter Trojans
- Centaurs
- Short-period comets
- Long-period comets
- ☆ Sun

Fig. 2. Legend of the symbols used thorough this paper.




Fig. 3. Examples of reflectivity spectra for a set of objects. The reflectivity is normalized to unity for the *V* filter, and the spectra are shifted for clarity. The physico-dynamical classes of the objects are identified by their symbols, which are explained in Fig. 2. The spectra for all the objects are available online on the MBOSS site.

we estimated the radius of the objects whose escape velocity $v_{\rm esc}$ is equal to the velocity of material ejection via cometary activity $v_{\rm ej}$. The escape velocity is given by

$$v_{\rm esc} = \sqrt{\frac{2GM}{R}},\tag{3}$$

where G is the gravitational constant, M the mass of the object, and R its radius. Assuming a density $\rho \sim 1000 \mathrm{kg} \ \mathrm{m}^{-1}$ to be able to infer the mass from the volume of the object,

$$v_{\rm esc} \sim 7.5 \ 10^{-4} R$$
 (4)

Table 4. Averge colours and dispersions of all MBOSSes (excluding Trojans), comparing those with M(1,1) < 5 and the others.

Colour	N	Aver./ σ	N	Aver./ σ	t-Prob	f-Prob
	M(1,1) < 5		N	$I(1,1) \ge 5$		
J-H	22	0.27 ± 0.25	84	0.41 ± 0.20	0.025	0.149
H - K	16	0.04 ± 0.07	31	0.07 ± 0.12	0.279	0.036

Notes: The q cut-off is set at the median value; N is the number of measurements; t-Prob and f-Prob are the probabilities that the two sub-samples are randomly extracted from the same distribution, as evaluated with the t- and f-tests. The other colours show insignificant differences.

in IS units. To estimate the velocity of material ejection, v_{ei} (in m s⁻¹), we used the relation

$$v_{\rm ej} = 580 \ r^{-0.5} \sqrt{\frac{\mu_0}{\mu}},$$
 (5)

where r is the heliocentric distance in AU, and μ_o/μ the ratio of the molecular mass of the species driving the activity to that of water. This relation was obtained by measuring the expansion of cometary comae, and is supported by a theoretical analysis —see Delsemme (1982) for a discussion. This velocity is the terminal velocity of the gas in the case of a small comet, and it is directly controlled by the thermal velocity of the sublimating gas. For the TNOs, we used $\mu_o/\mu = 0.64$ for CO. By equating $\nu_{\rm ej}$ to $\nu_{\rm esc}$, we obtained a critical radius R_c above which an object is likely to retain some of the material ejected by cometary activity

$$where R_c \sim \frac{6.2 \ 10^5}{\sqrt{r}}.\tag{6}$$

 R_c is in m and r in AU. Delsanti et al. (2004) discuss the evolution of R_c with different species and distance. For our purpose, it is enough to say that $R_c \sim 150$ km at r = 17 AU and 100 km at 43 AU. Stansberry et al. (2008), based on their large set of measurements with the Spitzer Space Telescope, indicate that large TNOs have higher than average albedos. We used p = 0.2 to convert R_c in an absolute magnitude, leading to

$$M(1,1) \sim 1.9 + 2.5 \log r,$$
 (7)

again with r in AU. This gives $M(1, 1) \sim 5$ at r = 17 AU, and 6 at 43 AU. In what follows, we took M(1, 1) = 5 as a conservative limit, below which objects are likely to keep at least part of their atmospheres in the Centaur and TN regions, thus be potentially affected by different resurfacing processes. This choice is partly arbitrary. We performed the following analysis with different cut-off values, leading to similar results.

To verify whether we were justified in separating the bright M(1,1) objects from the others, we compared their colours and gradients with those of the faint objects. The Trojans were not considered in this test, whose results are summarized in Table 4.

With the exception of the J-H colour, the bright and the faint objects have compatible mean colours and overall distribution. In the infrared, the bright objects have marginally bluer colours $(J - H = 0.27 \pm 0.25)$ than the faint ones $((J - H = 0.41 \pm 0.20)$; this has only a probability of 2.5% of occurring by chance, i.e. this difference is marginally significant. Similarly, the bright objects have a slightly broader distribution of H - K colour than the fainter ones (at the 3% level). When performing these tests on individual or groups of classes, the result was a slightly more significant for the resonant objects, but could not be tested for the other groups, which did not have enough bright objects. Nevertheless, removing the non-resonant objects increased the significance of the result for the resonant objects. Additionally, one must consider the possibility that the larger dispersion in H - K for the fainter objects is connected to the lower signal-to-noise ratio in K, as the sky is much brighter in this band and the instruments tend to be less sensitive there. Considering the colours of the various dynamical classes in the visible wavelength range for both the bright (M(1, 1) < 5)and faint objects, the colour distributions of the bright objects were found to be indistinguishable from those of the fainter ones, indicating that the object populations might be rather uniform despite some larger objects possibly having some different surface properties (for instance, albedo) and intrinsic activity possibly playing a role in resurfacing the bodies. We note, however, that the NIR colours of individual KBOs can be affected if a significant fraction of their surfaces is covered with ices which display strong absorptions (such as CH_4), particularly in H and K bands. For example, the colours of Pluto, Eris and 2005 FY_9 (see Brown 2008) show this effect. The presence of these ice absorption lines can be inferred from the unusual negative slope of the NIR spectral gradients (i.e. from *J* to *H* and *K* bands).

In summary, the infrared colours of the objects with bright absolute magnitudes differ slightly from those of the fainter objects. We therefore separated them for most colour tests, but we also re-incorporated them into some tests of the visible colours where more objects were helpful.

On the basis of these results, one might consider including the intrinsically brighter objects in the remainder of the study. However, we decided not to: if they were indeed similar, they would only marginally increase the size of the sample, without changing our conclusions. If they eventually turned out to be different, as possibly suggested by the NIR colours, we did the right thing in keeping them separate.

3.3. Global characteristics of individual classes

The MBOSS database that we analyse contains photometric measurements of ten different dynamical classes (see Fig. 2), i.e. the SPCs, the LPCs, the Jupiter Trojans, Centaurs, Kuiper Belt objects in 3:2 (Plutinos = Res. 3:2) and in other orbital resonances




Fig. 4. R - I (top) and spectral gradient (bottom) cumulative distributions (left) and histogram (right). Solar value is indicated by a dotted line. Similar plots are available on-line for the other colours.

(Res.others) with Neptune, the classical disk objects (CDOs) in two flavors of dynamically 'hot' and 'cold' CDOs, the scattered disk objects (SDOs) and the detached disk objects (DDOs).

Selected colour-colour diagrams of MBOSSes are shown in Fig. 1. Colour distributions of the various classes of objects are displayed in Fig. 4, for a selection of colours and spectral gradients S. At first sight, the ranges of visible colours are similar for the several dynamical groups (Plutinos and resonants, CDOs, SDOs, and Centaurs), ranging from slightly bluish (-10%/100 nm) compared to the Sun to very red (55%/100 nm). Jupiter Trojans do not appear to contain any very red objects (i.e. with spectral gradients > 20%/100 nm) as the other groups do. For LPCs, the number of measured objects (14) may be too small to provide representative results for the total population, although we note that the available data indicate that the spectral gradients are between about 0 and 10 %/100nm, in close agreement with the other groups.

The mean colours (and corresponding variances) for the various dynamical classes are listed in Table 7, top panel, using all data in the MBOSS2 database, and the values for the restricted data-set of objects with $M(1, 1) \ge 5$ mag are in Table 7, bottom panel. Many of these means are based on only a small number of objects and should be interpreted with care.

The average characteristics of the dynamical classes are also displayed as reflectivity spectra in Fig. 5. The red slope in the visible wavelength range, although different for individual groups, levels off in the NIR with transition between the I and J or H bands.




Fig. 5. Average reflectivity spectra for the different physico-dynamical classes. a: small objects (M(1, 1) > 5 mag), b: large objects only (M(1, 1) < 5 mag), c: all objects

3.4. Comparisons between classes

In our comparison, we considered the photometric properties of the dynamical classes all groups except LPCs, since are too sparse to expect firm conclusions.

3.4.1. Colours and spectral gradients

Comparing the colour distributions of the different physico-dynamical classes, we can make the following observations (see Fig. 4):

- The distributions of both the colours and the spectral gradients in the visible wavelength range differ among the groups, in terms of their extent, shape and peak location. Taking the spectral gradients as an example, the peak level increases starting with the Jupiter Trojans (5-10%/100 nm), and continuing to the SPCs, Centaurs, DDOs, SDOs, Plutinos (and possibly the resonant objects) (10-20%/100 nm), and the hot CDOs (20-30%/100 nm), before ending with the cold CDOs which are the reddest objects (25-35%/100 nm) among the MBOSSes. The colour ranges and distribution width can clearly be inferred from the cumulative distribution functions (see left panels of Fig. 4).
- Relevant secondary peaks in the frequency distributions may exist for Centaurs and Plutinos at higher reddening. At least for Centaurs, statistical arguments for a bimodal surface colour distribution have been presented in numerous papers (eg Tegler & Romanishin 1998, 2003; Peixinho et al. 2003; Delsanti et al. 2006; Tegler et al. 2008). Tegler et al. (2008) presented a

Table 5. Statistical comparisons of the spectral gradient distributions of pairs of MBOSS populations: t-test, f-test, and KS test.

			Resonant	Classical	Classical	Scattered	Detached	Centaurs	Short	Long	Trojans
			(others)	cold	hot	disk	disk		period	period	
						objects	objects		comets	comets	
N			15	43	28	23	17	26	44	0	60
Resonant	31	t	0.701	0.002	0.713	0.340	0.056	0.768	0.006		0.000
3:2		f	0.367	0.209	0.505	0.777	0.264	0.163	0.334		0.000
		KS	0.810	0.000	0.817	0.612	0.023	0.438	0.072		0.000
			Indist.	Not comp.	Indist.	Indist.	Marginal	Indist.	Marginal	-/-	Not comp.
Resonant	15	t		0.019	0.485	0.211	0.034	0.968	0.005		0.000
others		f		0.986	0.166	0.523	0.881	0.057	0.810		0.000
		KS		0.019	0.680	0.183	0.009	0.101	0.020		0.000
				Marginal	Indist.	Indist.	Marginal	Indist.	Marginal	-/-	Not comp.
Classical	43	t		-	0.002	0.000	0.000	0.038	0.000		0.000
Cold		f		•	0.051	0.402	0.839	0.006	0.747		0.000
		KS			0.006	0.000	0.000	0.000	0.000		0.000
						Not comp.	Not comp.	Not comp.	Not comp.	-/-	Not comp.
Classic	28	t				0.602	0.160	0.560	0.039		0.000
Hot		f				0.377	0.106	0.471	0.094		0.000
		KS				0.161	0.002	0.284	0.040		0.000
						Indist.	Marginal	Indist.	Marginal	-/-	Not comp.
Scattered	23	t					0.355	0.286	0.114		0.001
Disk		f					0.406	0.125	0.569		0.000
Objects		KS					0.280	0.358	0.359		0.000
							Indist.	Indist.	Indist.	-/-	Not comp.
Detached	17	t						0.064	0.583		0.020
Disk		f						0.031	0.662		0.000
Objects		KS						0.182	0.444		0.000
Ū								Indist.	Indist.	-/-	Not comp.
Centaurs	26	t							0.015		0.000
		f						•	0.014		0.000
		KS							0.035		0.000
									Marginal	-/-	Not comp.
Short	44	t									0.090
Period		f						•			0.000
Comets		ĸs									0.000
										-/-	Not comp.

Notes: The three tests are the Student t- and f-tests and the KS test. The number N indicates how many objects have measurements. The result is the probability that the two distributions are randomly extracted from the same population. The label indicates whether the populations are significantly different (i.e., not compatible), marginally different, or statistically indistinguishable (see text for details).

detailed analysis of the Centaurs' bimodality, based on a sample of 26 objects, i.e. almost as large as the one presented here (29 objects, so we do not present a new analysis). They concluded that the B - R distribution is bimodal with a confidence level of 99.5%, with 10 red Centaurs and 16 gray ones. The red ones have marginally smaller orbital inclinations, and higher albedos at the 99% confidence level.

- The frequency distribution in J - H instead gives a rather uniform picture for all dynamical groups with respect to colour range and peak position (with a few singular exceptions for Plutinos and CDOs).

This global characterization indicates that parameters of the visible spectral energy distribution are better diagnostics of both the global surface reflectivity and differences among MBOSSes than the NIR colours. However, we note that the NIR spectral data of MBOSSes is more sensitive to compositional differences than visible one, because stronger absorption bands, in particular of icy compounds, are found in the NIR wavelength domain (Barucci et al. 2008).

Table 6. Statistical comparisons of the J-H colour distributions of pairs of MBOSS populations: t-test, f-test, and KS test.

			Classic	Classic	Centaurs
			cold	hot	
	N		28	34	17
Resonant	21	t	0.498	0.119	0.308
3:2		f	0.000	0.961	0.000
		KS	0.449	0.677	0.984
			Not comp.	Indist.	Not comp.
Classic	28	t		0.121	0.332
cold		f	-	0.000	0.703
		KS		0.710	0.284
				Not comp.	Indist.
Classic	34	t		_	0.271
hot		f	-	-	0.000
		KS			0.499
					Not comp.

Notes: The three tests are the Student t- and f-tests and the KS test. The N number indicates how many objects have measurements. The classes that are not listed do not have sufficient measurements for these tests to be performed. The result is the probability that the two distributions are randomly extracted from the same population. The label indicates whether the populations are significantly different, marginally different, or statistically indistinguishable (see text for details).

Table 7. Mean colours and variances, for the MBOSS populations: all objects

colour	Res. 3:2	Res.others	Classic cold	Classic hot	Scattered	Detached	J.Trojans	Centaurs	SPCs	LPCs
Full data	base						-			
U-B	0	2	3	11	7	2	0	0	0	0
		0.273 ± 0.090	0.593 ± 0.123	0.258 ± 0.168	0.334 ± 0.144	0.180 ± 0.156				
B-V	38	16	41	38	27	22	74	28	17	1
	0.868 ± 0.170	0.975 ± 0.142	0.996 ± 0.140	0.864 ± 0.163	0.858 ± 0.153	0.869 ± 0.109	0.777 ± 0.091	0.892 ± 0.195	0.863 ± 0.209	0.810 ± 0.000
V - R	38	21	49	41	25	23	80	30	43	2
	0.558 ± 0.101	0.603 ± 0.109	0.630 ± 0.086	0.510 ± 0.145	0.536 ± 0.096	0.519 ± 0.083	0.445 ± 0.048	0.567 ± 0.131	0.494 ± 0.105	0.422 ± 0.045
V-I	40	23	73	51	28	24	80	26	22	1
	1.101 ± 0.202	1.181±0.164	1.220±0.159	1.010 ± 0.249	1.037 ± 0.189	1.039 ± 0.170	0.861 ± 0.090	1.077 ± 0.252	0.947 ± 0.112	0.878 ± 0.000
V - J	16	5	5	13	9	4	12	10	1	0
	1.838 ± 0.493	2.138 ± 0.327	1.960 ± 0.421	1.509 ± 0.431	1.753 ± 0.481	1.614±0.601	1.551 ± 0.120	1.769 ± 0.498	1.630 ± 0.000	
R-I	36	19	36	35	22	21	80	26	23	1
	0.536 ± 0.122	0.554 ± 0.090	0.588 ± 0.095	0.506 ± 0.139	0.544 ± 0.108	0.515±0.099	0.416 ± 0.057	0.525 ± 0.151	0.447 ± 0.101	0.430 ± 0.000
J-H	21	13	28	34	13	10	12	17	1	0
	0.442 ± 0.285	0.437 ± 0.065	0.398 ± 0.082	0.316±0.290	0.400 ± 0.124	0.381±0.126	0.434 ± 0.064	0.375 ± 0.075	0.360 ± 0.000	
H - K	11	4	4	8	8	4	12	15	1	0
	0.043 ± 0.059	0.052 ± 0.023	0.079 ± 0.030	0.077 ± 0.117	0.060 ± 0.151	0.035 ± 0.228	0.139 ± 0.041	0.085 ± 0.143	0.140 ± 0.000	
Grt	45	25	82	56	29	26	80	30	43	2
	18.633±10.659	23.471±9.362	25.515±9.217	15.387±11.887	15.258±9.974	15.459 ± 8.873	7.024 ± 3.786	19.320±13.842	11.483±8.307	5.188 ± 3.692
M(1,1)	38	21	46	45	24	23	60	29	134	13
	6.680±1.981	6.235 ± 1.393	6.600 ± 0.614	5.928±1.703	7.168 ± 1.225	5.718 ± 2.181	11.356±1.057	9.456±1.654	16.131±1.667	13.550 ± 3.601

Table 7. Mean colours and variances, for the MBOSS populations, continued: Objects with M(1, 1) > 5 mag

colour	Res. 3:2	Res.others	Classic cold	Classic hot	Scattered	Detached	J.Trojans	Centaurs	SPCs	LPCs
Objects v	with $M(1,1) \ge 5$ m	ag								
U - B	0	1	2	3	7	2	0	0	0	0
		0.210 ± 0.000	0.525 ± 0.049	0.187 ± 0.103	0.334 ± 0.144	0.180 ± 0.156				
B-V	26	10	34	22	23	15	60	28	16	1
	0.858 ± 0.175	0.943 ± 0.131	1.011±0.139	0.919 ± 0.167	0.876 ± 0.143	0.836 ± 0.084	0.789 ± 0.092	0.892 ± 0.195	0.872 ± 0.213	0.810 ± 0.000
V - R	31	15	42	29	23	16	60	30	43	2
	0.555 ± 0.102	0.590 ± 0.102	0.642 ± 0.082	0.523 ± 0.152	0.532 ± 0.098	0.513 ± 0.077	0.461 ± 0.040	0.567 ± 0.131	0.494 ± 0.105	0.422 ± 0.045
V-I	31	13	37	25	22	15	60	26	22	1
	1.098 ± 0.205	1.114±0.162	1.226 ± 0.145	1.050 ± 0.258	1.068 ± 0.186	1.010±0.159	0.898 ± 0.062	1.077 ± 0.252	0.947 ± 0.112	0.878 ± 0.000
V - J	9	1	4	2	6	1	12	10	1	0
	1.912±0.528	2.295 ± 0.000	1.932 ± 0.480	1.807 ± 0.024	1.796±0.369	1.646 ± 0.000	1.551±0.120	1.769 ± 0.498	1.630 ± 0.000	
R-I	30	13	33	25	20	14	60	26	23	1
	0.538 ± 0.127	0.521 ± 0.084	0.578 ± 0.090	0.529 ± 0.137	0.549 ± 0.113	0.510 ± 0.104	0.437 ± 0.044	0.525 ± 0.151	0.447 ± 0.101	0.430 ± 0.000
J-H	14	5	17	17	9	5	12	17	1	0
	0.518 ± 0.319	0.467 ± 0.064	0.401 ± 0.071	0.342 ± 0.296	0.417 ± 0.133	0.414 ± 0.026	0.434 ± 0.064	0.375 ± 0.075	0.360 ± 0.000	
H - K	6	1	3	0	5	1	12	15	1	0
	0.056 ± 0.076	0.080 ± 0.000	0.070 ± 0.030		-0.004 ± 0.114	0.260 ± 0.000	0.139 ± 0.041	0.085 ± 0.143	0.140 ± 0.000	
Grt	31	15	43	28	23	17	60	30	43	2
	19.051±10.558	20.240±8.496	26.488 ± 8.567	17.795±12.258	16.526±10.121	13.971±8.172	8.538 ± 2.759	19.320±13.842	11.483±8.307	5.188 ± 3.692
M(1,1)	32	15	45	34	23	17	60	29	134	13
	7.392 ± 0.802	6.974 ± 0.795	6.635 ± 0.571	6.717 ± 0.833	7.275 ± 1.124	6.746 ± 0.572	11.356±1.057	9.456 ± 1.654	16.131±1.667	13.550 ± 3.601

Notes: The number of objects included in each average is indicated. Grt is the spectral gradient S. M(1, 1) is the R absolute magnitude.

We now compare the populations using the statistical tests described in section 3.1. First of all, we stress that a common feature of all the statistical tests applied is the estimate of a probability that the two populations compared are *not* randomly extracted from the same population. In other words, the statistical tests can very firmly establish whether two populations significantly differ. A contrario, no statistical test can prove that two populations are identical. For instance, the colour distributions of the Centaurs and Jupiter Trojans (see Tab. 5) are radically different with a very high significance level: it is totally improbable that these two populations were randomly extracted from the same reservoir. However, the spectral gradient distributions of SDOs and DDOs indicate that these objects could have been randomly extracted from a same pool, even though this may be just a coincidence.

Table 5 summarizes the results of the comparison between the spectral gradient S distribution of the MBOSS populations using these tests. Following the discussion in Section 3.2, the tests were restricted to the objects with M(1,1) > 5 mag. For each pair, the table lists the probabilities that the two classes were randomly extracted from a common population, based on the mean spectral gradient (t-test), the spectral gradient variance (f-test), and the overall spectral gradient distribution (KS test). Low probabilities indicate that the two considered distributions are incompatible. The pairs for which at least one of the tests returns a probability < 0.001 were flagged as "not compatible", indicating they are significantly different. Those where a test indicates a probability < 0.05 are marked as marginally different, suggesting that there is a difference, but that it is not very strongly significant. The others are statistically indistinguishable using these tests, which again does not mean that the objects are similar, just that these tests cannot prove they are different. Similar tests were performed on the individual colours, with similar results. We discuss here only the spectral gradients, as they encompass most of the information present in the visible spectrum range.

In the infrared, the tests are possible only in J - H, and only some of the classes have enough measurements for the tests to be meaningful. The results are presented in Table 6. The statistical tests indicate the following results:

- The spectral gradient distributions of the various resonant classes are statistically indistinguishable. To increase the size of the sample, all the resonant objects were divided only between Plutinos (3:2) and "other resonant objects".
- The spectral gradient distributions of Plutinos and other resonant objects, SDOs, hot CDOs, and Centaurs are indistinguishable.
- Additionally, the Plutinos and the hot CDOs had indistinguishable J H distributions.
- The spectral gradient distribution of DDOs is indistinguishable from that of SDOs and only marginally distinguishable from those of Plutinos and resonant objects, as well as hot CDOs.
- The spectral gradient distribution of cold CDOs is incompatible with the ones of hot CDOs, Plutinos, SDOs, DDOs, and Centaurs, and is only marginally compatible with that of the resonant objects.
- Additionally, the cold CDOs J H distribution is incompatible with that of the Plutinos and the hot CDOs.
- The spectral gradient distribution of Jupiter Trojans clearly indicates its complete incompatibility with all other MBOSS populations.
- The spectral gradient distribution of SPCs significantly differs from those of the Jupiter Trojans and cold CDOs. It is only
 marginally different from those of the hot CDOs, Plutinos, and other resonant objects as well as Centaurs, and it cannot be
 distinguished from that of SDOs and DDOs.
- The J-H colours of the Centaurs are incompatible with those of the Plutinos and the hot CDOs.

In conclusion, Plutinos, other resonant objects, hot CDOs, SDOs, and DDOs may have rather similar spectral gradients in the visible, while cold CDOs and Jupiter Trojans form two separate groups of spectral properties in the visible wavelength range. The peak in the spectral gradient distribution of hot CDOs seems to be shifted slightly to higher reddening, although for the moment this result is not yet at a statistically significant level. Short-period comets may agree with the first populations mentioned above.

3.4.2. Absolute magnitude

The absolute magnitude M(1,1) of an object depends on the reflectivity function across the surface, i.e. a first order approximation on the object size, the geometric albedo, and the photometric phase function. Given the relationship between M(1,1), size, albedo, and phase function, one can assume that the M(1,1) distribution is dominated by the effects of the size, except for the largest bodies, which often have high albedos (Stansberry et al. 2008). The solar phase angle for KBOs is always relatively small, and the phase function is moderately steep, so the effect on M(1,1) hardly exceeds 0.2-0.3mag. Hence, we assume that the M(1,1) distribution of KBOs reflects –at least to zeroth order– the size distribution of the objects, with the understatement that a detailed analysis requires more accurate size estimates of the objects. Such a careful study can be found in Stansberry et al. (2008).

The M(1,1) distributions are strongly biased by cumulative selection effects: while there are few bright and large objects in a given class, they are more easy to discover, and also more easy to observe, so they are likely to be represented in this data-set. A contrario, the discovery surveys are incomplete for fainter objects, and faint objects are less likely to be picked by observers performing colour measurements. Consequently, the M(1,1) distributions are likely to be fairly complete at the bright end, but have a complex non-completeness at the faint end. Meech et al. (2004) discussed and simulated many of these selection effects.

Fig. 6 shows the frequency histogram and cumulative distribution function of M(1, 1) for the various MBOSS groups, now covering all objects in the MBOSS database (i.e. not excluding those with M(1, 1) > 5 mag).

The selection bias against small KBOs is clear, with no object fainter than $M(1,1) \sim 8.5$ in the distant classes – this corresponds to actual magnitudes fainter than 23.5 for Plutinos, and even fainter for CDOs, i.e. about the limiting magnitude for colour observations performed with fairly short exposures on a 8 meter class telescope. On the other hand, Centaurs and SPCs may be seen as a sample of smaller KBOs that is scattered towards the inner planetary system. Meech et al. (2004) found that the shape of the SPC size distribution is incompatible with that of the other TNOs.

The slope k of the logarithmic cumulative M(1,1) distribution relates to the power law q of the cumulative radius distribution $N_c(a) \propto a^{-q}$ as q = 5k. Measuring the k slopes on the straight parts of the distributions in Fig. 6 leads to values $q \sim 0.8$ for the SPCs,




Fig. 6. R absolute mag nitude M(1, 1) cumulative distributions (logarithmic, a), and histogram (b).

the Centaurs, the faint SDOs, the DDOs, the hot CDOs, and the resonant objects; the steep slope of the cold CDOs and the bright SDOs corresponds to $q \sim 2$. Only the steepest part of the SPCs distribution reaches $q \sim 3$. Analyses based on controlled samples designed for size analyses have found much steeper power-laws of from q = 3.0 (Trujillo et al. 2001) to 3.5 (Kenyon & Bromley 2004), i.e. distributions much richer in small/faint objects. This confirms that the sample used for colour measurements is heavily biased against faint objects —the distributions may be fairly complete only for the brightest SPCs.

Therefore, the size distributions in Fig. 6 should only be used for internal comparison, with the hope that the selection effects affected them in a similar way, and/or restricting the analysis to the bright end of the distributions.

The observed Jupiter Trojans (M(1,1)) between 8 and 13 mag) are clearly smaller than KBOs, which cover the M(1,1) range from about 0 to 9 mag. The bulk of that shift is caused by the proximity of the Trojans, which enables objects fainter by ~ 4 mag to be observed.

It is noteworthy that cold CDOs display a rather peaked M(1,1) distribution and do not contain objects with M(1,1) brighter than about 5 mag, while Plutinos, hot CDOs, and DDOs contain objects of M(1,1) from 2 to -1 mag. The rather steep slope in the cumulative distribution function of M(1,1) of the cold CDOs reflects the likely difference in the size range of these KBOs. In particular, it differs from that of hot CDOs, which should be exposed to very similar selection effects, i.e. it is real. The brightest




Fig. 7. The spectral gradient of the objects [%/100 nm] as a function of the main orbital elements, the orbital excitation \mathcal{E} , and the absolute R magnitude M(1,1). Similar plots are available on-line for the various colours.

resonant objects reach M(1, 1) of about 3mag, while the SDOs are more similar in brightness to cold CDOs, despite having a much less pronounced peak level on a slightly wider distribution plateau.

From the M(1,1) distributions of MBOSSes, we conclude that the absence of larger bodies among the cold CDOs and their presence among other populations of the Kuiper Belt may suggest different formation environments for both groups of objects that may have favored the growth of larger bodies for Plutinos, SDOs, DDOs, and possibly hot CDOs compared to cold CDOs. Jupiter Trojans are a population of smaller size bodies, again arguing for a formation environment and evolution scenario that were distinct from that of the other MBOSSes (now including cold CDOs). Since it is very likely that the large bodies in these groups may have survived impact events widely unaffected over the lifetime of the solar system, one can assume that they represent the original population of planetesimals. The Centaurs reach the magnitude/size range where the vast majority of larger KBOs are detected, but they do not include one of the rare big ones. This might well be compatible with the scattering scenario for Centaurs but needs quantitive confirmation by dynamical calculations.

Altogether, the M(1, 1) distributions give the impression that the MBOSSes may have been formed in at least three (maybe four) different environments, i.e. one for the Jupiter Trojans, one for the cold CDOs, maybe one for the hot CDOs, and one for the rest of the bodies. Since KBOs now reside and for a very long time resided in the same distance range, one may consider Plutinos, resonant objects, hot CDOs, and SDOs as immigrant population of the Kuiper Belt compared to the cold CDOs that may represent "aborigines KBOs". Interestingly, the DDOs also appear to be 'sizewise' in accord with the Kuiper Belt "immigrants", although they are found at a much more distant location.

3.5. Dependences on dynamical parameters

The colour distribution (say, the B-V distribution) of a class of objects (say, the Centaurs) is considered as a function of another parameter of the objects (say, the perihelion distance q) to search for correlations as possible indicators of any physical dependences. The traditional Pearson correlation factor is not robust for non-linear dependencies, so a simpler and more robust estimator was used: the sample is divided into two sub-samples, using the median value of the independent variable (in the example, the median q of the Centaurs, i.e. 16.22 AU). The colour distribution of the two sub-samples are then compared through t-test and t-test and the Pearson correlation parameter is also estimated. Visually, this can be done by comparing the left-hand side with the right-hand side of all the sub-panels of Figure 7.

For cold CDOs: The variances in the colours B-V, B-R and B-I are different for the inner (smaller variance) and the outer (larger variance) population among the cold CDOs (considering both their semi-major axis and perihelion distance). These differences have a good level of significance, as listed in Tab. 8. The effect is only marginally observed in V-R and V-I. The average colours of the sub-samples are indistinguishable: only the breaths of the distributions are different

Since similar results are found neither in other colours nor the spectral gradients in the visible, the effect might be related to a wavelength dependent phenomenon that is mostly restricted to the B filter, for instance a stronger unknown continuum absorber in the outer cold CDOs.

Table 8. Averge colours and dispersions for the cold CDOs, comparing those with smaller a (or q) and those with large values.

Colour	N	Aver./σ	N	Aver./σ	t-Prob	f-Prob
	a	ı < 43.95	a	$a \le 43.95$		
B-V	16	0.99 ± 0.16	18	1.03 ± 0.11	0.148	0.148
B-R	16	1.59 ± 0.26	17	1.69 ± 0.09	0.151	0.000
B-I	9	2.18 ± 0.37	14	2.27 ± 0.15	0.489	0.003
V - R	21	0.63 ± 0.10	21	0.65 ± 0.06	0.310	0.034
V-I	17	1.20 ± 0.16	20	1.24 ± 0.14	0.395	0.538
	9	<i>y</i> < 40.91	q	$t \le 40.91$		
B-V	17	0.95±0.17	17	1.07±0.06	0.013	0.000
B-R	16	1.56 ± 0.25	17	1.72 ± 0.07	0.024	0.000
B-I	13	2.15 ± 0.30	10	2.34 ± 0.13	0.060	0.017
V - R	19	0.61 ± 0.08	23	0.67 ± 0.08	0.033	0.802
V-I	18	1.20 ± 0.18	19	1.25 ± 0.10	0.278	0.010

Notes: The a and q cut-off are set at the median value; N is the number of measurements; t-Prob and f-Prob are the probabilities that the two sub-samples are randomly extracted from the same distribution, evaluated with the t- and f-tests. The other colours show insignificant differences.

Table 9. Average colours and dispersions for the hot CDO, comparing those with smaller q and those with large values.

Colour	N	Aver./σ	N	Aver./σ	t-Prob	f-Prob
	q	< 39.39	q	≤ 39.39		J
B-V	10	0.80 ± 0.12	12	1.02±0.13	0.000	0.795
B-R	10	1.28 ± 0.21	12	1.64 ± 0.18	0.000	0.666
B-I	8	1.71 ± 0.28	9	2.37 ± 0.27	0.001	0.838
V - R	14	0.47 ± 0.14	15	0.57 ± 0.15	0.071	0.792
V - I	12	0.94 ± 0.24	13	1.15 ± 0.24	0.037	0.967

Notes: The q cut-off is set at the median value; N is the number of measurements; t-Prob and f-Prob are the probabilities that the two sub-samples are randomly extracted from the same distribution, evaluated with the t- and f-tests. The other colours show insignificant differences.

Table 10. Average colours and dispersions for the hot CDOs, comparing those with smaller *i* and those with large values.

Colour	N	Aver./ σ	N	Aver./ σ	<i>t</i> -Prob	f-Prob
		i < 17.0		$i \le 17.0$		
B-V	14	0.99±0.16	8	0.80 ± 0.09	0.002	0.096
B-R	14	1.60 ± 0.21	8	1.26 ± 0.20	0.002	0.940
B-I	11	2.14 ± 0.33	6	1.70 ± 0.31	0.020	0.978
V - R	15	0.61 ± 0.08	14	0.43 ± 0.16	0.001	0.012
V-I	13	1.20 ± 0.15	12	0.89 ± 0.26	0.002	0.064
Grad.	15	24.59 ± 7.99	13	9.95±11.80	0.001	0.165

Notes: The q cut-off is set at the median value; N is the number of measurements; t-Prob and f-Prob are the probabilities that the two sub-samples are randomly extracted from the same distribution, evaluated with the t- and f-tests. The other colours show insignificant differences.

For hot CDOs: Hot CDOs with a perihelion closer to the Sun are significantly bluer than the ones that do not get close, see Tab. 9, confirming earlier suggestions by Boehnhardt et al. (2004) based on a smaller object sample. This result is visible in the B - V, B - R, and B - I colours (e.g. $B - V = 0.80 \pm 0.12$ for q < 39.4 AU, and $B - V = 1.02 \pm 0.13$ mag for the others). This result is based on fairly small samples, which decreases the strength of its significance.

Hot CDOs with high inclinations or with large orbital excitation parameters are bluer than the others, see Tab. 10, a finding seen for several colours (B - V, B - R, V - R, V - I) as well as for the spectral gradient in the visible. The sample sizes are small, but sufficient. This result was already known (Trujillo & Brown 2002; Doressoundiram et al. 2008) and is generally expressed as an anti-correlation between reddening of the objects and their orbital energy \mathcal{E} . The usual interpretation scenario is colour changes due to resurfacing by impact ejecta if during a collision between a hot CDO and another KBO subsurface material with different colour properties is ejected and deposited on the surface by impact excavations. Since cold CDOs do not display such an anti-correlation despite of their occupying the same distance range as the hot CDOs do, it may be more an effect of orbital energy, i.e. collision energy rather than collision frequency.

For others: Other combinations of dynamical parameters and colours, spectral gradient, and M(1, 1) data of different dynamical classes do not show any statistically significant signal indicating possible correlations and physical dependencies.

Table 11. Average M(1,1) and dispersions for the hot CDOs, comparing those with blue vs red colours.

Colour	Colour	N	Aver./σ	N	Aver./σ	t-Prob
	cut-off		Blue		Red	
Hot TNO	Os					
B-V	0.92	11	6.92±0.67	11	6.54 ± 0.62	0.183
B-R	1.55	11	7.09 ± 0.45	11	6.37 ± 0.64	0.007
B-I	1.92	8	7.13 ± 0.52	9	6.41 ± 0.67	0.025
V - R	0.56	14	6.88 ± 0.61	15	6.34 ± 0.62	0.026
V-I	1.10	12	6.87 ± 0.61	13	6.29 ± 0.65	0.031
R-I	0.54	12	6.85 ± 0.53	13	6.30 ± 0.72	0.039
Resonan	t objects					
B-V	0.88	18	7.61 ± 0.81	18	6.79 ± 0.78	0.004
B-R	1.40	18	7.61 ± 0.81	18	6.79 ± 0.78	0.004
B-I	1.89	15	7.40 ± 0.84	16	6.79 ± 0.81	0.052
V - R	0.56	23	7.48 ± 0.72	23	7.04 ± 0.88	0.073
V-I	1.06	22	7.42 ± 0.72	22	7.08 ± 0.91	0.170
R-I	0.51	21	7.47 ± 0.78	22	7.01 ± 0.83	0.065

Notes: The objects with M(1, 1) < 5 were excluded from this test; the colour cut-off is set at the median value for the considered colour; N is the number of measurements; t-Prob are the probabilities that the two sub-samples are randomly extracted from the same distibution, evaluated with the t-test. The other colours show insignificant differences.

3.6. Absolute magnitude

3.6.1. Absolute magnitude vs orbital parametres

For the comparison between M(1,1) and dynamical parameters, we consider the small objects $(M(1,1) \ge 5 \text{ mag})$ only and we find that, for Centaurs, SDOs, and DDOs, when considered together, objects with small semi-major axis a and perihelion distance q display a significantly fainter absolute magnitude M(1,1) than the distant ones. While this is a statistically strong result, it disappears when the three classes are considered separately. Thus, it can simply be seen as the selection bias that makes the more distant and thus fainter objects harder to measure than the closer ones.

3.6.2. Absolute magnitude of the blue objects, vs red objects:

We now compare the absolute magnitude M(1, 1) distribution of a class of objects as a function of their colour, to explore whether the redder objects have the same M(1, 1) distribution as the bluer objects. Considering only the small objects ($M(1, 1) \ge 5$ mag), the blue hot CDOs and the blue resonant objects have fainter absolute magnitudes than the redder objects of the same type (see Tab. 11).

3.6.3. Others parameters

Other combinations of the absolute magnitude with other parameters do not show any statistically significant signal.

4. Interpretation scenarios

In the following discussion, we try to elaborate and and understand the results from the statistical analysis described in Section 3 in the framework of two rather different scenarios, i.e. surface colouration by on-going external processes and surface colours as a result of the primordial constitution of the bodies.

4.1. Surface colouration by external processes

Colour diversity of MBOSSes was originally attributed to the action of external processes on their surfaces. These process included resurfacing by impacts, which were expected to produce the neutral-bluish colours of fresh ice excavated and/or spread over the surface, and an irradiation process, producing red colours via radiation aging (eg Luu & Jewitt 1996, for a model). When restricting our MBOSS database samples to objects with $M(1,1) \ge 5$ mag, as done during most of our analysis, a third process, i.e. resurfacing due to atmospheric deposits of material from intrinsic activity, was considered less important for the subsequent discussion, since it should only affect large-sized bodies, the smaller ones losing the material into space.

A critical argument for significant changes in the surface reflectivity due to impacts comes from a lightcurve analysis of MBOSSes: asymmetric or partial impact resurfacing of the body surfaces should cause colour variegations over the rotation of the bodies that are, however, usually not seen (Sekiguchi et al. 2002, S. Sonnett, priv. comm., J. L. Ortiz, priv. comm.). Nevertheless, other authors report colour changes with rotation phase, e.g. for Haumea (Lacerda et al. 2008). Thus, strong variegation is certainly not widespread.

In the colouration scenario based on external processes, DDOs are expected to represent the end state of the radiation reddening of the surface, since one can assume that, for these objects, impact resurfacing does not occurs at a significant level, simply because

the spatial number density of DDOs seems very low and collisions should therefore happen extremely rarely over the lifetime of the solar system. From our statistical analysis, we found that DDOs are not the reddest objects among MBOSSes. Cold CDOs instead show by far the reddest surface colours. To explain the smaller reddening of DDOs in this colouring scenario, one has to assume that cold CDOs do not represent the end state of radiation reddening (despite showing the reddest colours), but high radiation doses, similar to those to which DDOs are exposed, decrease the spectral gradients again. Laboratory experiments (de Bergh et al. 2008) have found materials that display such behaviour, although the end state of these material has rather neutral spectral gradients in the visible wavelength range. However, it remains unclear whether these materials exist in DDOs and, if not, whether the laboratory results apply more generally to more or even all possible surface compounds considered to exist in DDOs. It is also noteworthy to mention the coincidence of the peaks in the spectral gradient distributions of DDOs as well as SDOs, Plutinos, resonant objects, and Centaurs; this may be by chance despite the different importance of collisional resurfacing for these groups.

The colour cascade, with peaks in the colour histograms decreasing in reddening from cold CDOs to hot CDOs, then SDOs, Plutinos, resonant objects, and parts of the Centaurs, must be attributable to the increasing importance of impact resurfacing. However, Centaurs live in a rather empty zone of the planetary system, hence impact resurfacing must contribute less to their evolution. Here, the peak in the spectral gradient diagram could be attributed to resurfacing from intrinsic activity. The very red Centaurs instead represent the inactive objects that are subject to only radiation reddening, though not with extremely high doses (otherwise they could be expected to show similar reddening to DDOs). Among the six known Centaurs showing cometary activity, currently only one has red colours. A larger sample of colours of cometary active Centaurs is therefore needed.

Jupiter Trojans differ in terms of their surface reddening from the MBOSSes. It is remarkable that Jupiter Trojans do not show very 'red' surfaces as objects in all other MBOSS populations do, despite their high exposure to radiation from the Sun (owing to their closer distance by a factor of about 100). Thus, we conclude that this difference may be due to their having different material properties from the other MBOSSes.

An interesting result is seen for the SPCs: the population's peak is well within the range of those of SDOs, Plutinos, moderately red Centaurs, as well as DDOs. Impact resurfacing should be negligible for SPCs and instead resurfacing due to activity should be far more important. On the other hand, owing to the fast, continuous mass-loss on short time-scales (the lifetime for active SPCs is estimated to be $100\,000$ years or less), the surface deposits should represent the intrinsic colour of the cometary material. Radiation aging is unimportant over the lifetime of SPC nuclei in the inner planetary system since cometary activity will result in fast and continuous resurfacing of the body. Hence, SPCs give the impression that their nuclei may consist –on a global scale– of material with rather homogeneous colour properties. As SPCs to a large extent originate from the Kuiper Belt (Gomes et al. 2008; Morbidelli et al. 2008a) and because of their colour similarity to the immediate dynamical relatives therein (SDOs, Plutinos) and the Centaurs from where SPCs are captured, it is attractive to assume that the colours of these relatives should also be globally uniform throughout the bodies. Jewitt (2002a) discusses the missing very red SPCs, suggesting that the red surfaces are rapidly buried as water ice sublimation starts at r < 6 AU.

4.2. Global colouration in the formation era

An alternative interpretation of the global results for the colour and M(1, 1) distribution is that the measured colours actually reflect the global characteristics of the objects, that is throughout the object and not only to its (aged) surface. This hypothesis is supported by the interpretation of SPC colours and the similarities of the peaks in the colour distribution of the MBOSS populations. We note that Brown et al. (2011) introduced a similar hypothesis for the colour diversity among Kuiper Belt objects based on sublimation properties of super-volatile ices. We describe below what can be seen as arguments in support of such a 'primordial' colouration scheme for the MBOSSes, where 'primordial' indicates that the colours of the body materials originate from a time before the body was formed, i.e. the formation period of the planetary system, and have remained largely unaltered ever since.

An additional assumption needed is that the surface materials are not —or at least not by a large amount—affected by external processes such as reddening due to cosmic radiation. The global character of the material colours also implies that impact resurfacing would reproduce the original pre-impact surface colours of the bodies and that large-scale colour diversities on the surface should not occur. Intrinsic activity in MBOSSes will expel gaseous species and solids of which at least the solids may resemble the original colours, while the gases, if not lost to interplanetary space, may condense on the body as surface frost, in which case the surface colour may change. However, our selection criterion to use only objects with M(1,1) > 5 mag for the statistical analysis should significantly reduce the possibility that MBOSSes that may display intrinsic activity 'contaminate' the study results.

In the context of this primordial colouration scenario, the findings from our statistical analysis of the MBOSS colours allow us to draw some interesting conclusions:

- At least four different colour populations may have existed in the protoplanetary disk, i.e. the cold CDOs, the hot CDOs, the SDO/DDO/Plutino populations, and the Jupiter Trojans. The four populations are characterized by different surface colours and spectral gradients of decreasing amplitude in the visible wavelength region, but may have very similar colours in the NIR.
- The colour diversities among MBOSS populations with different surface colours call for formation environments at different and disjunct distances from the Sun. Migration transport may have shifted the original populations to their current locations in the planetary system, that is have placed cold and hot CDOs in the same distance range. This is very much in line with the Nice model —see Section 7.1 of Morbidelli et al. (2008a) for an overall description, or Levison et al. (2008) for more details, which gives an integrated overview of the dynamical history of the outer solar system. In the Nice model, the cold CDOs form a distinct population formed at larger distances from the Sun than the other classes, and in which the hot CDOs are then implanted in the later stages of Neptune's outward migration. The Nice model also specifically accounts for the separate nature of the Jupiter Trojans (Morbidelli et al. 2005) and is able to produce DDOs from a population at closer distances to the Sun than the cold CDOs. The Nice model lists an impressive list of successes in reproducing the dynamical structure of the current solar system;

the observed similarity of the SDO, DDO, and resonant populations could help refining the overall picture painted by the model. However, DDOs and hot CDOs may represent a challenge for migration scenarios, in particular if one wants to include their colour similarities with other MBOSS populations.

- Similar colour properties among MBOSS populations at different locations in the outer planetary system may indicate a common formation region of the bodies. If so, this could apply to both Plutinos and DDOs, objects nowadays found at rather different distances from the Sun. Migration (for the Plutinos) and scattering (for the DDOs) might again have been involved in separating the bodies formed in a common source region.
- The migration and scattering processes have kept a population of the same formation origin together and shifted them to confined destination regions with no obvious signs of intermixing from other sides. This conclusion suggests that the migration and scattering processes in the outer solar system might not have been very violent since they have retained the colour characteristics as population entities.
- The colour distributions for the populations with 'temporary' dynamical object association, i.e. SDOs, Centaurs, and SPCs, should reflect the efficiency of the injection and ejection processes of bodies from the various feeding populations, i.e. Plutinos and both hot and cold CDOs.

There are supportive and also critical arguments from the statistical analysis for the conclusions above. In particular, the similarities and differences in the M(1,1) distributions provide compatible pieces of evidence: the disjunct M(1,1) distributions of the Jupiter Trojans and both the hot and cold CDOs (the latter having been noted in various papers and summarized in Doressoundiram et al. 2008) may call for different formation environments at different solar distances, as also suggested by Morbidelli et al. (2008b). On the other side, the existence of large objects with M(1,1) < 3 mag among Plutinos and DDOs constitutes an independent supportive argument for a formation region that would be common among these populations and distinct from the cold CDOs, as noted by Levison & Stern (2001) and developed by Gomes (2003). Despite the possibly different formation environments of Plutinos and DDOs on the one hand and hot CDOs on the other as suggested by their colour differences, the presence of large objects (M(1,1) < 3 mag) in all three populations suggests that the planetesimal formation and destruction processes were similarly efficient. On the other side, no obvious explanation —except an ad-hoc assumption as an outcome of the formation—can be offered for the anti-correlation between surface colours and excitation parameter as well as the inclinations of the hot CDO population, since the migration process should be 'colour-blind'. While the existence of very red objects among hot CDOs, DDOs, and Plutinos could be attributed to the contamination of cold CDOs, the presence of bodies with blue, neutral surfaces in all populations except for cold CDOs and DDOs may indicate that there are colour contaminants either from the primordial environment or due to dynamical injection during a later phase.

5. A summarizing discussion

The MBOSS database first published in 2003 has since been updated using subsequently published magnitude and colour measurements of TNOs, Centaurs, Jupiter Trojans, and both short- and long-period comets. A qualifying criterion has been applied to the data in the database in order to distinguish noisy measurement sets from more consistent ones. we have performed a new statistical analysis of the qualified colours, spectral gradient, and absolute R magnitudes M(1,1) data in the database with the aim of identifying group properties among the various dynamical MBOSS populations, i.e. for the hot and cold classical disk objects (CDOs), the scattered disk objects (SDOs), the Plutinos, and resonant objects, as well as related objects outside the Kuiper belt such as detached disk objects (DDOs), Centaurs, and SPCs, using a classification based on the SSBN08 system Gladman et al. (2008). Furthermore, Jupiter Trojans are included in the analysis since, from all that is known, these objects are unrelated to the MBOSSes, although represent an abundant population at the border of this region in the solar system. To avoid or at least to reduce the impact on the surface colours due to intrinsic activity, the statistical sample was restricted to objects with normalized (but not yet phase corrected) brightness $M(1,1) \ge 5$ mag.

Our statistical analysis has uncovered indications of indistinguishable and incompatible colour distributions and both the differences and similarities of the M(1, 1) parameters among the MBOSS populations. These results are consistent with a scheme of four different colour (and M(1, 1)) groups in the MBOSS regime, i.e. (1) cold CDOs, (2) hot CDOs, (3) Plutinos, resonant objects, SDOs, DDOs, Centaurs, and SPCs, and (4) Jupiter Trojans. Additional correlations between photometric and dynamical parameters of the various populations were searched for, but only a single well-established one was found in the data of the hot CDOs, for which colour anti-correlates with inclination and the collision energy parameter.

The statistical results have been discussed in the context of two basic interpretation scenarios, i.e. by explaining the surface colours of the MBOSSes as being (1) due to the interplay of resurfacing of impact ejecta (blue colour shift) and radiation reddening (red colour shift) and (2) of primordial nature imposed by the material properties of the formation environment. The two scenarios have the principle capability to explain our findings based on the analysis of the MBOSS database. However, they can also be challenged, each of them in a different way. The resurfacing scenario has to work out whether and under which assumption it is able to reproduce the actually observed colour, spectral gradient, and M(1, 1) distributions of the MBOSSes and in particular that it can explain the relatively moderate reddening of the DDOs in this picture. In this respect, further lab measurements of the radiation reddening of reference materials under high dose levels could play an important role.

The primordial surface-colour scenario provides a good explanation of the apparent uniformity of surface colours among SPCs despite the cometary activity that should be able to excavate material layers of different colours if they exist throughout the nuclei. The primordial surface-colour scenario works with ad hoc assumptions about the colour, spectral gradient, and M(1,1) parameter distributions that require justification. Brown et al. (2011) indicated some first arguments to this respect. However, beyond that, the scenario interpretation relies on the object migration for the transport of the MBOSSes from their formation environment to the various regions where they are found today. Object migration in the early planetary system has already been introduced to

explain the existence of the hot CDOs in the immediate neighborhood of the cold CDOs in the Kuiper Belt. More comprehensive models have been introduced (see Morbidelli et al. (2008b) and Gomes et al. (2008) for reviews). These dynamical models are at the moment—mostly 'colour-blind' and 'size-blind' apart from an a-priori postulation of colour differences for hot and cold CDOs. Our analysis of the MBOSS colours and M(1,1) size parameters would require at least three, possibly four, different source regions in the primordial disk. It is not very plausible that the MBOSSes remained in these original source regions, but it is instead very likely that they were redistributed after formation to the locations in and out of the Kuiper belt where we find them today. A comprehensive model scenario is not yet available that is able to consistently explain the formation environment, the physical properties and the dynamical history of the MBOSS population. Nonetheless, the dynamics, colour, and size distributions together (possibly also albedos) of the measured populations are to be seen as important and constraining benchmarks for these model scenarios —not only qualitatively, but also quantitatively.

In parallel, considerable progress has been achieved in the taxonomic classification of the objects. This process aims to sort the objects among categories defined by similar colour properties (independently of the physico-dynamical classes —see Barucci et al. 2005a), that is an approach orthogonal to the one presented here, which aims to compare the photometric properties of physicodynamical classes. The multi-pronged approaches of dynamical simulations and modelling, taxonomic classification, and photometric analysis is moving towards an integrated understanding of the MBOSS populations.

Acknowledgements. H. Boehnhardt wishes to thank the European Southern Observatory for supporting this work during his research stay in September and November 2011 at ESO Headquarters in Garching/Germany. We are grateful for the very valuable comments and input provided by the anonymous referee. This research has made use of NASA's Astrophysics Data System.

References

```
Barucci, M. A., Belskaya, I. N., Fulchignoni, M., & Birlan, M. 2005a, AJ, 130, 1291
```

Barucci, M. A., Boehnhardt, H., Dotto, E., et al. 2002, Astron. Astrophys., 392, 335

Barucci, M. A., Brown, M. E., Emery, J. P., & Merlin, F. 2008, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P., Morbidelli, A., & Dotson, R., Ed., U. Arizona Press), 143-160

Barucci, M. A., Cruikshank, D. P., Dotto, E., et al. 2005b, A&A, 439, L1

Barucci, M. A., Doressoundiram, A., Tholen, D., Fulchignoni, M., & Lazzarin, M. 1999, Icarus, 142, 476

Barucci, M. A. & Fulchignoni, M. 1990, in Asteroids, Comets III (Uppsala universitet Reprocentralen), 7-10

Barucci, M. A., Merlin, F., Dotto, E., Doressoundiram, A., & de Bergh, C. 2006, A&A, 455, 725

Barucci, M. A., Romon, J., Doressoundiram, A., & Tholen, D. J. 2000, Astron. J., 120, 496

Bauer, J. M., Meech, K. J., Fernández, Y. R., et al. 2003, Icarus, 166, 195 Benecchi, S. D., Noll, K. S., Grundy, W. M., et al. 2009, Icarus, 200, 292

Benecchi, S. D., Noll, K. S., Stephens, D. C., Grundy, W. M., & Rawlins, J. 2011, Icarus, 213, 693

Binzel, R. P. 1992, Icarus, 99, 238

Boehnhardt, H., Barucci, M. A., & Delsanti, A. e. a. 2004, in The First Decadal review of the Edgeworth-Kuiper Belt (Davies, J.K., Barrera, L.J. Ed., Kluwer scientific Publishers), 145

Boehnhardt, H., Delsanti, A., Barucci, A., et al. 2002, A&A, 395, 297

Boehnhardt, H., Rainer, N., Birkle, K., & Schwehm, G. 1999, Astron. Astrophys., 341, 912

Boehnhardt, H., Tozzi, G. P., Birkle, K., et al. 2001, Astron. Astrophys., 378, 653

Brown, M. E. 2008, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P., Morbidelli, A., & Dotson, R., Ed., U. Arizona Press),

Brown, M. E., Schaller, E. L., & Fraser, W. C. 2011, ApJ, 739, L60

Brown, M. E., Trujillo, C. A., & Rabinowitz, D. L. 2005, ApJ, 635, L97

Brown, W. R. & Luu, J. X. 1997, Icarus, 126, 218

Buie, M. W. & Bus, S. J. 1992, Icarus, 100, 288

Buie, M. W., Grundy, W. M., Young, E. F., Young, L. A., & Stern, S. A. 2006, AJ, 132, 290

Buratti, B. J., Hillier, J. K., Heinze, A., et al. 2003, Icarus, 162, 171

Campins, H., A'Hearn, M. F., & McFadden, L.-A. 1987, ApJ, 316, 847

Carraro, G., Maris, M., Bertin, D., & Parisi, M. G. 2006, A&A, 460, L39

Chamberlin, A. B., McFadden, L.-A., Schulz, R., Schleicher, D. G., & Bus, S. J. 1996, Icarus, 119, 173

Consolmagno, G. J., Tegler, S. C., Rettig, T., & Romanishin, W. 2000, Bull. Am. Astron. Soc., 32

Coradini, A. 1977, Computers and Geosciences, 3, 85

Davies, J. K. 2000, in Minor Bodies in the Outer Solar System, Fitzsimmons, Jewitt and West (Ed.) (Springer), 9-24

Davies, J. K., Green, S., McBride, N., et al. 2000, Icarus, 146, 253

Davies, J. K., McBride, N., Ellison, S. L., Green, S. F., & Ballantyne, D. R. 1998, Icarus, 134, 213

Davies, J. K., McBride, N., & Green, S. F. 1997, Icarus, 125, 61

Davies, J. K., Sykes, M. V., & Cruikshank, D. P. 1993, Icarus, 102, 166 Davies, J. K., Tholen, D. J., Whiteley, R. J., et al. 2001, Icarus, 150, 69

de Bergh, C., Delsanti, A., Tozzi, G. P., et al. 2005, A&A, 437, 1115

de Bergh, C., Schmitt, B., Moroz, L. V., Quirico, E., & Cruikshank, D. P. 2008, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P., Morbidelli, A., & Dotson, R., Ed., U. Arizona Press), 483-506

Delahodde, C. E., Meech, K. J., Hainaut, O. R., & Dotto, E. 2001, Astron. Astrophys., 376, 672

Delsanti, A., Hainaut, O., Jourdeuil, E., et al. 2004, Astron. Astrophys., 417, 1145

Delsanti, A., Peixinho, N., Boehnhardt, H., et al. 2006, AJ, 131, 1851

Delsanti, A. C., Boehnhardt, H., Barrera, L., et al. 2001, Astron. Astrophys., 380, 347

Delsemme, A. H. 1982, in IAU Colloq. 61: Comet Discoveries, Statistics, and Observational Selection (L. L. Wilkening, Ed.), 85-130

DeMeo, F. E., Fornasier, S., Barucci, M. A., et al. 2009, A&A, 493, 283

Dones, L., Weissman, P. R., Levison, H. F., & Duncan, M. J. 2004, in Comets II (Festou, M. C., Keller, H. U., & Weaver, H. A., Ed.), 153-174

Doressoundiram, A., Barucci, M. A., Romon, J., & Veillet, C. 2001, Icarus, 154, 277

Doressoundiram, A., Barucci, M. A., Tozzi, G. P., et al. 2005a, Planet. Space Sci., 53, 1501

Doressoundiram, A., Boehnhardt, H., Tegler, S. C., & Trujillo, C. 2008, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P., Morbidelli, A., & Dotson, R., Ed., U. Arizona Press), 91-104

Doressoundiram, A., Peixinho, N., de Bergh, C., et al. 2002, Astron. J., 124, 2279

```
Doressoundiram, A., Peixinho, N., Doucet, C., et al. 2005b, Icarus, 174, 90
Doressoundiram, A., Peixinho, N., Moullet, A., et al. 2007, AJ, 134, 2186
Doressoundiram, A., Tozzi, G. P., Barucci, M. A., et al. 2003, Astron. J., 125, 2721
Dotto, E., Barucci, M. A., Boehnhardt, H., et al. 2003, Icarus, 162, 408
Dotto, E., Fornasier, S., Barucci, M. A., et al. 2006, Icarus, 183, 420
Dumas, C., Merlin, F., Barucci, M. A., et al. 2007, A&A, 471, 331
Duncan, M. J. & Levison, H. F. 1997, Science, 276, 1670
Farnham, T. L. & Davies, J. K. 2003, Icarus, 164, 418
Ferrin, I., Rabinowitz, D., Schaefer, B., et al. 2001, Astrophys. J., 548, L243
Fink, U., Hoffmann, M., Grundy, W., Hicks, M., & Sears, W. 1992, Icarus, 97, 145
Fornasier, S., Doressoundiram, A., Tozzi, G. P., et al. 2004, aa, 421, 353
Fornasier, S., Dotto, E., Hainaut, O., et al. 2007, Icarus, 190, 622
Fulchignoni, M., Belskaya, I., Barucci, M. A., de Sanctis, M. C., & Doressoundiram, A. 2008, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt,
   H., Cruikshank, D. P., Morbidelli, A., & Dotson, R., Ed., U. Arizona Press), 181-192
Fulchignoni, M., Birlan, M., & Barucci, M. A. 2000, Icarus, 146, 204
Gil-Hutton, R. & Licandro, J. 2001, Icarus, 152, 246
Gladman, B., Kavelaars, J. J., Nicholson, P. D., Loredo, T. J., & Burns, J. A. 1998, Astron. J., 116, 2042
Gladman, B., Marsden, B. G., & Vanlaerhoven, C. 2008, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P., Morbidelli, A.,
   & Dotson, R., Ed., U. Arizona Press), 43-57
Gomes, R. S. 2003, Icarus, 161, 404
Gomes, R. S., Fern Ndez, J. A., Gallardo, T., & Brunini, A. 2008, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P.,
   Morbidelli, A., & Dotson, R., Ed., U. Arizona Press), 259-273
Green, S. F., McBride, N., O Ceallaigh, D. P., et al. 1997, Mon. Not. R. Astron. Soc., 290, 186
Gutiérrez, P. J., Ortiz, J. L., Alexandrino, E., Roos-Serote, M., & Doressoundiram, A. 2001, A&A, 371, L1
Hainaut, O. R. 2001, priv. Comm.
Hainaut, O. R., Delahodde, C. E., Boehnhardt, H., et al. 2000, Astron. Astrophys., 356, 1076
Hainaut, O. R. & Delsanti, A. C. 2002, A&A, 389, 641, paper I
Hartmann, W. K., Cruikshank, D. P., Degewij, J., & Capps, R. W. 1981, Icarus, 47, 333
Jewitt, D. 2009a, AJ, 137, 4296
Jewitt, D. 2009b, AJ, 137, 4296
Jewitt, D. & Kalas, P. 1998, Astrophys. J., 499, L103
Jewitt, D. & Luu, J. 1989, AJ, 97, 1766
Jewitt, D. & Luu, J. 1998, Astron. J., 115, 1667
Jewitt, D. & Meech, K. J. 1986, Astrophys. J., 310, 937
Jewitt, D., Peixinho, N., & Hsieh, H. H. 2007, AJ, 134, 2046
Jewitt, D., Sheppard, S., & Fernández, Y. 2003, AJ, 125, 3366
Jewitt, D. C. 2002a, AJ, 123, 1039
Jewitt, D. C. 2002b, Astron. J., 123, 1039
Jewitt, D. C. & Luu, J. X. 2001, Astron. J., 122, 2099
Jewitt, D. C. & Meech, K. J. 1988, Astrophys. J., 328, 974
Jewitt, D. C. & Sheppard, S. S. 2002, Astron. J., 123, 2110
Kavelaars, J., Jones, L., Gladman, B., Parker, J. W., & Petit, J.-M. 2008, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P.,
   Morbidelli, A., & Dotson, R., Ed., U. Arizona Press), 59-69
Keller, H. U. & Thomas, N. 1989, A&A, 226, L9
Kenyon, S. J. & Bromley, B. C. 2004, AJ, 128, 1916
Lacerda, P., Jewitt, D., & Peixinho, N. 2008, AJ, 135, 1749
Lamy, P. & Toth, I. 2009, Icarus, 201, 674
Lamy, P. L., Toth, I., Fernandez, Y. R., & Weaver, H. A. 2004, The sizes, shapes, albedos, and colors of cometary nuclei (Festou, M. C., Keller, H. U., & Weaver,
   H. A.), 223-264
Lamy, P. L., Toth, I., Jorda, L., et al. 2002, Icarus, 156, 442
Landolt, A. 1992, Astrophys. J., 104, 340
Levison, H. F., Duncan, M. J., Brasser, R., & Kaufmann, D. E. 2010, Science, 329, 187
Levison, H. F., Morbidelli, A., Vanlaerhoven, C., Gomes, R., & Tsiganis, K. 2008, Icarus, 196, 258
Levison, H. F. & Stern, S. A. 2001, AJ, 121, 1730
Lowry, S. C., Fitzsimmons, A., & Collander-Brown, S. 2003, A&A, 397, 329
Lowry, S. C. & Weissman, P. R. 2003, Icarus, 164, 492
Luu, J. & Jewitt, D. 1990, Icarus, 86, 69
Luu, J. X. 1993, Icarus, 104, 138
Luu, J. X. & Jewitt, D. C. 1996, Astron. J., 111, 499
McBride, N., Green, S. F., Davies, J. K., et al. 2003, Icarus, 161, 501
McKinnon, W. B., Prialnik, D., Stern, S. A., & Coradini, A. 2008, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P.,
   Morbidelli, A., & Dotson, R., Ed., U. Arizona Press), 213-241
Meech, K. J., Hainaut, O. R., & Marsden, B. G. 2004, Icarus, 170, 463
Meech, K. J. & Svoren, J. 2004, in Comets II (Festou, M. C., Keller, H. U., & Weaver, H. A., Ed.), 317-335
Millis, R. L., A'Hearn, M. F., & Campins, H. 1988, Astrophys. J., 324, 1194
Morbidelli, A., Levison, H. F., & Gomes, R. 2008a, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P., Morbidelli, A., &
   Dotson, R., Ed., U. Arizona Press), 275-292
Morbidelli, A., Levison, H. F., & Gomes, R. 2008b, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P., Morbidelli, A., &
   Dotson, R., Ed., U. Arizona Press), 275-292
Morbidelli, A., Levison, H. F., Tsiganis, K., & Gomes, R. 2005, Nature, 435, 462
Mueller, B. E. A., Tholen, D. J., Hartmann, W. K., & Cruikshank, D. P. 1992, Icarus, 97
```

Ortiz, J. L., Sota, A., Moreno, R., et al. 2004, A&A, 420, 383

Parker, J. W., Stern, S. A., Festou, M. C., A'Hearn, M. F., & Weintraub, D. 1997, Astron. J., 113, 1899

Peixinho, N., Boehnhardt, H., Belskaya, I., et al. 2004, Icarus, 170, 153

Peixinho, N., Doressoundiram, A., Delsanti, A., et al. 2003, A&A, 410, L29

Neese, C. 2011, NASA Planetary Data System, 150

Müller, T. G., Lellouch, E., Böhnhardt, H., et al. 2009, Earth Moon and Planets, 105, 209 N. McBride, J. K. Davies, S. F. G. & Foster, M. J. 1999, Mon. Not. R. Astron. Soc., 306, 799

```
Peixinho, N., Lacerda, P., & Jewitt, D. 2008, AJ, 136, 1837
```

Peixinho, N., Lacerda, P., Ortiz, J. L., et al. 2001, Astron. Astrophys., 371, 753

Perna, D., Barucci, M. A., Fornasier, S., et al. 2010, A&A, 510, A53

Protopapa, S., Alvarez-Candal, A., Barucci, M. A., et al. 2009, A&A, 501, 375

Protopapa, S., Boehnhardt, H., Herbst, T. M., et al. 2008, A&A, 490, 365

Rabinowitz, D., Tourtellotte, S., Brown, M., & Trujillo, C. 2004, IAU Circ., 8295, 2

Rabinowitz, D. L., Schaefer, B. E., Schaefer, M., & Tourtellotte, S. W. 2008, AJ, 136, 1502

Rabinowitz, D. L., Schaefer, B. E., & Tourtellotte, S. W. 2007, AJ, 133, 26

Reyment, R. & Joreskog, K. 1993 (Cambridge University Press)

Romanishin, W. & Tegler, S. C. 1999, Nature, 398, 129

Romanishin, W., Tegler, S. C., & Consolmagno, G. J. 2010, AJ, 140, 29 Romanishin, W., Tegler, S. C., Levine, J., & Butler, N. 1997, Astron. J., 113, 1893

Romon-Martin, J., Barucci, M. A., de Bergh, C., et al. 2002, Icarus, 160, 59

Schaefer, B. E. & Rabinowitz, D. L. 2002, Icarus, 160, 52

Schaller, E. L. 2010, in European Physical Journal Web of Conferences, Vol. 9, European Physical Journal Web of Conferences, 267

Schaller, E. L. & Brown, M. E. 2007, ApJ, 659, L61

Sekiguchi, T., Boehnhardt, H., Hainaut, O. R., & Delahodde, C. E. 2002, A&A, 385, 281

Sheppard, S. S. 2010, AJ, 139, 1394

Sicardy, B., Ortiz, J. L., Assafin, M., et al. 2011, Nature, 478, 493

Snodgrass, C., Carry, B., Dumas, C., & Hainaut, O. 2010, A&A, 511, A72+

Snodgrass, C., Fitzsimmons, A., & Lowry, S. C. 2005, A&A, 444, 287

Stansberry, J., Grundy, W., Brown, M., et al. 2008, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P., Morbidelli, A., & Dotson, R., Ed., U. Arizona Press), 161-179

Stern, S. A. & Tholen, D. J. 1998, Pluto and Charon (Stern, S. A. & Tholen, D. J., Ed.)

Tegler, S. C., Bauer, J. M., Romanishin, W., & Peixinho, N. 2008, in The Solar System Beyond Neptune (Barucci, M. A., Boehnhardt, H., Cruikshank, D. P., Morbidelli, A., & Dotson, R., Ed., U. Arizona Press), 161-179

Tegler, S. C. & Romanishin, W. 1998, Nature, 392, 49

Tegler, S. C. & Romanishin, W. 2000, Nature, 407, 979

Tegler, S. C. & Romanishin, W. 2003, Icarus, 161, 181

Tegler, S. C., Romanishin, W., & Consolmagno, S. J. 2003, Astrophys. J., 599, L49

Tholen, D. J. 1984, PhD thesis, University of Arizona

Tholen, D. J. & Barucci, M. A. 1989, in Asteroids II (The University of Arizona Press), 298-315

Trujillo, C. A. & Brown, M. E. 2002, ApJ, 566, L125

Trujillo, C. A., Brown, M. E., Barkume, K. M., Schaller, E. L., & Rabinowitz, D. L. 2007, ApJ, 655, 1172

Trujillo, C. A., Jewitt, D. C., & Luu, J. X. 2001, AJ, 122, 457

Trujillo, C. A., Sheppard, S. S., & Schaller, E. L. 2011, ApJ, 730, 105

Weintraub, D. A., Tegler, S. C., & Romanishin, W. 1997, Icarus, 128, 456

Appendix A: Online material

The following tables list the colours of the objects in the database at the time this analysis was performed (Table A.1), as well as the corresponding references (Table A.2). These tables do not appear in the journal; are available electronically from the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/. The up-to-date version of the database is available from http://www.eso.org/~ohainaut/MBOSS.

Table A.1. Object class and colour data

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	$Grt^{(3)} \pm \sigma$	B-V ±σ	V-R ±σ	R-I ±σ	V-J ±σ	J-H ±σ	H-K $\pm\sigma$
1P/Halley	SP Comet	2	13.558 ± 0.596	4.266 ± 2.015	0.720 ± 0.040	0.410 ± 0.030	0.390 ± 0.060	_	_	_
2P/Encke	SP Comet	6	15.207 ± 0.596	5.117 ± 2.574	0.780 ± 0.020	0.424 ± 0.053	0.408 ± 0.060	_	_	_
4P/Faye	SP Comet	2	16.020 ± 0.009	7.951 ± 0.000	_	0.450 ± 0.040	_	_	_	_
6P/dArrest	SP Comet	3	16.239 ± 0.009	13.264 ± 2.147	0.770 ± 0.040	0.563 ± 0.067	0.450 ± 0.040	_		_
7P/PW	SP Comet	2	15.185 ± 0.009	4.264 ± 2.620	_	0.400 ± 0.050	0.410 ± 0.060	_	_	_
8P/Tuttle	SP Comet	2	12.799 ± 0.009	15.869 ± 2.326	0.890 ± 0.040	0.530 ± 0.040	0.530 ± 0.060	_	_	_
9P/Tempel 1	SP Comet	14	14.560 ± 0.460	9.959 ± 0.826		0.468 ± 0.010	0.469 ± 0.013		_	_
10P/Tempel 2	SP Comet	10	14.287 ± 0.304	13.740 ± 1.578	0.800 ± 0.020	0.521 ± 0.037	0.520 ± 0.030		_	_
14P/Wolf	SP Comet	2	15.423 ± 0.009	17.666 ± 3.210	_	0.570 ± 0.070	0.510 ± 0.060	_	_	_
15P/Finlay	SP Comet	1	17.488 ± 0.009	_	_	_	_	_	_	_
16P/Brooks2	SP Comet	1	16.107 ± 0.009			_	_		_	
17P/Holmes	SP Comet	2	16.094 ± 0.009	18.608 ± 0.000		0.560 ± 0.020	_		_	
19P/Borrelly	SP Comet	1	15.860 ± 0.009	_	_	_	_	_	_	_
21P/GZ	SP Comet	2	17.259 ± 0.009	12.661 ± 2.057	0.800 ± 0.030	0.500 ± 0.020	_	_	_	_
22P/Kopff	SP Comet	4	16.093 ± 0.156	12.577 ± 2.730	0.795 ± 0.056	0.519 ± 0.046	0.450 ± 0.055	_	_	_
4P/Schaumasse	SP Comet	i	17.744 ± 0.009		_	_	_	_	_	_
26P/GS	SP Comet	2	16.642 ± 0.136	5.227 ± 0.000	_	0.420 ± 0.139	_		_	
28P/Neujmin1	SP Comet	11	12.580 ± 0.009	9.967 ± 4.007	_	0.494 ± 0.072	0.440 ± 0.079		_	
29P/SW1	Centaurs	3	11.322 ± 0.009	14.948 ± 1.099	0.790 ± 0.030	0.536 ± 0.030	0.497 ± 0.025	_		
30P/Reinmuth1	SP Comet	1	14.602 ± 0.009	——————————————————————————————————————	0.770 ± 0.030	0.550 ± 0.050	0.477 ± 0.023		_	
31P/SW2	SP Comet	1	14.802 ± 0.009 14.803 ± 0.009	_	_	_	_	_	_	_
2P/ComasSola	SP Comet	1	14.478 ± 0.009	_	_	_	_		_	
3P/Daniel	SP Comet	1		_	_	_	_		_	_
	SP Comet	1	17.053 ± 0.009	_	_	_	_	_		_
6P/Whipple			15.432 ± 0.009	2 101 - 2 015	_	0.200 + 0.020	0.660 + 0.060	_	_	_
7P/Forbes	SP Comet	2	17.348 ± 0.009	3.181 ± 2.015		0.290 ± 0.030	0.660 ± 0.060		_	_
9P/Oterma	Centaurs	3	12.464 ± 0.009	3.094 ± 3.964	0.890 ± 0.070	0.386 ± 0.063	0.412 ± 0.114		_	_
OP/Vaisala1	SP Comet	1	15.983 ± 0.009	_		_	_		_	_
1P/TG	SP Comet	1	18.034 ± 0.009	_	_	_	_	_	_	_
2P/Neujmin3	SP Comet	1	17.259 ± 0.009	_	_	_	_	_	_	_
3P/WH	SP Comet	1	15.983 ± 0.009			_	_		_	_
4P/Reinmuth2	SP Comet	2	16.225 ± 0.009	24.892 ± 0.000	_	0.620 ± 0.080	_	_	_	_
5P/HMP	SP Comet	2	17.744 ± 0.009	0.768 ± 2.444	1.080 ± 0.050	0.440 ± 0.050	0.210 ± 0.050	_	_	_
6P/Wirtanen	SP Comet	3	18.369 ± 0.009	4.079 ± 0.000	_	0.407 ± 0.081	_	_	_	_
P/AJ	SP Comet	2	15.024 ± 0.009	3.454 ± 7.417	0.780 ± 0.080	0.400 ± 0.080	_		_	_
18P/Johnson	SP Comet	1	14.970 ± 0.009	_	_	_	_	_	_	_
19P/AR	SP Comet	3	14.120 ± 0.277	9.452 ± 2.141	0.770 ± 0.030	0.471 ± 0.032	0.444 ± 0.064	1.630 ± 0.200	0.360 ± 0.141	0.140 ± 0.141
50P/Arend	SP Comet	2	17.371 ± 0.009	24.132 ± 4.627		0.810 ± 0.100	0.260 ± 0.090		_	
51P/Harrington	SP Comet	1	15.648 ± 0.009			_	_		_	
52P/HA	SP Comet	1	16.690 ± 0.009	_	_	_	_	_	_	_
3P/VanBiesbroec	SP Comet	2	14.647 ± 0.009	-2.672 ± 0.000	_	0.328 ± 0.081	_	_	_	_
55P/TT	SP Comet	2	15.543 ± 0.482	10.765 ± 2.444	0.750 ± 0.050	0.510 ± 0.050	0.420 ± 0.050	_	_	_
66P/SB	SP Comet	1	16.294 ± 0.009	10.705 ± 2.444 —					_	
7P/dTND	SP Comet	1	16.239 ± 0.009 16.239 ± 0.009	_	_	_	_		_	
8P/JN	SP Comet	1	18.369 ± 0.009	_	_		_	<u> </u>	_	
9P/KK	SP Comet	2	17.771 ± 0.009	$-$ 12.937 \pm 3.558	_	0.620 ± 0.070	0.270 ± 0.080	_	_	_
50P/Tsuchinshan2	SP Comet	1	17.771 ± 0.009 17.744 ± 0.009	12.937 ± 3.336	_	0.020 ± 0.070	0.270 ± 0.000	_	_	_
51P/SS	SP Comet	1	17.744 ± 0.009 18.229 ± 0.009	_	_	_	_	_	_	_
					_	_	_	_	_	_
52P/Tsuchinshan1	SP Comet	1	17.744 ± 0.009	10.054 + 2.272		0.500 + 0.050	0.420 + 0.040		_	
53P/Wild1	SP Comet	2	16.453 ± 0.009	10.054 ± 2.273	_	0.500 ± 0.050	0.420 ± 0.040		_	

Table A.1. continued.

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	Grt $\pm \sigma$	B-V ±σ	V-R ±σ	R-I ±σ	V-J ±σ	J-H ±σ	H-K $\pm\sigma$
64P/SG	SP Comet	1	16.239 ± 0.009	_	_	_		_	_	
55P/Gunn	SP Comet	1	13.853 ± 0.009	_	_	_	_	_	_	_
57P/CG	SP Comet	2	15.754 ± 0.009	13.630 ± 0.000	_	0.510 ± 0.060	_	_	_	_
68P/Klemola	SP Comet	1	15.547 ± 0.009	_	_	_	_	_	_	_
69P/Taylor	SP Comet	1	14.947 ± 0.009	_	_	_	_	_	_	_
70P/Kojima	SP Comet	2	15.912 ± 0.009	22.758 ± 0.000	_	0.600 ± 0.090	_	_	_	_
71P/Clark	SP Comet	2	18.097 ± 0.009	27.065 ± 0.000	_	0.640 ± 0.070	_	_	_	_
72P/DF	SP Comet	1	17.744 ± 0.009	_	_	_	_	_	_	_
73P/SW3	SP Comet	1	16.850 ± 0.050	10.751 ± 0.000	_	0.480 ± 0.235	_	_	_	
73P/SW3C	SP Comet	1	18.097 ± 0.009	_	_	_	_	_	_	_
74P/SC	SP Comet	1	15.518 ± 0.009	_	_	_	_	_	_	
75P/Kohoutek	SP Comet	1	15.754 ± 0.009	_	_	_	_	_	_	_
76P/WKI	SP Comet	1	19.667 ± 0.009						_	
77P/Longmore	SP Comet	1	15.358 ± 0.009	_		_	_	_		
78P/Gehrels2	SP Comet	1	16.498 ± 0.009	_		_				_
79P/dTH	SP Comet	1	16.529 ± 0.009	_						_
81P/Wild2	SP Comet	1	15.754 ± 0.009	_	_	_	_	_	_	
31P/Wild2 32P/Gehrels3	SP Comet	1		_	_	_	_	_	_	_
,			17.943 ± 0.009		_		_	_	_	_
84P/Giclas	SP Comet	2	17.488 ± 0.009	-3.328 ± 0.000	1.500 . 0.000	0.320 ± 0.030	_	_	_	_
36P/Wild3	SP Comet	3	19.092 ± 0.009	18.211 ±35.399	1.580 ± 0.080	0.555 ± 0.385	_	_	_	
87P/Bus	SP Comet	2	20.024 ± 0.009	16.889 ± 0.000	_	0.543 ± 0.020	_	_	_	_
88P/Howell	SP Comet	1	15.866 ± 0.009	_	_	_	_	_	_	_
39P/Russell2	SP Comet	1	16.864 ± 0.009	_	_	_		_	_	_
90P/Gehrels1	SP Comet	1	14.602 ± 0.009	_	_	_		_	_	_
91P/Russell3	SP Comet	1	16.690 ± 0.009	_	_	_	_	_	_	_
92P/Sanguin	SP Comet	2	16.882 ± 0.009	16.937 ± 1.982	_	0.540 ± 0.040	0.540 ± 0.040	_	_	_
93P/HL	SP Comet	1	_	-7.552 ± 0.000		0.267 ± 0.075			_	_
94P/Russell4	SP Comet	1	15.866 ± 0.009	_	_	_	_	_	_	
96P/Machholz1	SP Comet	2	14.734 ± 0.009	6.036 ± 0.000	_	0.429 ± 0.027	_	_	_	
97P/MB	SP Comet	1	16.107 ± 0.009	_		_			_	
98P/Takamizawa	SP Comet	1	14.418 ± 0.009	_	_	_	_	_	_	
99P/Kowal1	SP Comet	1	13.853 ± 0.009	_	_	_	_	_	_	
100P/Hartley1	SP Comet	1	16.690 ± 0.009	_					_	
101P/Chernykh	SP Comet	1	15.358 ± 0.009						_	
103P/Hartley2	SP Comet	1	17.744 ± 0.009	_		_				
104P/Kowal2	SP Comet	1	17.744 ± 0.009 17.259 ± 0.009	_			_		_	
105P/SB	SP Comet	1	17.259 ± 0.009 17.259 ± 0.009							
106P/Schuster	SP Comet	2	17.239 ± 0.009 17.394 ± 0.009	$\frac{-}{12.396 \pm 2.913}$	1.010 ± 0.060	0.520 ± 0.060	0.450 ± 0.060	_		
						0.320 ± 0.000	0.430 ± 0.000	_	_	
107P/WH	SP Comet	6	15.865 ± 0.220	0.119 ± 4.538	0.674 ± 0.089	0.361 ± 0.048		_	_	_
109P/ST	SP Comet	2	12.002 ± 0.376	18.608 ± 0.000	_	0.560 ± 0.050	_	_	_	
10P/Hartley3	SP Comet	2	15.597 ± 0.009	30.401 ± 0.000	_	0.670 ± 0.090	_	_	_	_
11P/HRC	SP Comet	1	18.369 ± 0.009		_			_	_	_
12P/UN	SP Comet	2	17.488 ± 0.009	15.593 ± 0.000	_	0.530 ± 0.040		_	_	_
13P/Spitaler	SP Comet	1	17.053 ± 0.009	—	<u> </u>	<u> </u>	-	_	_	_
14P/WS	SP Comet	2	17.799 ± 0.009	12.158 ± 1.087	0.850 ± 0.030	0.460 ± 0.020	0.540 ± 0.020	_	_	_
115P/Maury	SP Comet	1	17.033 ± 0.009	_	_	_	_	_	_	_
116P/Wild4	SP Comet	1	14.539 ± 0.009	_	_	_	_	_	_	_
117P/HRA	SP Comet	1	14.304 ± 0.009	_	_	_	_	_	_	
18P/SL	SP Comet	1	15.358 ± 0.009	_	_	_	_	_	_	_

Table A.1. continued.

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	Grt $\pm \sigma$	B-V ±σ	V-R ±σ	R-I ±σ	V-J ±σ	J-H $\pm\sigma$	H-K $\pm\sigma$
119P/PH	SP Comet	1	15.270 ± 0.009	_	_	_	_	_	_	_
120P/Mueller1	SP Comet	1	16.379 ± 0.009	_	_	_	_		_	_
121P/SH	SP Comet	1	16.212 ± 0.009	_	_	_	_	_	_	_
123P/WH	SP Comet	1	15.547 ± 0.009	_		_	_			
24P/Mrkos	SP Comet	1	16.239 ± 0.009	_		_	_			
25P/Spacewatch	SP Comet	1	17.744 ± 0.009	_	_	_	_	_	_	_
26P/IRAS	SP Comet	1	16.280 ± 0.009	_	_	_	_	_	_	_
28P/SH	SP Comet	1	15.451 ± 0.009	_	_	_	_	_	_	_
29P/SL	SP Comet	1	15.358 ± 0.009	_	_	_	_	_	_	_
30P/McNH	SP Comet	1	15.983 ± 0.009	_	_	_	_	_	_	_
31P/Mueller2	SP Comet	1	17.744 ± 0.009	_	_	_	_		_	
32P/HRA	SP Comet	1	17.488 ± 0.009	_	_	_	_		_	
34P/KV	SP Comet	1	16.529 ± 0.009	_	_	_	_	_	_	_
35P/SL	SP Comet	1	16.239 ± 0.009	_	_	_	_	_	_	_
36P/Mueller3	SP Comet	1	15.866 ± 0.009	_	_			_	_	_
37P/SL	SP Comet	1	14.947 ± 0.009	_	_	_	_	_	_	_
38P/SL	SP Comet	1	17.744 ± 0.009	_			_		_	
39P/VO	SP Comet	1	15.185 ± 0.009	_		_	_		_	
40P/BS	SP Comet	1	15.451 ± 0.009	_		_	_		_	_
41P/Machholz2	SP Comet	1	17.259 ± 0.009	_	_	<u> </u>	_	<u> </u>	<u> </u>	_
43P/KM	SP Comet	5	17.239 ± 0.009 13.598 ± 0.009	20.354 ± 1.102	0.820 ± 0.023	0.580 ± 0.020	0.560 ± 0.021			_
44P/Kushida	SP Comet	1	16.864 ± 0.009	20.554 ± 1.102 —	0.020 ± 0.023	0.380 ± 0.020	0.300 ± 0.021	_	_	_
47P/KM	SP Comet	1	20.648 ± 0.009	_	_	_	_	_	_	_
48P/AL	SP Comet	1	15.648 ± 0.009	_	_	_	_	_	_	
52P/HL	SP Comet	1	13.946 ± 0.009 13.946 ± 0.009	_	_	_	_	_	_	_
54P/Brewington	SP Comet	1	16.379 ± 0.009	_	_	_	_	_	_	
		2		26 202 + 1 076	0.870 + 0.040	0.605 + 0.020	0.735 ± 0.060	_	_	
66P/2001T4	Centaurs		_	36.293 ± 1.976	0.870 ± 0.040	0.695 ± 0.030		_	_	_
04P/Wild2	SP Comet	1 2	8.729 ± 0.042	9.261 ± 0.863	0.707 . 0.020	0.510 . 0.022	0.436 ± 0.093	1 577 : 0 026	0.420 + 0.042	0.125 + 0.02
172-Aneas	J.Trojan				0.727 ± 0.030	0.510 ± 0.022	0.400 ± 0.030	1.577 ± 0.036	0.430 ± 0.042	0.135 ± 0.036
173-Anchises	J.Trojan	1	8.793 ± 0.059	4.286 ± 1.443	0.811 ± 0.034	0.402 ± 0.035	0.403 ± 0.052			
647-Menelaus	J.Trojan	1		6.803 ± 1.630		0.428 ± 0.043	0.438 ± 0.061			
871-Astyanax	J.Trojan	3	11.149 ± 0.093	7.777 ± 2.485	0.716 ± 0.071	0.456 ± 0.053	0.424 ± 0.089	1.422 ± 0.103	0.373 ± 0.053	0.123 ± 0.04
993 RO	Res. 3:2	2	8.612 ± 0.134	24.485 ± 4.231	0.850 ± 0.070	0.590 ± 0.137	0.480 ± 0.113	_	_	_
994 ES ₂	Classic Cold	1	7.509 ± 0.130					_	_	_
994 EV ₃	Classic Cold	4	7.048 ± 0.178	30.400 ± 7.250	1.065 ± 0.110	0.588 ± 0.156	0.800 ± 0.126	_	_	_
994 TA	Centaurs	2	11.421 ± 0.126	31.545 ± 6.243	1.261 ± 0.139	0.672 ± 0.080	0.740 ± 0.210	_	_	_
995 DB ₂	Classic Cold	2	8.058 ± 0.120	54.335 ± 0.000	_	_	_	_	_	_
995 DC ₂	Classic Cold	5	6.809 ± 0.204	35.475 ± 7.654	_	0.770 ± 0.160	0.580 ± 0.160	_	_	_
995 FB ₂₁	Classic Cold	4	7.017 ± 0.099	_	_	_	_	_	_	_
995 HM ₅	Res. 3:2	8	7.849 ± 0.109	9.432 ± 3.661	0.649 ± 0.102	0.460 ± 0.097	0.428 ± 0.126	_	1.180 ± 0.470	_
995 WY ₂	Classic Cold	2	6.935 ± 0.149	19.923 ± 11.862	1.030 ± 0.280	0.600 ± 0.230	0.510 ± 0.280	_	_	_
996 KV ₁	Classic Hot	1	_	_	_	_	_		0.430 ± 0.256	
996 RQ ₂₀	Classic Hot	7	6.872 ± 0.100	22.364 ± 4.307	0.935 ± 0.141	0.558 ± 0.096	0.591 ± 0.109		0.380 ± 0.226	
996 RR ₂₀	Res. 3:2	4	6.811 ± 0.252	36.280 ± 5.193	1.143 ± 0.106	0.730 ± 0.095	0.628 ± 0.125	_	_	_
996 TC ₆₈	Classic Hot	1	6.734 ± 0.073	22.758 ± 0.000	_	0.600 ± 0.078	_	_	_	_
996 TK ₆₆	Classic Cold	4	6.190 ± 0.116	29.906 ± 4.054	0.993 ± 0.062	0.680 ± 0.067	0.551 ± 0.097	_	_	_
996 TS ₆₆	Classic Hot	7	5.947 ± 0.130	26.561 ± 4.871	1.028 ± 0.079	0.672 ± 0.106	0.637 ± 0.088	1.824 ± 0.178	0.650 ± 0.071	
997 CT ₂₉	Classic Cold	4	6.452 ± 0.173	32.698 ± 4.508	<u> </u>	0.762 ± 0.105	0.490 ± 0.120	<u> </u>	0.410 ± 0.238	
997 CV ₂₉	Classic Hot	2	7.154 ± 0.030	28.171 ± 2.056	1.210 ± 0.010	0.650 ± 0.020	_	_	0.360 ± 0.243	_

Table A.1. continued.

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	Grt ±σ	B-V ±σ	V-R ±σ	R-I ±σ	V-J ±σ	J-H ±σ	H-K ±σ
1997 GA ₄₅	Classic Hot	1	7.744 ± 0.500	_	_	_	_	_	_	_
1997 QH ₄	Classic Hot	4	7.128 ± 0.175	27.144 ± 4.569	1.088 ± 0.143	0.641 ± 0.092	0.631 ± 0.107		_	_
1997 RL ₁₃	Classic Hot	1	9.361 ± 0.300	_	_	_	_		_	_
1997 RT ₅	Classic Hot	4	6.951 ± 0.379	15.015 ± 4.506	1.075 ± 0.121	0.474 ± 0.107	0.539 ± 0.091	_	0.380 ± 0.231	_
1997 RX ₉	Classic Hot	1	7.802 ± 0.100		_	_	_	_	_	_
1997 SZ ₁₀	Res.(other)	1	8.148 ± 0.060	28.167 ± 2.909	1.140 ± 0.080	0.650 ± 0.030	_	_	_	_
1998 FS ₁₄₄	Classic Hot	4	6.717 ± 0.105	23.851 ± 3.089	0.950 ± 0.044	0.588 ± 0.060	0.510 ± 0.085		0.390 ± 0.228	
1998 KG ₆₂	Classic Cold	5	6.367 ± 0.283	24.857 ± 3.762	1.039 ± 0.105	0.609 ± 0.095	0.610 ± 0.078		0.410 ± 0.231	
1998 KS ₆₅	Classic Cold	2	7.166 ± 0.040	26.149 ± 1.858	1.090 ± 0.040	0.640 ± 0.020	_		_	
1998 KY ₆₁	Classic Cold	1	_	_	_	_	_	_	0.350 ± 0.278	_
1998 UR ₄₃	Res. 3:2	4	8.124 ± 0.134	14.454 ± 7.356	0.784 ± 0.101	0.583 ± 0.120	0.354 ± 0.175	_	_	_
1998 UU ₄₃	Res.(other)	2	6.925 ± 0.110	14.799 ± 3.950	_	0.610 ± 0.120	0.400 ± 0.114	_	0.490 ± 0.273	_
1998 WS ₃₁	Res. 3:2	2	7.931 ± 0.203	18.738 ± 6.008	0.726 ± 0.080	0.606 ± 0.102	0.439 ± 0.172	_	_	_
1998 WU ₂₄	Scattered	1	— — — — — — — — — — — — — — — — — — —	12.934 ± 1.361	0.780 ± 0.034	0.530 ± 0.037	0.460 ± 0.051	1.670 ± 0.043	0.430 ± 0.125	0.240 ± 0.189
1998 WV ₂₄	Classic Cold	2	7.126 ± 0.067	12.734 ± 1.301 11.777 ± 3.357	0.770 ± 0.034 0.770 ± 0.010	0.502 ± 0.058	0.450 ± 0.031 0.450 ± 0.106		— U.123	5.2-10 ± 0.107 —
1998 W V ₂₄ 1998 WV ₃₁	Res. 3:2	2	7.639 ± 0.007	13.835 ± 4.507	0.770 ± 0.010 0.790 ± 0.099	0.502 ± 0.038 0.521 ± 0.075	0.430 ± 0.100 0.481 ± 0.144			
1998 WW ₃₁	Classic Hot	1	7.039 ± 0.072 —	8.864 ± 0.000	0.790 ± 0.099 —	0.321 ± 0.073	0.461 ± 0.144 —	_	_	
1998 WW ₃₁ 1998 WW _{31+B}	Classic Hot	1	_	2.628 ± 0.000	_	_	_	_	_	
1998 W W 31+B 1998 W X ₂₄	Classic Hot Classic Cold	2	6.241 ± 0.099	2.628 ± 0.000 30.875 ± 3.793	1.090 ± 0.050	0.727 ± 0.086	0.500 ± 0.092	_	_	_
					1.090 ± 0.030				0.400 ± 0.235	_
1998 WX ₃₁	Classic Cold	3	6.159 ± 0.095	29.252 ± 3.312	_	0.638 ± 0.083	0.653 ± 0.081	_		_
1998 WY ₂₄	Classic Cold	2	6.441 ± 0.086	29.095 ± 3.185		0.620 ± 0.092	0.680 ± 0.078	_	0.400 ± 0.240	
1998 WZ ₃₁	Res. 3:2	3	8.045 ± 0.115	6.687 ± 4.513	0.727 ± 0.093	0.489 ± 0.114	0.339 ± 0.114	_	0.400 ± 0.234	
1998 XY ₉₅	Detached	2	6.438 ± 0.143	41.429 ± 5.409	0.930 ± 0.233	0.720 ± 0.145	0.752 ± 0.126		_	_
1999 CB ₁₁₉	Classic Hot	1	6.737 ± 0.076	33.869 ± 2.678	1.212 ± 0.102	0.714 ± 0.076	0.645 ± 0.062			
1999 CD ₁₅₈	Res.(other)	5	4.822 ± 0.099	16.860 ± 2.607	0.864 ± 0.056	0.520 ± 0.053	0.575 ± 0.064	1.862 ± 0.076	0.423 ± 0.135	0.048 ± 0.116
1999 CF ₁₁₉	Detached	3	6.933 ± 0.133	14.704 ± 4.163		0.622 ± 0.115	0.365 ± 0.106		0.380 ± 0.226	
1999 CH ₁₁₉	Classic Hot	1	_		_	_	_	_	0.320 ± 0.266	_
1999 CJ ₁₁₉	Classic Cold	1	_	_	_	_	_	_	0.450 ± 0.265	_
1999 CL ₁₁₉	Classic Hot	2	5.699 ± 0.072	19.530 ± 3.359	_	0.497 ± 0.077	0.665 ± 0.094	_	0.450 ± 0.282	_
1999 CQ ₁₃₃	Classic Hot	1	_				_		0.460 ± 0.282	
1999 CX ₁₃₁	Res.(other)	3	6.910 ± 0.107	19.521 ± 4.293	0.918 ± 0.124	0.664 ± 0.128	0.434 ± 0.105		0.370 ± 0.238	_
1999 HJ ₁₂	Classic Cold	2	_	30.391 ± 0.000	_	_	_	_	0.480 ± 0.598	_
1999 HS ₁₁	Classic Cold	3	6.331 ± 0.079	31.102 ± 3.766	1.121 ± 0.082	0.698 ± 0.056	0.600 ± 0.090		_	
1999 HV ₁₁	Classic Cold	1	7.003 ± 0.050	21.706 ± 2.057	1.110 ± 0.060	0.590 ± 0.020	_		_	
1999 OD ₄	Classic Hot	1	_	_	_	_	_	_	0.460 ± 0.282	_
1999 OE ₄	Classic Cold	4	6.831 ± 0.177	13.989 ± 7.765	1.103 ± 0.236	0.567 ± 0.159	0.414 ± 0.180	_	0.360 ± 0.216	_
1999 OH₄	Classic Hot	2	6.321 ± 0.123	-7.656 ± 9.026	_	0.200 ± 0.134	0.430 ± 0.335	_	0.330 ± 0.328	_
1999 OJ ₄	Classic Cold	5	6.912 ± 0.062	26.192 ± 3.113	1.098 ± 0.157	0.668 ± 0.072	0.549 ± 0.073	_	0.230 ± 0.197	_
1999 OJ _{4+B}	Classic Cold	1	—	34.360 ± 0.000					—	_
1999 OM ₄	Classic Cold	1	7.502 ± 0.100	18.976 ± 3.909	1.137 ± 0.135	0.602 ± 0.104	0.499 ± 0.088			
1999 RC ₂₁₅	Classic Cold	2	6.571 ± 0.126	33.893 ± 4.703	— — — — — — — — — — — — — — — — — — —	0.810 ± 0.134	0.520 ± 0.120	_	0.320 ± 0.225	
1999 RC ₂₁₅ 1999 RT ₂₁₄	Classic Cold	1	0.571 ± 0.120 —	35.732 ± 0.000		J.010 ± 0.134	0.520 ± 0.120		0.320 ± 0.223	
1999 RT ₂₁₄ 1999 RX ₂₁₄	Classic Cold	3	6.413 ± 0.052	20.207 ± 2.817	1.054 ± 0.072	0.593 ± 0.065	0.530 ± 0.066	_	_	
	Classic Hot	2	7.010 ± 0.065	18.063 ± 3.131	0.693 ± 0.072	0.565 ± 0.003 0.565 ± 0.080	0.530 ± 0.000 0.530 ± 0.084	_	_	
1999 RY ₂₁₄								_	_	_
1999 TR ₁₁	Res. 3:2	2	8.144 ± 0.152	37.094 ± 4.048	1.020 ± 0.080	0.750 ± 0.097	0.650 ± 0.122	_	_	
1999 XY ₁₄₃	Classic Hot	1	5.924 ± 0.065	26.014 ± 2.399		0.590 ± 0.071	0.660 ± 0.064	_	- 0.460 - 0.265	
2000 AF ₂₅₅	Detached	1	_				_	_	0.460 ± 0.265	
2000 CE ₁₀₅	Classic Cold	2	_	14.512 ± 0.000	_	_	_	_	0.160 ± 0.184	_
2000 CF ₁₀₅	Classic Cold	6		24.245 ± 0.000	_	_	_	_	0.400 ± 0.266	_

Table A.1. continued.

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	Grt ±σ	B-V ±σ	V-R ±σ	R-I ±σ	V-J ±σ	J-H ±σ	H-K ±σ
2000 CF _{105+B}	Classic Cold	1	_	21.377 ± 0.000	_	_	_	_	_	
2000 CG ₁₀₅	Classic Hot	3	6.124 ± 0.311	_	_	0.107 ± 0.246	0.293 ± 0.265	_	0.510 ± 0.298	_
000 CK ₁₀₅	Res. 3:2	2	6.370 ± 0.118	25.479 ± 4.225	_	0.590 ± 0.127	0.650 ± 0.114	_	0.390 ± 0.256	_
000 CL_{104}	Classic Cold	4	6.642 ± 0.229	23.634 ± 4.673	1.223 ± 0.167	0.600 ± 0.119	0.612 ± 0.128	_	0.450 ± 0.285	_
000 CN ₁₀₅	Classic Cold	4	5.261 ± 0.072	29.677 ± 1.174	1.100 ± 0.020	0.636 ± 0.072	0.640 ± 0.020	_	_	_
000 CO ₁₀₅	Classic Hot	2	5.594 ± 0.078	22.327 ± 2.904	_	0.620 ± 0.085	0.550 ± 0.078	_	0.380 ± 0.228	
000 CP ₁₀₄	Classic Hot	2	_	18.106 ± 0.000	_	_	_		0.440 ± 0.289	
000 CQ ₁₀₅	Detached	5	5.965 ± 0.049	7.864 ± 2.902	0.671 ± 0.023	0.449 ± 0.038	0.346 ± 0.101	_	0.420 ± 0.262	
000 CQ ₁₁₄	Classic Cold	3	6.660 ± 0.147	34.780 ± 4.289	_	0.690 ± 0.156	0.650 ± 0.134	_	0.420 ± 0.403	_
000 CQ_{114+B}	Classic Cold	1	_	33.683 ± 0.000		_	_		_	_
000 FS ₅₃	Classic Cold	2	7.300 ± 0.080	34.603 ± 1.976	1.060 ± 0.040	0.710 ± 0.020	_	_	_	_
000 FV ₅₃	Res. 3:2	1	7.550 ± 0.087	17.018 ± 4.070	_	0.448 ± 0.094	0.679 ± 0.116	_	_	
000 FZ ₅₃	Centaurs	2	10.990 ± 0.085	14.413 ± 3.939	_	0.686 ± 0.090	0.238 ± 0.103			
000 GV ₁₄₆	Classic Cold	1	— — — — — — — — — — — — — — — — — — —	35.732 ± 0.000	_	—			_	
000 KK ₄	Classic Hot	3	5.982 ± 0.103	17.292 ± 2.477	0.910 ± 0.040	0.580 ± 0.102	0.640 ± 0.064	1.790 ± 0.106	_	
000 KK ₄	Classic Hot	1	J.702 ± 0.103	27.864 ± 0.000	0.710 ± 0.040	0.300 ± 0.102	- -			
000 KL ₄ 000 OU ₆₉	Classic Cold	3	6.839 ± 0.120	10.128 ± 0.000	_	0.473 ± 0.130	_	_	0.400 ± 0.235	
		2	0.039 ± 0.120		_	0.473 ± 0.130	_	_	0.400 ± 0.233 0.560 ± 0.336	
000 PD ₃₀	Classic Cold	7	 5 706 + 0 001	19.719 ± 0.000	0.752 + 0.060	0.272 + 0.077	0.410 + 0.040	1 (4(+ 0.000		0.260 + 0.07
000 PE ₃₀	Detached	1	5.796 ± 0.081	2.353 ± 3.158	0.752 ± 0.069	0.373 ± 0.077	0.410 ± 0.049	1.646 ± 0.080	0.452 ± 0.114	0.260 ± 0.07
000 PH ₃₀	Detached	1	_		_	_	_	_	0.550 ± 0.361	
000 QL ₂₅₁	Classic Cold	1	_	6.266 ± 0.000	_	_	_	_	_	_
000 QL_{251+B}	Classic Cold	1	_	12.082 ± 0.000	_	_	_	_	_	_
001 FL ₁₈₅	Classic Cold	1	<u> </u>	23.082 ± 0.000	— — — — — — — — — — — — — — — — — — —	<u> </u>	_	_	_	_
001 FM ₁₉₄	Detached	1	7.582 ± 0.040	7.035 ± 2.909	0.760 ± 0.030	0.440 ± 0.030	_	_	_	_
001 FU ₁₇₂	Res. 3:2	1	8.165 ± 0.181	22.552 ± 8.420	_	0.670 ± 0.192	0.470 ± 0.225	_	_	_
001 KA ₇₇	Classic Hot	4	5.050 ± 0.094	37.759 ± 3.264	1.104 ± 0.107	0.704 ± 0.064	0.716 ± 0.120	_	_	_
001 KB ₇₇	Res. 3:2	6	7.373 ± 0.063	11.050 ± 3.211	_	0.434 ± 0.083	0.548 ± 0.089		_	
001 KD ₇₇	Res. 3:2	9	5.926 ± 0.104	24.005 ± 2.575	1.123 ± 0.054	0.624 ± 0.070	0.565 ± 0.069	2.294 ± 0.092	0.366 ± 0.141	_
001 KG ₇₇	Scattered	1	8.340 ± 0.120	7.034 ± 5.595	0.810 ± 0.040	0.440 ± 0.060	_	_	_	_
001 KY ₇₆	Res. 3:2	1	_	_		_	_		0.390 ± 0.228	
001 OG ₁₀₉	Classic Cold	1	_	_	_	_	_		0.350 ± 0.300	_
001 OK ₁₀₈	Classic Cold	1	_	_				_	0.430 ± 0.275	_
001 QC ₂₉₈	Classic Hot	3	6.421 ± 0.030	10.333 ± 2.434	0.750 ± 0.080	0.490 ± 0.030	0.480 ± 0.040	_	0.430 ± 0.323	_
$OO1 QC_{298+B}$	Classic Hot	1		6.266 ± 0.000	_	_	_		— U.430 ± 0.323	_
001 QC_{298+B} 001 QD_{298}	Classic Hot	1	6.185 ± 0.170	30.399 ± 8.330	0.970 ± 0.130	0.670 ± 0.090	_	_	_	_
001 QD ₂₉₈ 001 QF ₂₉₈	Res. 3:2	8	5.128 ± 0.126	2.287 ± 2.687	0.570 ± 0.130 0.644 ± 0.070	0.381 ± 0.062	0.360 ± 0.085	1.126 ± 0.124	0.317 ± 0.172	0.165 ± 0.11
001 QX ₃₂₂	Detached	3	6.388 ± 0.080	18.862 ± 5.260	0.914 ± 0.058	0.561 ± 0.002 0.563 ± 0.056	- 0.000	1.120 ± 0.124	0.317 ± 0.172 0.410 ± 0.247	0.105 ± 0.11
001 QX ₃₂₂ 001 XR ₂₅₄	Classic Cold	1	0.366 ± 0.060	15.002 ± 3.200 15.011 ± 0.000	0.914 ± 0.036	0.303 ± 0.030	_	_	0.410 ± 0.247 —	_
		1	_		_	_	_	_	_	_
001 XR _{254+B}	Classic Cold	•	_	16.024 ± 0.000	_	_	_	_	0.270 + 0.226	_
001 XU ₂₅₄	Classic Hot	1	10.000 : 0.000	20.752 + 6.501	1 170 : 0 020	0.750 : 0.070	_		0.370 ± 0.226	
001 XZ ₂₅₅	Centaurs	1	10.800 ± 0.080	39.753 ± 6.501	1.170 ± 0.020	0.750 ± 0.070			_	
002 CB ₂₄₉	Centaurs	1	10.587 ± 0.109	5.278 ± 4.487		0.492 ± 0.116	0.300 ± 0.114	_	_	_
002 DH ₅	Centaurs	1	10.190 ± 0.061	4.151 ± 2.833	0.663 ± 0.072	0.391 ± 0.069	0.416 ± 0.081	_	_	_
002 GB_{32}	Scattered	1	7.637 ± 0.010	18.001 ± 1.087	0.880 ± 0.028	0.510 ± 0.020	0.610 ± 0.020	_	_	_
002 PP ₁₄₉	Classic Hot	1	7.256 ± 0.030	-6.534 ± 3.793	0.850 ± 0.100	0.280 ± 0.040	_	_	_	_
003 FZ ₁₂₉	Detached	1	6.982 ± 0.020	10.329 ± 1.812	0.840 ± 0.057	0.480 ± 0.040	0.460 ± 0.030	_	_	_
003 GH ₅₅	Classic Cold	1	5.949 ± 0.050	25.973 ± 5.595	1.120 ± 0.050	0.630 ± 0.060	_	_	_	_
003 HB ₅₇	Scattered	1	7.389 ± 0.020	13.225 ± 1.525	0.830 ± 0.042	0.480 ± 0.030	0.540 ± 0.030	_	_	_
003 HX ₅₆	Classic Hot	1	7.030 ± 0.209	-1.963 ±13.676		0.350 ± 0.226	0.260 ± 0.459		_	_

Table A.1. continued.

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	Grt $\pm \sigma$	B-V ±σ	V-R ±σ	R-I $\pm \sigma$	V-J ±σ	J-H $\pm\sigma$	H-K $\pm\sigma$
2003 QA ₉₂	Classic Cold	1	_	25.973 ± 3.793	1.040 ± 0.030	0.630 ± 0.040	_	_	_	_
003 QD_{112}	Centaurs	1	11.641 ± 0.030	_	_	_	_	_	_	_
2003 QK ₉₁	Detached	1	6.967 ± 0.030	11.829 ± 1.812	0.870 ± 0.057	0.500 ± 0.040	0.470 ± 0.030			
2003 QQ ₉₁	Classic Hot	1	_	13.624 ± 7.416	0.670 ± 0.060	0.510 ± 0.080	_			_
$2003 \ QW_{111}$	Res.(other)	1	_	27.864 ± 0.000	_	_	_			_
2003 QW_{111+B}	Res.(other)	1	_	30.391 ± 0.000	_	_	_	_	_	
2003 QW ₉₀	Res. 3:2	2	_	15.349 ± 0.000		_		1.930 ± 0.162		
2003 QY ₉₀	Classic Cold	1	_	32.348 ± 0.000		_				
2003 QY_{90+B}	Classic Cold	1	_	32.348 ± 0.000	_	_	_	_	_	_
2003 SQ ₃₁₇	Classic Hot	1	6.161 ± 0.020	_	_	_	_	_	-0.450 ± 0.197	_
2003 TH ₅₈	Res. 3:2	1	7.007 ± 0.045	5.615 ± 2.028	0.610 ± 0.064	0.380 ± 0.045	0.480 ± 0.045	1.160 ± 0.098	1.280 ± 0.201	
2003 TJ ₅₈	Classic Cold	1	—	19.719 ± 0.000	_	_	_	_	_	_
2003 TJ _{58+B}	Classic Cold	1	_	18.639 ± 0.000	_	_	_	_		
2003 UY ₂₉₁	Classic Cold	1		20.163 ± 3.383	0.880 ± 0.099	0.510 ± 0.070	0.670 ± 0.070			
2003 UZ ₁₁₇	Classic Cold	4	4.961 ± 0.020	-0.537 ± 1.824	0.676 ± 0.031	0.310 ± 0.070 0.350 ± 0.028	0.334 ± 0.039	_	_	_
2003 VL ₁₇₉	Classic Cold	1	7.701 E 0.020	7.953 ± 9.243	0.810 ± 0.031 0.810 ± 0.040	0.350 ± 0.028 0.450 ± 0.100	0.334 ± 0.039	_	_	
2003 1L ₁₇₉ 2004 OJ ₁₄	Detached	1	6.991 ± 0.020	1.933 ± 9.243 15.681 ± 1.525	0.810 ± 0.040 0.900 ± 0.042	0.430 ± 0.100 0.520 ± 0.030	0.540 ± 0.030	_	_	_
			0.991 ± 0.020			0.320 ± 0.030	0.340 ± 0.030	_	_	_
2004 PB ₁₀₈	Classic Hot	1	_	19.177 ± 0.000	_	_	_	_	_	_
2004 PB _{108+B}	Classic Hot	1		21.377 ± 0.000	_	_	_			_
2004 PT ₁₀₇	Classic Hot	1	5.847 ± 0.010					_	0.540 ± 0.228	_
2004 PY ₄₂	1	_	17.581 ± 1.360	0.770 ± 0.030	0.550 ± 0.030	0.540 ± 0.020		_	_	
2004 VN ₁₁₂	Detached	1	-	12.396 ± 2.913	0.900 ± 0.085	0.520 ± 0.060	0.450 ± 0.060			_
2004 XR ₁₉₀	Detached	1	3.937 ± 0.030	10.679 ± 1.982	0.790 ± 0.057	0.450 ± 0.040	0.520 ± 0.040	-	-	
2005 CB ₇₉	Classic Hot	1	4.375 ± 0.028	1.054 ± 1.574	0.740 ± 0.036	0.350 ± 0.036	0.380 ± 0.036	1.040 ± 0.076	-0.510 ± 0.175	_
2005 EO ₂₉₇	Detached	1	7.220 ± 0.030	13.673 ± 2.955	0.840 ± 0.071	0.480 ± 0.050	0.570 ± 0.080	_	_	_
2005 EO ₃₀₄	Classic Cold	1	_	38.554 ± 0.000	_	_	_	_	_	_
2005 EO _{304+B}	Classic Cold	1	_	33.683 ± 0.000	_	_	_			
2005 GE ₁₈₇	Res. 3:2	1	7.214 ± 0.030	_	_	_	_	_	0.660 ± 0.144	_
2005 PU ₂₁	Scattered	1	6.091 ± 0.020	30.943 ± 1.087	1.140 ± 0.028	0.650 ± 0.020	0.680 ± 0.020	_	_	_
2005 SD ₂₇₈	Detached	1	5.916 ± 0.020	17.797 ± 1.087	0.970 ± 0.028	0.560 ± 0.020	0.530 ± 0.020	_	_	_
2006 SQ ₃₇₂	Scattered	1	7.667 ± 0.020	24.677 ± 1.702	1.030 ± 0.042	0.590 ± 0.030	0.650 ± 0.040			
2007 TG ₄₂₂	Scattered	1	6.186 ± 0.010	13.918 ± 1.982	0.880 ± 0.057	0.510 ± 0.040	0.510 ± 0.040	_	_	_
2007 VJ ₃₀₅	Detached	1	6.713 ± 0.020	14.913 ± 1.525	0.920 ± 0.042	0.520 ± 0.030	0.520 ± 0.030	_	_	_
2008 KV ₄₂	Scattered	1	8.565 ± 0.040	8.461 ± 2.913	0.820 ± 0.085	0.470 ± 0.060	0.420 ± 0.060			
2008 OG ₁₉	Detached	1	4.611 ± 0.010	18.921 ± 0.677	0.940 ± 0.014	0.530 ± 0.010	0.590 ± 0.010	_		
2008 YB ₃	SP Comet	1	9.715 ± 0.010	10.521 ± 0.677 10.549 ± 0.677	0.800 ± 0.014	0.460 ± 0.010	0.490 ± 0.010			_
2060-Chiron	Centaurs	34	6.092 ± 0.069	0.114 ± 0.996	0.700 ± 0.020	0.361 ± 0.017	0.325 ± 0.023	1.199 ± 0.110	0.294 ± 0.079	0.065 ± 0.09
2223-Sarpedon	J.Trojan	1	9.265 ± 0.046	8.785 ± 1.244	0.753 ± 0.032	0.465 ± 0.025	0.323 ± 0.023 0.440 ± 0.041		0.274 ± 0.077	0.005 ± 0.07
2357-Phereclos	J.Trojan	1	8.817 ± 0.078	7.601 ± 1.876	0.733 ± 0.032 0.718 ± 0.059	0.403 ± 0.023 0.427 ± 0.045	0.463 ± 0.068	_	_	
		1		-0.087 ± 1.847	0.718 ± 0.039 0.677 ± 0.052	0.427 ± 0.045 0.352 ± 0.045	0.403 ± 0.008 0.339 ± 0.067	_	_	_
3548-Eurybates	J.Trojan	-	0.490 + 0.090					_	_	_
1035-1986 WD	J.Trojan	2	9.489 ± 0.080	10.259 ± 1.995	0.752 ± 0.040	0.484 ± 0.043	0.451 ± 0.069			_
1829-Sergestus	J.Trojan	1	11.122 ± 0.068	4.196 ± 1.856	0.851 ± 0.050	0.420 ± 0.039	0.372 ± 0.065		_	_
5130-Ilioneus	J.Trojan	1	9.656 ± 0.048	9.188 ± 1.378	0.763 ± 0.034	0.481 ± 0.026	0.424 ± 0.046			
5145-Pholus	Centaurs	41	7.165 ± 0.076	48.354 ± 1.930	1.261 ± 0.121	0.788 ± 0.036	0.822 ± 0.054	2.612 ± 0.048	0.391 ± 0.047	-0.037 ± 0.0
5244-Amphilochos	J.Trojan	1	_	3.187 ± 1.754	_	0.407 ± 0.044	0.363 ± 0.064	_	_	_
5258-1989 AU ₁	J.Trojan	1	_	8.233 ± 1.590	_	0.466 ± 0.041	0.425 ± 0.059	_	_	_
5511-Cloanthus	J.Trojan	1	10.287 ± 0.047	10.774 ± 1.229	0.906 ± 0.027	0.442 ± 0.027	0.526 ± 0.042	_	_	_
6545-1986 TR ₆	J.Trojan	1	10.219 ± 0.073	10.572 ± 1.962	0.734 ± 0.041	0.499 ± 0.042	0.436 ± 0.069	_	_	_
5998-Tithonus	J.Trojan	1	11.284 ± 0.059	8.155 ± 1.282	0.787 ± 0.040	0.455 ± 0.032	0.438 ± 0.046	_	_	_

Table A.1. continued.

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	Grt $\pm \sigma$	B-V ±σ	V-R ±σ	R-I $\pm \sigma$	V-J ±σ	J-H $\pm\sigma$	H-K $\pm\sigma$
7066-Nessus	Centaurs	17	9.054 ± 0.075	39.777 ± 3.448	1.090 ± 0.040	0.763 ± 0.059	0.689 ± 0.087	2.290 ± 0.040	0.309 ± 0.268	-0.089 ± 0.385
7352-1994 CO	J.Trojan	1	9.873 ± 0.056	4.815 ± 1.629	0.713 ± 0.036	0.417 ± 0.034	0.397 ± 0.056	_		_
3405-Asbolus	Centaurs	43	8.999 ± 0.059	12.765 ± 1.370	0.738 ± 0.038	0.508 ± 0.053	0.505 ± 0.038	1.654 ± 0.048	0.386 ± 0.103	0.170 ± 0.171
0030-1989 UX ₅	J.Trojan	1	11.121 ± 0.047	11.783 ± 1.115	0.887 ± 0.024	0.493 ± 0.027	0.480 ± 0.039		_	_
9430-Erichthonio	J.Trojan	1	11.306 ± 0.106	10.723 ± 3.060	0.742 ± 0.079	0.488 ± 0.060	0.456 ± 0.109		_	_
9818-Eurymachos	J.Trojan	1	_	-0.162 ± 1.824	0.673 ± 0.052	0.339 ± 0.046	0.355 ± 0.067	_	_	_
10199-Chariklo	Centaurs	64	6.508 ± 0.016	13.595 ± 0.883	0.802 ± 0.049	0.491 ± 0.016	0.520 ± 0.014	1.685 ± 0.110	0.414 ± 0.046	0.097 ± 0.048
0370-Hylonome	Centaurs	25	9.257 ± 0.159	9.668 ± 6.478	0.690 ± 0.110	0.469 ± 0.154	0.496 ± 0.189	1.310 ± 0.100	0.180 ± 0.085	0.270 ± 0.092
1089-1994 CS ₈	J.Trojan	1	10.681 ± 0.098	4.732 ± 2.619	0.689 ± 0.073	0.423 ± 0.056	0.384 ± 0.094			_
1351-1997 TS ₂₅	J.Trojan	1	10.793 ± 0.076	9.325 ± 2.033	0.739 ± 0.044	0.498 ± 0.044	0.402 ± 0.072			_
1488-1988 RM ₁₁	J.Trojan	2	11.524 ± 0.194	6.507 ± 2.832	0.777 ± 0.110	0.436 ± 0.062	0.420 ± 0.103	_	_	_
1663-1997 GO ₂₄	J.Trojan	1	10.838 ± 0.052	6.580 ± 1.438	0.837 ± 0.030	0.409 ± 0.030	0.463 ± 0.049	_	_	
2917-1998 TG ₁₆	J.Trojan	2	11.388 ± 0.073	11.839 ± 1.962	0.724 ± 0.042	0.537 ± 0.042	0.410 ± 0.069	1.707 ± 0.066	0.475 ± 0.086	0.205 ± 0.081
2921-1998 WZ ₅	J.Trojan	2	10.948 ± 0.073	3.492 ± 1.876	0.673 ± 0.040	0.403 ± 0.047	0.380 ± 0.068	_	_	— — — — — — — — — — — — — — — — — — —
3463-Antiphos	J.Trojan	2	11.083 ± 0.066	7.093 ± 1.913	0.692 ± 0.045	0.449 ± 0.034	0.412 ± 0.066	1.417 ± 0.048	0.343 ± 0.055	0.103 ± 0.052
3862-1999 XT ₁₆₀	J.Trojan J.Trojan	1	11.003 ± 0.000 —	2.203 ± 1.754		0.381 ± 0.034	0.370 ± 0.064			
4707-2000 CC ₂₀	J.Trojan	2	11.410 ± 0.112	4.232 ± 1.902	0.752 ± 0.041	0.412 ± 0.035	0.370 ± 0.004 0.385 ± 0.067	_	_	_
5094-1999 WB ₂	J.Trojan	3	11.582 ± 0.112	5.186 ± 2.495	0.652 ± 0.065	0.472 ± 0.065 0.477 ± 0.065	0.322 ± 0.007	1.401 ± 0.066	0.342 ± 0.062	0.095 ± 0.058
5502-1999 NV ₂₇	J.Trojan J.Trojan	3	9.943 ± 0.075	7.368 ± 1.724	0.766 ± 0.047	0.477 ± 0.003 0.445 ± 0.036	0.322 ± 0.094 0.430 ± 0.060	1.565 ± 0.046	0.342 ± 0.002 0.492 ± 0.041	0.053 ± 0.036 0.152 ± 0.034
5535-2000 AT ₁₇₇	J.Trojan J.Trojan	4	10.555 ± 0.075	9.836 ± 2.098	0.739 ± 0.047	0.470 ± 0.030 0.470 ± 0.049	0.450 ± 0.000 0.461 ± 0.082	1.681 ± 0.045	0.500 ± 0.041	0.132 ± 0.034 0.123 ± 0.047
	Classic Cold	5	6.979 ± 0.094			0.470 ± 0.049 0.707 ± 0.093	0.401 ± 0.082 0.651 ± 0.166		0.300 ± 0.046	0.123 ± 0.047
5760-1992 QB ₁	Res. 3:2	5 6		27.538 ± 4.806 12.407 ± 4.044	0.869 ± 0.143 0.802 ± 0.071			_	0.450 ± 0.253	_
5788-1993 SB			8.026 ± 0.143			0.475 ± 0.077	0.514 ± 0.114			0.040 + 0.10
5789-1993 SC	Res. 3:2	17	6.740 ± 0.077	34.698 ± 4.012	1.045 ± 0.117	0.688 ± 0.074	0.697 ± 0.093	2.236 ± 0.200	0.412 ± 0.240	-0.040 ± 0.19
5807-1994 GV ₉	Classic Cold	1	6.813 ± 0.091	38.552 ± 0.000	_	0.740 ± 0.099		_	0.460 - 0.705	_
5809-1994 JS	Res.(other)	2	7.479 ± 0.160	28.194 ± 6.000		0.760 ± 0.170	0.480 ± 0.149		0.460 ± 0.705	_
5810-1994 JR ₁	Res. 3:2	9	6.873 ± 0.079	15.313 ± 4.751	1.010 ± 0.180	0.614 ± 0.114	0.538 ± 0.098		0.430 ± 0.242	
5820-1994 TB	Res. 3:2	15	7.499 ± 0.116	37.669 ± 3.728	1.107 ± 0.126	0.697 ± 0.075	0.739 ± 0.099	2.368 ± 0.162	0.428 ± 0.119	0.120 ± 0.120
5836-1995 DA ₂	Res.(other)	6	7.809 ± 0.091	16.356 ± 5.630		0.550 ± 0.110	0.500 ± 0.160			
5874-1996 TL ₆₆	Scattered	16	5.208 ± 0.131	1.451 ± 2.858	0.687 ± 0.072	0.369 ± 0.052	0.370 ± 0.077	1.452 ± 0.114	0.267 ± 0.144	-0.018 ± 0.13
5875-1996 TP ₆₆	Res. 3:2	7	6.958 ± 0.076	30.636 ± 3.598	1.031 ± 0.112	0.655 ± 0.076	0.673 ± 0.086	2.309 ± 0.060	0.170 ± 0.078	0.020 ± 0.092
5883-1997 CR ₂₉	Classic Hot	3	7.076 ± 0.135	20.484 ± 7.048	0.750 ± 0.152	0.538 ± 0.157	0.620 ± 0.182		0.490 ± 0.320	
5977-1998 MA ₁₁	J.Trojan	1	10.610 ± 0.046	8.770 ± 1.049	0.748 ± 0.033	0.465 ± 0.025	0.441 ± 0.036	_	_	_
6684-1994 JQ ₁	Classic Cold	8	6.646 ± 0.113	31.346 ± 4.713	1.134 ± 0.068	0.736 ± 0.101	0.650 ± 0.123	_	_	_
7416-1988 RR ₁₀	J.Trojan	1	12.739 ± 0.097	12.134 ± 2.732	0.742 ± 0.071	0.488 ± 0.055	0.498 ± 0.097	_	_	_
8060-1999 XJ ₁₅₆	J.Trojan	1	_	3.456 ± 1.719	0.758 ± 0.056	0.412 ± 0.043	0.364 ± 0.063	_	_	_
.8137-2000 OU ₃₀	J.Trojan	1	11.131 ± 0.057	9.115 ± 1.218	0.733 ± 0.039	0.496 ± 0.031	0.409 ± 0.044		_	_
8268-Dardanos	J.Trojan	1	12.080 ± 0.070	12.538 ± 1.495	0.795 ± 0.042	0.529 ± 0.040	0.451 ± 0.056	_		_
8493-Demoleon	J.Trojan	2	10.813 ± 0.083	3.154 ± 2.460	0.703 ± 0.073	0.395 ± 0.047	0.380 ± 0.087	1.383 ± 0.045	0.384 ± 0.046	0.091 ± 0.040
8940-2000 QV ₄₉	J.Trojan	1	11.615 ± 0.097	8.449 ± 2.608	0.709 ± 0.072	0.465 ± 0.055	0.429 ± 0.093		_	_
9255-1994 VK ₈	Classic Cold	4	7.016 ± 0.163	24.540 ± 5.861	1.010 ± 0.060	0.659 ± 0.095	0.490 ± 0.255	_	0.520 ± 0.297	_
9299-1996 SZ ₄	Res. 3:2	5	8.330 ± 0.166	13.725 ± 4.694	0.754 ± 0.100	0.522 ± 0.061	0.446 ± 0.154	_	_	_
9308-1996 TO ₆₆	Classic Hot	16	4.520 ± 0.042	2.077 ± 2.168	0.671 ± 0.057	0.389 ± 0.043	0.356 ± 0.053	0.997 ± 0.101	_	_
9521-Chaos	Classic Hot	13	4.503 ± 0.090	23.292 ± 2.396	0.932 ± 0.054	0.608 ± 0.041	0.571 ± 0.087	1.835 ± 0.087	0.400 ± 0.095	0.033 ± 0.094
0000-Varuna	Classic Hot	8	3.455 ± 0.090	26.843 ± 1.917	0.906 ± 0.052	0.637 ± 0.040	0.628 ± 0.040	2.010 ± 0.050	0.564 ± 0.070	-0.038 ± 0.10
0108-1995 QZ ₉	Res. 3:2	3	7.889 ± 0.399	14.511 ± 3.460	0.880 ± 0.040	0.515 ± 0.050		_	_	-
0161-1996 TR ₆₆	Res.(other)	1	_	_	_	_	_	_	0.470 ± 0.296	_
0738-1999 XG ₁₉₁	J.Trojan	2	11.536 ± 0.073	10.132 ± 1.962	0.776 ± 0.041	0.472 ± 0.042	0.467 ± 0.069	1.593 ± 0.062	0.520 ± 0.084	0.113 ± 0.08
3549-Epicles	J.Trojan	1	11.812 ± 0.073 11.812 ± 0.118	7.916 ± 2.720	0.770 ± 0.041 0.800 ± 0.071	0.472 ± 0.042 0.485 ± 0.068	0.387 ± 0.009	1.393 ± 0.002 —		
3694-1997 KZ ₃	J.Trojan J.Trojan	1	11.340 ± 0.053	8.337 ± 1.151	0.723 ± 0.035	0.483 ± 0.008 0.474 ± 0.029	0.387 ± 0.101 0.418 ± 0.041	1.657 ± 0.033	0.493 ± 0.033	0.211 ± 0.028

Table A.1. continued.

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	Grt $\pm \sigma$	B-V ±σ	V-R ±σ	R-I ±σ	V-J ±σ	J-H $\pm\sigma$	H-K $\pm\sigma$
24233-1999 XD ₉₄	J.Trojan	1	11.433 ± 0.071	9.103 ± 1.978	0.704 ± 0.051	0.481 ± 0.037	0.418 ± 0.069	_	_	_
4341-2000 AJ ₈₇	J.Trojan	1	12.013 ± 0.067	2.026 ± 1.928	0.713 ± 0.043	0.369 ± 0.035	0.390 ± 0.067	_	_	_
24380-2000 AA ₁₆₀	J.Trojan	1	_	1.556 ± 1.804	0.734 ± 0.055	0.391 ± 0.048	0.336 ± 0.068	_	_	_
24390-2000 AD ₁₇₇	J.Trojan	2	11.595 ± 0.066	12.512 ± 1.913	0.700 ± 0.042	0.513 ± 0.034	0.462 ± 0.066	1.684 ± 0.048	0.461 ± 0.055	0.140 ± 0.052
24420-2000 BU ₂₂	J.Trojan	1	_	2.931 ± 1.742	0.937 ± 0.066	0.441 ± 0.042	0.304 ± 0.063		_	
24426-2000 CR ₁₂	J.Trojan	1	_	5.604 ± 1.660	0.717 ± 0.054	0.414 ± 0.043	0.424 ± 0.062		_	_
24444-2000 OP ₃₂	J.Trojan	1	11.395 ± 0.093	6.264 ± 2.536	0.712 ± 0.071	0.437 ± 0.053	0.409 ± 0.090	_	_	
24452-2000 QU ₁₆₇	J.Trojan	1	11.776 ± 0.099	6.333 ± 2.377	0.872 ± 0.056	0.441 ± 0.056	0.406 ± 0.087	_	_	_
4467-2000 SS ₁₆₅	J.Trojan	1	11.800 ± 0.104	11.221 ± 2.463	0.927 ± 0.079	0.460 ± 0.060	0.513 ± 0.091	_	_	_
4835-1995 SM ₅₅	Classic Hot	10	4.332 ± 0.040	0.272 ± 1.805	0.652 ± 0.032	0.357 ± 0.043	0.356 ± 0.052	1.010 ± 0.050	-0.270 ± 0.163	
4952-1997 OJ ₄	Res. 3:2	4	7.622 ± 0.194	4.696 ± 4.772	0.763 ± 0.114	0.431 ± 0.114	0.396 ± 0.102	_	_	_
4978-1998 HJ ₁₅₁	Classic Cold	2	7.008 ± 0.050	29.628 ± 2.427	1.110 ± 0.030	0.710 ± 0.030	_	_	_	_
5347-1999 RQ ₁₁₆	J.Trojan	1	11.545 ± 0.075	10.900 ± 2.252	0.618 ± 0.058	0.488 ± 0.042	0.461 ± 0.079	_	_	_
6181-1996 GQ ₂₁	Res.(other)	8	4.796 ± 0.051	37.628 ± 1.969	1.011 ± 0.068	0.726 ± 0.043	0.694 ± 0.056	2.438 ± 0.062	0.497 ± 0.082	0.055 ± 0.096
6308-1998 SM ₁₆₅	Res.(other)	8	5.549 ± 0.107	31.425 ± 2.295	0.989 ± 0.114	0.641 ± 0.056	0.666 ± 0.072	2.436 ± 0.002 2.295 ± 0.087	0.546 ± 0.086	0.080 ± 0.07
6308-1998 SM _{165+B}	Res.(other)	1		31.423 ± 2.293 31.037 ± 0.000						
6375-1999 DE ₉	Res.(other)	13	4.809 ± 0.045	20.625 ± 1.625	0.967 ± 0.043	0.579 ± 0.036	0.568 ± 0.043	1.717 ± 0.193	0.327 ± 0.106	0.025 ± 0.10
8958-2001 CQ ₄₂	J.Trojan	13	4.007 ± 0.043	-2.591 ± 1.987	0.730 ± 0.043 0.730 ± 0.054	0.364 ± 0.045	0.308 ± 0.043 0.230 ± 0.071	1.717 ± 0.173	0.327 ± 0.100	0.025 ± 0.10
8978-Ixion	Res. 3:2	8	3.429 ± 0.050	-2.391 ± 1.987 22.481 ± 2.259	0.730 ± 0.034 1.009 ± 0.051	0.504 ± 0.043 0.610 ± 0.030	0.230 ± 0.071 0.580 ± 0.040	1.590 ± 0.064	0.277 ± 0.069	-0.010 ± 0.07
9981-1999 TD ₁₀	Scattered			12.194 ± 1.263		0.510 ± 0.030 0.502 ± 0.032	0.580 ± 0.040 0.511 ± 0.051		0.277 ± 0.009 0.425 ± 0.093	-0.010 ± 0.07
		18	8.693 ± 0.039		0.808 ± 0.095			1.790 ± 0.057		0.102 + 0.06
0698-Hippokoon	J.Trojan	3	11.977 ± 0.125	7.486 ± 2.170	0.715 ± 0.067	0.458 ± 0.048	0.412 ± 0.079	1.527 ± 0.074	0.396 ± 0.080	0.183 ± 0.06
1820-1999 RT ₁₈₆	J.Trojan	1	12.450 ± 0.168	10.495 ± 4.232	0.889 ± 0.093	0.520 ± 0.091	0.396 ± 0.153	_	_	
1821-1999 RK ₂₂₅	J.Trojan	1	11.849 ± 0.173	8.202 ± 3.877	0.980 ± 0.111	0.440 ± 0.097	0.461 ± 0.145	_		
1824-Elatus	Centaurs	14	10.462 ± 0.122	27.348 ± 1.859	1.020 ± 0.060	0.620 ± 0.048	0.630 ± 0.038	_	0.392 ± 0.062	0.020 ± 0.08
2430-2000 RQ ₈₃	J.Trojan	1	12.293 ± 0.074	8.679 ± 1.818	0.772 ± 0.038	0.474 ± 0.045	0.425 ± 0.067			—
2532-Thereus	Centaurs	221	9.218 ± 0.015	10.537 ± 1.186	0.763 ± 0.072	0.501 ± 0.016	0.488 ± 0.036	1.350 ± 0.050	0.442 ± 0.074	0.110 ± 0.07
2615-2001 QU ₂₇₇	J.Trojan	1	11.405 ± 0.050	9.346 ± 1.235	0.807 ± 0.052	0.452 ± 0.028	0.474 ± 0.043			_
2794-1989 UE ₅	J.Trojan	1	12.744 ± 0.094	6.794 ± 2.116	0.923 ± 0.065	0.393 ± 0.056	0.486 ± 0.080	_	_	_
2929-1995 QY ₉	Classic Cold	5	7.365 ± 0.062	18.741 ± 10.170	0.740 ± 0.200	0.561 ± 0.110	_	2.027 ± 0.201	_	_
3001-1997 CU ₂₉	Classic Cold	6	6.216 ± 0.104	28.823 ± 3.429	1.157 ± 0.145	0.645 ± 0.066	0.634 ± 0.092	_	0.410 ± 0.238	_
3128-1998 BU ₄₈	Scattered	9	6.861 ± 0.137	26.535 ± 4.487	1.044 ± 0.071	0.631 ± 0.101	0.644 ± 0.104	2.275 ± 0.059	0.505 ± 0.098	-0.157 ± 0.10
3340-1998 VG ₄₄	Res. 3:2	12	6.322 ± 0.073	20.986 ± 3.166	0.930 ± 0.055	0.573 ± 0.047	0.598 ± 0.098	_	0.398 ± 0.128	0.010 ± 0.07
4785-2001 RG ₈₇	J.Trojan	1	12.513 ± 0.076	4.113 ± 1.599	0.728 ± 0.050	0.386 ± 0.043	0.419 ± 0.060	_	_	
5671-1998 SN ₁₆₅	Classic Cold	6	5.679 ± 0.320	6.857 ± 3.068	0.712 ± 0.095	0.444 ± 0.078	0.437 ± 0.083	1.270 ± 0.050	_	_
8083-Rhadamanth	Classic Hot	1	7.430 ± 0.063	11.286 ± 3.072	0.650 ± 0.085	0.527 ± 0.069	0.412 ± 0.075	_	_	_
8084-1999 HB ₁₂	Res.(other)	3	6.632 ± 0.079	15.815 ± 3.668	0.893 ± 0.064	0.544 ± 0.065	0.481 ± 0.094		_	
8628-Huya	Res. 3:2	58	4.678 ± 0.099	22.400 ± 2.192	0.963 ± 0.042	0.609 ± 0.090	0.593 ± 0.055	1.970 ± 0.050	0.330 ± 0.109	_
9285-2001 BP ₇₅	J.Trojan	1	_	-0.195 ± 1.869	0.810 ± 0.059	0.381 ± 0.044	0.291 ± 0.067	_	_	_
0314-1999 KR ₁₆	Scattered	5	5.612 ± 0.092	42.033 ± 2.385	1.123 ± 0.083	0.738 ± 0.057	0.750 ± 0.040	_	0.560 ± 0.071	0.020 ± 0.07
2301-2001 UR ₁₆₃	Res.(other)	6	3.799 ± 0.102	44.811 ± 2.993	1.290 ± 0.109	0.840 ± 0.048	0.673 ± 0.121	2.381 ± 0.058	0.476 ± 0.115	
2355-Typhon	Scattered	12	7.252 ± 0.054	12.657 ± 1.354	0.758 ± 0.039	0.525 ± 0.022	0.414 ± 0.053	1.560 ± 0.045	0.476 ± 0.113 0.406 ± 0.087	0.160 ± 0.07
2355-Typhon+B	Scattered	1	7.232 ± 0.034 —	9.763 ± 0.000	0.730 ± 0.037					
3212-2000 AL ₁₁₃	J.Trojan	1	_	2.314 ± 1.748	_	0.402 ± 0.043	0.343 ± 0.064	_	_	_
4594-1999 OX ₃	Scattered	30	7.052 ± 0.085	34.216 ± 2.104	-1.138 ± 0.067	0.402 ± 0.043 0.705 ± 0.056	0.642 ± 0.066	2.217 ± 0.127	0.399 ± 0.105	-0.023 ± 0.09
-	Classic Cold		1.032 ± 0.083		1.136 ± 0.007	0.705 ± 0.030	0.042 ± 0.000	2.21/ ± 0.12/	0.399 ± 0.103 0.428 ± 0.291	-0.023 ± 0.05
5802-2000 PV ₂₉		2	4 050 + 0 060	20 224 + 1 729	1.020 + 0.047	0.602 + 0.022	0.610 + 0.040	2 215 + 0 101		
7171-1999 TC ₃₆	Res. 3:2	23	4.858 ± 0.060	29.334 ± 1.738	1.029 ± 0.047	0.693 ± 0.032	0.619 ± 0.049	2.215 ± 0.101	0.337 ± 0.092	0.009 ± 0.09
7171-1999 TC _{36+B}	Res. 3:2	1	<u> </u>	19.177 ± 0.000	0.004 : 0.050	0.622 - 0.022	0.617 : 0.054	1.711 - 0.070	0.265 : 0.100	
7932-2000 GN ₁₇₁	Res. 3:2	13	6.183 ± 0.045	26.540 ± 1.817	0.924 ± 0.059	0.622 ± 0.038	0.617 ± 0.054	1.711 ± 0.079	0.365 ± 0.108	0.060 ± 0.09
17967-2000 SL ₂₉₈	J.Trojan	1	11.843 ± 0.113	11.415 ± 2.726	0.899 ± 0.058	0.489 ± 0.069	0.476 ± 0.102	_	_	_

Table A.1. continued.

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	Grt $\pm \sigma$	B-V ±σ	V-R $\pm \sigma$	R-I ±σ	V-J ±σ	J-H $\pm\sigma$	H-K $\pm\sigma$
48249-2001 SY ₃₄₅	J.Trojan	1	12.550 ± 0.072	11.716 ± 1.707	0.758 ± 0.050	0.530 ± 0.037	0.420 ± 0.060	_	_	_
48252-2001 TL ₂₁₂	J.Trojan	1	12.719 ± 0.153	8.694 ± 3.314	0.949 ± 0.100	0.467 ± 0.093	0.436 ± 0.129		_	_
18639-1995 TL ₈	Scattered	4	4.664 ± 0.087	22.529 ± 6.004	1.008 ± 0.159	0.621 ± 0.091	0.551 ± 0.120	2.420 ± 0.050	0.412 ± 0.131	-0.020 ± 0.083
19036-Pelion	Centaurs	3	10.420 ± 0.247	8.735 ± 3.519	0.746 ± 0.080	0.556 ± 0.078	0.368 ± 0.102	_	_	_
50000-Quaoar	Classic Hot	4	2.220 ± 0.029	29.224 ± 1.706	0.958 ± 0.035	0.650 ± 0.020	0.610 ± 0.028	2.180 ± 0.058	0.360 ± 0.050	0.030 ± 0.05
51359-2000 SC ₁₇	J.Trojan	1	11.985 ± 0.233	7.745 ± 5.725	0.864 ± 0.201	0.447 ± 0.131	0.438 ± 0.210		_	
52747-1998 HM ₁₅₁	Classic Cold	1	7.417 ± 0.100	24.891 ± 4.691	0.930 ± 0.090	0.620 ± 0.050	_		_	
52872-Okyrhoe	Centaurs	18	10.768 ± 0.093	10.846 ± 2.450	0.743 ± 0.065	0.486 ± 0.061	0.474 ± 0.070		0.457 ± 0.068	0.159 ± 0.080
52975-Cyllarus	Centaurs	9	8.634 ± 0.100	32.286 ± 3.654	1.096 ± 0.095	0.680 ± 0.085	0.669 ± 0.090	2.425 ± 0.077	0.450 ± 0.092	-0.100 ± 0.08
53469-2000 AX ₈	J.Trojan	1	_	0.771 ± 1.818	0.663 ± 0.051	0.356 ± 0.045	0.361 ± 0.067	_	_	_
54520-2000 PJ ₃₀	Scattered	1	_	_	_	_	_	_	0.410 ± 0.304	-0.100 ± 0.08 -0.153 ± 0.099
64598-Bienor	Centaurs	15	7.573 ± 0.104	8.793 ± 2.427	0.711 ± 0.059	0.476 ± 0.046	0.400 ± 0.079	1.684 ± 0.091	0.379 ± 0.078	0.153 ± 0.099
55565-2002 AW ₁₉₇	Classic Hot	9	3.087 ± 0.038	22.129 ± 1.429	0.913 ± 0.041	0.602 ± 0.031	0.581 ± 0.037	1.740 ± 0.054	0.329 ± 0.070	0.215 ± 0.094
55576-Amycus	Centaurs	19	7.774 ± 0.047	33.885 ± 1.517	1.112 ± 0.041	0.702 ± 0.031 0.702 ± 0.030	0.668 ± 0.039		0.299 ± 0.077	0.004 ± 0.064
55636-2002 TX ₃₀₀	Classic Hot	8	3.237 ± 0.047	-0.195 ± 1.517	0.679 ± 0.038	0.762 ± 0.030 0.359 ± 0.024	0.323 ± 0.041	_	0.299 ± 0.077	J.00+ ± 0.00·
55637-2002 UX ₂₅	Classic Hot	8	3.497 ± 0.043	15.518 ± 1.290	1.007 ± 0.038 1.007 ± 0.043	0.539 ± 0.024 0.540 ± 0.030	0.323 ± 0.041 —	_	0.404 ± 0.049	
55638-2002 VE ₉₅	Res. 3:2	2	3.497 ± 0.020 —	36.638 ± 2.876	1.007 ± 0.043 1.127 ± 0.074	0.540 ± 0.050 —	_	2.220 ± 0.057	0.404 ± 0.049 0.400 ± 0.057	0.020 ± 0.06
7.0		1	-11.534 ± 0.059	13.013 ± 1.374		0.494 ± 0.033	0.509 ± 0.049			0.020 ± 0.00
56968-2000 SA ₉₂	J.Trojan	-			0.986 ± 0.040			_	0.400 + 0.240	_
58534-Logos	Classic Cold	7	6.758 ± 0.181	26.351 ± 4.789	0.990 ± 0.127	0.729 ± 0.120	0.602 ± 0.119		0.498 ± 0.340	_
58534-Logos+B	Classic Cold	1		17.578 ± 0.000				_		_
59358-1999 CL ₁₅₈	Scattered	1	6.653 ± 0.090	5.546 ± 2.064	0.800 ± 0.060	0.390 ± 0.040	0.470 ± 0.060		_	_
50454-2000 CH ₁₀₅	Classic Cold	2	6.376 ± 0.093	25.350 ± 3.158	1.019 ± 0.086	0.643 ± 0.096	0.583 ± 0.095	_	_	_
60458-2000 CM ₁₁₄	Detached	2	6.972 ± 0.040	11.545 ± 1.858	0.730 ± 0.020	0.500 ± 0.020	_	_	_	
50458-2000 CM _{114+B}	Detached	1		13.041 ± 0.000	_	_	_	_	_	_
60558-Echeclus	Centaurs	8	9.555 ± 0.155	10.399 ± 3.404	0.854 ± 0.081	0.465 ± 0.076	0.486 ± 0.095	1.480 ± 0.072	0.466 ± 0.125	0.340 ± 0.08
50608-2000 EE ₁₇₃	Scattered	3	8.147 ± 0.180	13.198 ± 1.827	0.665 ± 0.032	0.488 ± 0.029	0.543 ± 0.056	_	_	_
50620-2000 FD ₈	Res.(other)	3	6.507 ± 0.221	30.813 ± 3.016	1.151 ± 0.110	0.664 ± 0.078	0.648 ± 0.070		_	
50621-2000 FE ₈	Res.(other)	3	6.510 ± 0.062	11.433 ± 1.500	0.750 ± 0.024	0.480 ± 0.020	0.500 ± 0.050	_	0.470 ± 0.277	_
53252-2001 BL ₄₁	Centaurs	10	11.278 ± 0.063	13.807 ± 2.421	0.718 ± 0.049	0.509 ± 0.044	0.381 ± 0.160	_	0.365 ± 0.071	0.255 ± 0.10
55150-2002 CA ₁₂₆	J.Trojan	1	_	3.241 ± 1.724	0.651 ± 0.051	0.377 ± 0.044	0.407 ± 0.064		_	
55225-2002 EK ₄₄	J.Trojan	1	_	1.982 ± 1.783	0.693 ± 0.052	0.401 ± 0.044	0.334 ± 0.065		_	
55489-Ceto	Scattered	3	6.286 ± 0.041	15.666 ± 0.871	0.860 ± 0.030	0.560 ± 0.030	_	_	_	_
55489-Ceto+B	Scattered	1	_	13.527 ± 0.000	_	_	_	_	_	_
66452-1999 OF ₄	Classic Cold	3	6.323 ± 0.080	26.708 ± 3.878	1.032 ± 0.106	0.673 ± 0.083	0.601 ± 0.080	_	0.380 ± 0.253	_
66652-Borasisi	Classic Cold	7	5.418 ± 0.056	33.830 ± 2.714	0.820 ± 0.170	0.646 ± 0.058	0.647 ± 0.062	2.010 ± 0.067	0.457 ± 0.162	0.100 ± 0.094
66652-Borasisi+B	Classic Cold	1		41.482 ± 0.000	0.020 ± 0.170 —	0.040 ± 0.030	0.047 ± 0.002 —	2.010 ± 0.007		0.100 ± 0.05
59986-1998 WW ₂₄	Res. 3:2	3	7.963 ± 0.085	17.966 ± 2.895	0.756 ± 0.112	0.463 ± 0.081	0.659 ± 0.104		_	
59987-1998 WA ₂₅	Classic Cold	1	7.703 ± 0.003	17.056 ± 0.000 17.056 ± 0.000	0.750 ± 0.112	0.403 ± 0.001	0.037 ± 0.104		_	
59988-1998 WA ₂₅	Res.(other)	2	7.352 ± 0.122	17.030 ± 0.000 13.652 ± 6.704	0.786 ± 0.129	0.492 ± 0.133	0.530 ± 0.183	_	_	_
51	, ,	1								_
59990-1998 WU ₃₁	Res. 3:2		8.204 ± 0.073	23.278 ± 3.272	0.720 ± 0.089	0.505 ± 0.086	0.722 ± 0.087	_	$-$ 0.425 \pm 0.105	_
73480-2002 PN ₃₄	Scattered	10	8.506 ± 0.030	12.613 ± 1.134	0.818 ± 0.095	0.520 ± 0.020			0.425 ± 0.105	_
6804-2000 QE	J.Trojan	1	12.095 ± 0.135	7.871 ± 2.878	0.803 ± 0.082	0.446 ± 0.070	0.443 ± 0.106		0.256 + 0.150	— 0.104 · 0.17
79360-1997 CS ₂₉	Classic Cold	15	4.990 ± 0.060	28.475 ± 2.382	1.055 ± 0.079	0.666 ± 0.054	0.609 ± 0.057	2.073 ± 0.134	0.356 ± 0.150	0.104 ± 0.176
79360-1997 CS _{29+B}	Classic Cold	1		31.690 ± 0.000	_			_	_	_
79978-1999 CC ₁₅₈	Res.(other)	2	5.355 ± 0.097	24.869 ± 2.974	0.996 ± 0.066	0.611 ± 0.052	0.619 ± 0.075	_		
79983-1999 DF ₉	Classic Hot	1	5.797 ± 0.110	33.944 ± 2.164	0.920 ± 0.060	0.710 ± 0.050	0.650 ± 0.060	_	_	_
30806-2000 CM ₁₀₅	Classic Cold	2	_	32.348 ± 0.000	_	_	_	_	0.360 ± 0.256	_
32075-2000 YW ₁₃₄	Res.(other)	14	4.382 ± 0.046	16.483 ± 2.187	0.922 ± 0.053	0.503 ± 0.052	0.581 ± 0.059	_	0.358 ± 0.153	_
32155-2001 FZ ₁₇₃	Scattered	9	5.813 ± 0.024	15.975 ± 1.446	0.864 ± 0.023	0.546 ± 0.026	0.508 ± 0.036	_	0.171 ± 0.118	

Table A.1. continued.

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	Grt $\pm \sigma$	B-V ±σ	V-R ±σ	R-I ±σ	V-J ±σ	J-H $\pm\sigma$	H-K $\pm \sigma$
32158-2001 FP ₁₈₅	Scattered	7	5.947 ± 0.054	16.009 ± 2.281	0.820 ± 0.048	0.572 ± 0.038	0.458 ± 0.078	_	_	_
3982-Crantor	Centaurs	17	8.662 ± 0.056	39.225 ± 1.992	1.105 ± 0.042	0.761 ± 0.039	0.667 ± 0.055		0.380 ± 0.067	-0.140 ± 0.06
4522-2002 TC ₃₀₂	Res.(other)	2	3.719 ± 0.020	28.979 ± 1.356	1.120 ± 0.028	0.640 ± 0.020	0.660 ± 0.020	_	_	_
4709-2002 VW ₁₂₀	J.Trojan	1	12.446 ± 0.158	12.795 ± 3.419	0.855 ± 0.087	0.462 ± 0.090	0.548 ± 0.130		_	_
35627-1998 HP ₁₅₁	Classic Cold	1	_	25.429 ± 0.000	_	_	_		_	_
5633-1998 KR ₆₅	Classic Cold	4	6.596 ± 0.206	30.583 ± 2.821	1.095 ± 0.080	0.628 ± 0.066	0.790 ± 0.081	_	_	_
6047-1999 OY ₃	Classic Hot	5	6.396 ± 0.147	-3.527 ± 2.148	0.726 ± 0.037	0.345 ± 0.046	0.277 ± 0.076	_	-0.380 ± 0.256	_
6177-1999 RY ₂₁₅	Classic Hot	3	7.429 ± 0.077	3.839 ± 3.476	0.719 ± 0.123	0.358 ± 0.090	0.631 ± 0.185	_	0.470 ± 0.286	_
7269-2000 OO ₆₇	Scattered	3	9.027 ± 0.155	26.481 ± 2.592	1.080 ± 0.092	0.654 ± 0.081	0.593 ± 0.055		_	
7555-2000 QB ₂₄₃	Scattered	6	8.437 ± 0.129	5.554 ± 5.948	0.763 ± 0.086	0.383 ± 0.083	0.729 ± 0.198	1.480 ± 0.108	0.592 ± 0.152	
8269-2001 KF ₇₇	Centaurs	1	10.038 ± 0.020	37.355 ± 1.301	1.080 ± 0.040	0.730 ± 0.010	_	_	_	_
8611-Teharonhia	Classic Cold	1	_	5.020 ± 0.000	_	_	_	_	_	_
8611-Teharonhia+B	Classic Cold	1	_	4.612 ± 0.000	_	_	_	_	_	_
0377-Sedna	Detached	3	1.077 ± 0.065	32.954 ± 2.972	1.131 ± 0.079	0.686 ± 0.077	0.657 ± 0.067	2.320 ± 0.060	0.290 ± 0.222	0.050 ± 0.314
0482-Orcus	Res. 3:2	6	1.982 ± 0.099	2.761 ± 1.831	0.664 ± 0.041	0.370 ± 0.039	0.390 ± 0.045	1.070 ± 0.042	0.120 ± 0.022	0.050 ± 0.051 0.053 ± 0.053
0568-2004 GV ₉	Classic Hot	2		20.819 ± 1.099	0.843 ± 0.028	_	—	1.570 ± 0.042 1.570 ± 0.058	0.340 ± 0.094	0.140 ± 0.09
1133-1998 HK ₁₅₁	Res. 3:2	6	6.940 ± 0.075	9.775 ± 2.589	0.645 ± 0.020 0.645 ± 0.121	0.520 ± 0.062	0.390 ± 0.061	1.570 ± 0.092	—	
1205-1998 US ₄₃	Res. 3:2	2	7.851 ± 0.078	4.463 ± 3.403	0.691 ± 0.102	0.446 ± 0.095	0.347 ± 0.094	- 1.570 ± 0.072	_	_
1554-1999 RZ ₂₁₅	Scattered	1	8.069 ± 0.079	19.081 ± 3.401	0.771 ± 0.02	0.575 ± 0.090	0.539 ± 0.091	_	<u> </u>	_
5626-2002 GZ ₃₂	Centaurs	18	6.733 ± 0.156	16.992 ± 2.947	0.674 ± 0.043	0.576 ± 0.056 0.576 ± 0.054	0.539 ± 0.091 0.538 ± 0.067	_	0.442 ± 0.113	_
9328-2001 UY ₁₂₃	J.Trojan	1	12.610 ± 0.103	12.496 ± 2.292	0.890 ± 0.058	0.570 ± 0.054 0.537 ± 0.056	0.434 ± 0.084	_	U.442 ± U.113	_
05685-2000 SC ₅₁	J.Trojan J.Trojan	1	12.010 ± 0.103 12.721 ± 0.097	8.131 ± 2.099	1.016 ± 0.055	0.337 ± 0.030 0.444 ± 0.059	0.454 ± 0.084 0.452 ± 0.081	_	_	_
11113-2001 VK ₈₅	J.Trojan J.Trojan	1	12.721 ± 0.097 12.760 ± 0.086	13.245 ± 1.864	0.822 ± 0.063	0.444 ± 0.039 0.462 ± 0.048	0.452 ± 0.061 0.558 ± 0.069	_	_	
05	Res. 3:2	7	7.245 ± 0.195	38.250 ± 4.976	0.822 ± 0.003 1.186 ± 0.118		0.338 ± 0.009 0.746 ± 0.113	2.435 ± 0.128	_	_
18228-1996 TQ ₆₆	Classic Hot	3				0.670 ± 0.096				_
18379-1999 HC ₁₂		3 1	7.856 ± 0.125	12.620 ± 6.899	0.894 ± 0.163	0.490 ± 0.138	0.343 ± 0.125	_	0.490 ± 0.298	
18702-2000 OM ₆₇	Detached		7.075 ± 0.030	14.444 ± 1.982	0.820 ± 0.057	0.470 ± 0.040	0.590 ± 0.040	_	_	_
19068-2001 KC ₇₇	Res.(other)	1	6.822 ± 0.030	18.607 ± 1.301	0.910 ± 0.010	0.560 ± 0.010	_	_		
19070-2001 KP ₇₇	Res.(other)	1		0.075 - 1.201			_	_	0.370 ± 0.235	_
19315-2001 SQ ₇₃	Centaurs	1	8.885 ± 0.030	8.875 ± 1.301	0.670 ± 0.020	0.460 ± 0.010	_			_
19951-2002 KX ₁₄	Classic Cold	3	_	26.057 ± 1.557	1.050 ± 0.030	0.610 ± 0.020	_	_	_	_
19979-2002 WC ₁₉	Res.(other)	1		28.487 ± 0.000			_	_		_
20061-2003 CO ₁	Centaurs	11	9.208 ± 0.040	11.701 ± 2.057	0.740 ± 0.030	0.490 ± 0.020			0.326 ± 0.091	
20132-2003 FY ₁₂₈	Detached	3	4.476 ± 0.010	21.441 ± 1.266	1.050 ± 0.028	0.600 ± 0.020	0.550 ± 0.030	1.640 ± 0.058	0.360 ± 0.078	0.110 ± 0.08
20178-2003 OP ₃₂	Classic Hot	1	_	3.135 ± 2.090	0.698 ± 0.052	_	_	_	_	_
20347-Salacia	Classic Hot	1		7.116 ± 0.000		_	_			_
20347-Salacia+B	Classic Hot	1		7.982 ± 0.000	_	_	_		_	_
20348-2004 TY ₃₆₄	Res. 3:2	1	_	28.425 ± 3.958	1.059 ± 0.089	_	_	_	_	_
20453-1988 RE ₁₂	J.Trojan	1	13.202 ± 0.189	6.453 ± 3.884	0.826 ± 0.132	0.388 ± 0.108	0.483 ± 0.151	_	_	_
21725-1999 XX ₁₄₃	Centaurs	10	8.522 ± 0.123	30.600 ± 5.205	1.060 ± 0.089	0.648 ± 0.116	0.679 ± 0.123	_	_	_
23509-2000 WK ₁₈₃	Classic Cold	1	_	29.751 ± 0.000	_	_	_	_	_	_
23509-2000 WK _{183+B}	Classic Cold	1	_	26.029 ± 0.000	_	_	_	_	_	_
24729-2001 SB ₁₇₃	J.Trojan	1	12.556 ± 0.107	10.320 ± 2.774	0.992 ± 0.060	0.503 ± 0.064	0.424 ± 0.101		_	
27546-2002 XU ₉₃	Scattered	1	7.943 ± 0.010	5.447 ± 1.087	0.760 ± 0.028	0.440 ± 0.020	0.380 ± 0.020		_	_
29772-1999 HR ₁₁	Classic Cold	1	_	28.264 ± 4.298	0.920 ± 0.120	0.530 ± 0.100	0.800 ± 0.070	_	_	_
30391-2000 JG ₈₁	Res.(other)	1	7.755 ± 0.115	3.743 ± 4.431	_	0.370 ± 0.128	0.430 ± 0.134	_	_	_
31695-2001 XS ₂₅₄	Res.(other)	1	_	_	_	_	_		0.420 ± 0.260	
34340-Pluto	Res. 3:2	4	-0.881 ± 0.400	7.607 ± 0.681	0.867 ± 0.016	0.515 ± 0.035	0.400 ± 0.010		_	_
34340-Pluto+B	Res. 3:2	1	_	3.334 ± 0.000	0.710 ± 0.002	_	_	_	_	_
34340-Pluto+C	Res. 3:2	1	_	-2.234 ± 0.000	0.644 ± 0.028	_	_	_	_	_

Table A.1. continued.

Object	Class ⁽¹⁾	Epochs ⁽²⁾	M11 ±σ	Grt ±σ	B-V ±σ	V-R ±σ	R-I ±σ	V-J ±σ	J-H ±σ	H-K ±σ
134340-Pluto+D	Res. 3:2	1	_	18.074 ± 0.000	0.907 ± 0.031	_	_	_	_	_
134860-2000 OJ ₆₇	Classic Cold	5	6.001 ± 0.120	26.378 ± 3.102	1.050 ± 0.060	0.670 ± 0.050	0.600 ± 0.070	_	0.305 ± 0.162	0.070 ± 0.139
134860-2000 OJ _{67+B}	Classic Cold	1	_	33.683 ± 0.000	_	_	_	_	_	
135182-2001 QT ₃₂₂	Classic Cold	1	_	15.584 ± 11.076	0.710 ± 0.060	0.530 ± 0.120	_	_	_	
136108-Haumea	Classic Hot	3	0.217 ± 0.030	-0.010 ± 0.850	0.631 ± 0.025	0.370 ± 0.020	0.320 ± 0.020	1.051 ± 0.020	-0.044 ± 0.037	-0.111 ± 0.048
136199-Eris	Detached	64	-1.462 ± 0.036	3.866 ± 0.823	0.805 ± 0.015	0.389 ± 0.049	0.363 ± 0.061	0.849 ± 0.108	0.080 ± 0.072	-0.280 ± 0.085
136472-Makemake	Classic Hot	1	_	4.693 ± 0.900	0.828 ± 0.022		_	_	_	_
137294-1999 RE ₂₁₅	Classic Cold	3	6.415 ± 0.187	30.557 ± 3.448	1.003 ± 0.130	0.710 ± 0.074	0.571 ± 0.089	_	_	_
137295-1999 RB ₂₁₆	Res.(other)	1	7.670 ± 0.087	14.468 ± 2.704	0.897 ± 0.122	0.522 ± 0.073	0.506 ± 0.061	_	_	_
138537-2000 OK ₆₇	Classic Cold	5	6.082 ± 0.083	20.354 ± 3.262	0.821 ± 0.123	0.583 ± 0.072	0.524 ± 0.079	2.421 ± 0.088	0.455 ± 0.145	0.040 ± 0.078
144897-2004 UX ₁₀	Res. 3:2	1	_	20.664 ± 2.909	0.950 ± 0.020	0.580 ± 0.030	_	_	_	_
145451-2005 RM ₄₃	Scattered	2	_	1.330 ± 1.474	0.590 ± 0.038	_	_	0.910 ± 0.067	0.200 ± 0.085	0.280 ± 0.092
145452-2005 RN ₄₃	Classic Hot	2	_	16.523 ± -9.000	_	_	_	1.550 ± 0.058	0.310 ± 0.071	0.190 ± 0.071
145453-2005 RR ₄₃	Classic Hot	3	_	0.109 ± 3.025	0.790 ± 0.076	_	_	1.020 ± 0.067	_	_
145480-2005 TB ₁₉₀	Detached	1	4.169 ± 0.020	18.632 ± 1.525	0.980 ± 0.042	0.560 ± 0.030	0.550 ± 0.030	_	_	_
148209-2000 CR ₁₀₅	Detached	3	6.228 ± 0.040	14.424 ± 3.398	0.771 ± 0.082	0.509 ± 0.048	0.590 ± 0.090	_	0.410 ± 0.297	_
148780-Altjira	Classic Hot	1	_	29.751 ± 0.000	_		_	_	_	_
148780-Altjira+B	Classic Hot	1	_	35.732 ± 0.000	_	_	_	_	_	_
150642-2001 CZ ₃₁	Classic Hot	1	_	_	_	_	_	_	0.550 ± 0.099	0.160 ± 0.099
163135-2002 CT ₂₂	J.Trojan	1	_	1.927 ± 1.789	0.690 ± 0.053	0.382 ± 0.045	0.360 ± 0.066	_	_	_
163216-2002 EN ₆₈	J.Trojan	1	_	4.392 ± 1.713	_	0.443 ± 0.042	0.347 ± 0.062	_	_	_
168700-2000 GE ₁₄₇	Res. 3:2	1	8.002 ± 0.110	16.667 ± 3.950	_	0.550 ± 0.120	0.520 ± 0.114	_	_	_
168703-2000 GP ₁₈₃	Classic Cold	5	5.756 ± 0.060	8.353 ± 2.654	0.669 ± 0.074	0.445 ± 0.053	0.463 ± 0.066	_	_	_
181708-1993 FW	Classic Hot	8	6.527 ± 0.122	18.980 ± 3.484	1.023 ± 0.069	0.583 ± 0.070	0.439 ± 0.104	_	0.400 ± 0.250	_
181855-1998 WT ₃₁	Classic Hot	3	7.440 ± 0.088	10.112 ± 6.721	0.751 ± 0.144	0.502 ± 0.121	0.416 ± 0.166	_	_	_
181867-1999 CV ₁₁₈	Res.(other)	1	7.186 ± 0.063	31.744 ± 2.469	_	0.733 ± 0.067	0.582 ± 0.062	_	_	_
181871-1999 CO ₁₅₃	Classic Cold	1	_	38.554 ± 0.000	_	_	_	_	_	_
181874-1999 HW ₁₁	Detached	2	6.672 ± 0.077	12.562 ± 2.766	0.840 ± 0.042	0.499 ± 0.073	0.493 ± 0.083	_	_	
181902-1999 RD ₂₁₅	Detached	2	7.860 ± 0.079	7.547 ± 0.000	_	_	0.498 ± 0.058	_	_	_
182397-2001 QW ₂₉₇	Classic Hot	1	6.660 ± 0.050	25.220 ± 3.210	1.020 ± 0.099	0.580 ± 0.070	0.670 ± 0.060	_	_	- - - - - - - - -
182933-2002 GZ ₃₁	Detached	1	_	33.013 ± 0.000	_	_	_	_	_	_
192388-1996 RD ₂₉	J.Trojan	1	_	4.691 ± 1.684	0.741 ± 0.055	0.421 ± 0.042	0.388 ± 0.062	_	_	_
192929-2000 AT ₄₄	J.Trojan	1	_	-0.624 ± 1.876	0.707 ± 0.053	0.354 ± 0.045	0.318 ± 0.068		_	
208996-2003 AZ ₈₄	Res. 3:2	9	3.227 ± 0.050	23.884 ± 2.812	0.757 ± 0.146	0.620 ± 0.030	0.560 ± 0.042	1.210 ± 0.054	0.188 ± 0.165	0.060 ± 0.071

Online Material p 12

Table A.2. References used for each object, and number of epochs.

Object	References	Nr.
C/1983 H1 IRAS-Araki-Alcock	Lamy et al. (2004)	1
C/1983 J1 Sugano-Saigusa-Fujikawa	Lamy et al. (2004)	1
C/1984 K1 Shoemaker	Lamy et al. (2004)	1
C/1984 U1 Shoemaker	Lamy et al. (2004)	1
C/1986 P1 Wilson	Lamy et al. (2004)	1
C/1987 A1 Levy C/1987 H1 Shoemaker	Lamy et al. (2004) Lamy et al. (2004)	1 1
C/1988 B1 Shoemaker	Lamy et al. (2004)	1
C/1988 C1 Maury-Phinney	Lamy et al. (2004)	1
C/1995 O1 Hale-Bopp	Lamy & Toth (2009)	
, 11	Lamy et al. (2004)	2
C/1996 B2 Hyakutake	Lamy et al. (2004)	1
C/1999 S4 LINEAR	Lamy et al. (2004)	1
C/2000 B4 LINEAR	Bauer et al. (2003)	6
C/2001 M10 NEAT	Jewitt (2009b)	2
C/2001 OG108 LONEOS	Lamy et al. (2004)	1
P/1991 L3 Levy	Lamy et al. (2004)	1
P/1993 W1 Mueller	Lamy et al. (2004)	1
P/1994 A1 Kushida P/1994 J3 Shoemaker	Lamy et al. (2004)	1 1
P/1994 J3 Shoemaker P/1995 A1 Jedicke	Lamy et al. (2004) Lamy et al. (2004)	1
P/1996 A1 Jedicke	Lamy et al. (2004)	1
P/1997 C1 Gehrels	Lamy et al. (2004)	1
P/1997 G1 Montani	Lamy et al. (2004)	1
P/1997 V1 Larsen	Lamy et al. (2004)	1
P/1998 S1 LM	Lamy et al. (2004)	1
P/1999 D1 Hermann	Lamy et al. (2004)	1
P/1999 RO28 LONEOS	Lamy et al. (2004)	1
P/2004 A1	Jewitt (2009b)	1
1P/Halley	Lamy et al. (2004)	
AD /5	Keller & Thomas (1989)	2
2P/Encke	Jewitt (2002b)	
	Meech et al. (2004)	
	Lamy et al. (2004) Luu & Jewitt (1990)	6
4P/Faye	Lamy et al. (2004)	Ü
41/1 dyC	Lamy & Toth (2009)	2
6P/d'Arrest	Meech et al. (2004)	-
,	Jewitt (2002b)	
	Lamy et al. (2004)	3
7P/Pons-Winnecke	Snodgrass et al. (2005)	
	Lamy et al. (2004)	2
8P/Tuttle	Lamy et al. (2004)	
OD/ID 1.1	Lamy & Toth (2009)	2
9P/Tempel 1	Meech et al. (2004)	1.4
10D/Tompol 2	Lamy et al. (2004)	14
10P/Tempel 2	Jewitt & Luu (1989) Meech et al. (2004)	
	Lamy et al. (2004)	
	Jewitt & Meech (1988)	
	Lamy & Toth (2009)	10
14P/Wolf	Snodgrass et al. (2005)	
	Lamy et al. (2004)	2
15P/Finlay	Lamy et al. (2004)	1
16P/Brooks2	Lamy et al. (2004)	1
17P/Holmes	Lamy et al. (2004)	2
10D/D II	Lamy & Toth (2009)	2
19P/Borrelly	Lamy et al. (2004)	1
21P/GZ	Lamy et al. (2004)	2
22P/Kopff	Luu (1993) Meech et al. (2004)	2
221/Kopii	Lamy et al. (2004)	
	Lamy et al. (2004)	
	Lamy & Toth (2009)	4
24P/Schaumasse	Lamy et al. (2004)	1
26P/GS	Boehnhardt et al. (1999)	
	Lamy et al. (2004)	2

Table A.2. Continued

Object	References	Nr
28P/Neujmin 1	Jewitt & Meech (1988)	
	Delahodde et al. (2001)	
	Meech et al. (2004)	
	Lamy et al. (2004)	
200/61	Campins et al. (1987)	11
29P/Schwassmann-Wachmann 1	Bauer et al. (2003)	
	Jewitt (2009b)	2
20D/D : 411	Lamy et al. (2004)	3
30P/Reinmuth1	Lamy et al. (2004)	1
31P/Schwassmann-Wachmann 2	Lamy et al. (2004)	1
32P/ComasSola	Lamy et al. (2004)	1
33P/Daniel	Lamy et al. (2004)	1
36P/Whipple	Lamy et al. (2004)	1
37P/Forbes	Lamy et al. (2004)	2
2017/04	Lamy & Toth (2009)	2
39P/Oterma	Bauer et al. (2003)	
	Jewitt (2009b)	2
40D/57 : 1 1	Lamy et al. (2004)	3
40P/Vaisala 1	Lamy et al. (2004)	1
41P/Tuttle-Giacobini-Kresak	Lamy et al. (2004)	1
42P/Neujmin 3	Lamy et al. (2004)	1
43P/Wolf-Harrington	Lamy et al. (2004)	1
14P/Reinmuth 2	Lamy et al. (2004)	
45D(T) 1.161 D.11.6(1)	Lamy & Toth (2009)	2
45P/Honda-Mrkos-Pajduŝáková	Lamy et al. (2004)	2
A CD TITL	Lamy & Toth (2009)	2
46P/Wirtanen	Meech et al. (2004)	
	Lamy et al. (2004)	
45D/4 11 1 1 1	Lamy & Toth (2009)	3
47P/Ashbrook-Jackson	Lamy et al. (2004)	_
	Lamy & Toth (2009)	2
48P/Johnson	Lamy et al. (2004)	1
49P/Arend-Rigaux	Lowry et al. (2003)	
	Lamy et al. (2004)	_
-07/4	Millis et al. (1988)	3
50P/Arend	Lamy et al. (2004)	
*1D#1	Lamy & Toth (2009)	2
51P/Harrington	Lamy et al. (2004)	1
52P/Harrington-Abell	Lamy et al. (2004)	1
53P/Van Biesbroeck	Meech et al. (2004)	2
55D/T 1 T1	Lamy et al. (2004)	2
55P/Tempel-Tuttle	Lamy et al. (2004)	2
CD/01 14 D 1	Lamy & Toth (2009)	2
56P/Slaughter-Burnham	Lamy et al. (2004)	1
57P/du Toit-Neujmin-Delporte	Lamy et al. (2004)	1
58P/Jackson-Neujmin	Lamy et al. (2004)	1
59P/Kearns-Kwee	Lamy et al. (2004)	
COD/FE 1: 1 2	Lamy & Toth (2009)	2
60P/Tsuchinshan 2	Lamy et al. (2004)	1
61P/Shajn-Schaldach	Lamy et al. (2004)	1
52P/Tsuchinshan 1	Lamy et al. (2004)	1
53P/Wild 1	Lamy et al. (2004)	2
(AD/C:f+ C-1, 1	Lamy & Toth (2009)	2
64P/Swift-Gehrels	Lamy et al. (2004)	1
55P/Gunn	Lamy et al. (2004)	1
67P/Churyumov-Gerasimenko	Lamy et al. (2004)	_
COD/IZ11-	Lamy & Toth (2009)	2
58P/Klemola	Lamy et al. (2004)	1
59P/Taylor	Lamy et al. (2004)	1
70P/Kojima	Lamy et al. (2004)	^
TIDIOL I	Lamy & Toth (2009)	2
71P/Clark	Lamy et al. (2004)	_
70D/D : E ::1	Lamy & Toth (2009)	2
72P/Denning-Fujikawa	Lamy et al. (2004)	1
73P/Schwassmann-Wachmann 3	Boehnhardt et al. (1999)	1
73P/Schwassmann-Wachmann 3 C 74P/Smirnova-Chernykh	Lamy et al. (2004)	1
	Lamy et al. (2004)	1

Table A.2. Continued

Object	References	Nr.
75P/Kohoutek	Lamy et al. (2004)	1
76P/West-Kohoutek-Ikemura	Lamy et al. (2004)	1
77P/Longmore	Lamy et al. (2004)	1
78P/Gehrels 2	Lamy et al. (2004)	1
79P/du Toit-Hartley	Lamy et al. (2004)	1
81P/Wild 2	Lamy et al. (2004)	1
82P/Gehrels 3	Lamy et al. (2004)	1
84P/Giclas	Lamy et al. (2004) Lamy & Toth (2009)	2
86P/Wild 3	Meech et al. (2004)	2
801 / Wild 3	Lamy et al. (2004)	
	Lamy & Toth (2009)	3
87P/Bus	Meech et al. (2004)	3
0,1,245	Lamy et al. (2004)	2
88P/Howell	Lamy et al. (2004)	1
89P/Russell 2	Lamy et al. (2004)	1
90P/Gehrels 1	Lamy et al. (2004)	1
91P/Russell 3	Lamy et al. (2004)	1
92P/Sanguin	Snodgrass et al. (2005)	
	Lamy et al. (2004)	2
93P/Lovas 1	Meech et al. (2004)	1
94P/Russell 4	Lamy et al. (2004)	1
96P/Machholz 1	Meech et al. (2004)	
	Lamy et al. (2004)	2
97P/Metcalf-Brewington	Lamy et al. (2004)	1
98P/Takamizawa	Lamy et al. (2004)	1
99P/Kowal 1	Lamy et al. (2004)	1
100P/Hartley 1	Lamy et al. (2004)	1
101P/Chernykh	Lamy et al. (2004)	1
103P/Hartley 2	Lamy et al. (2004)	1
104P/Kowal 2 105P/Singer-Brewster	Lamy et al. (2004) Lamy et al. (2004)	1 1
106P/Schuster	Lamy et al. (2004) Lamy et al. (2004)	1
1001/Schuster	Lamy & Toth (2009)	2
107P/Wilson-Harrington	Meech et al. (2004)	2
10/1/ Wilson Hairington	Lowry & Weissman (2003)	
	Lamy et al. (2004)	
	Chamberlin et al. (1996)	6
109P/Swift-Tuttle	Green et al. (1997)	
	Lamy et al. (2004)	2
110P/Hartley 3	Lamy et al. (2004)	
	Lamy & Toth (2009)	2
111P/Helin-Roman-Crockett	Lamy et al. (2004)	1
112P/Urata-Niijima	Lamy et al. (2004)	
1100/0 1: 1	Lamy & Toth (2009)	2
113P/Spitaler	Lamy et al. (2004)	1
114P/Wiseman-Skiff	Lamy et al. (2004)	2
115D/M	Lamy & Toth (2009)	2
115P/Maury 116P/Wild 4	Lamy et al. (2004)	1 1
110P/Wild 4 117P/Helin-Roman-Alu 1	Lamy et al. (2004) Lamy et al. (2004)	1
118P/Shoemaker-Levy 4	Lamy et al. (2004) Lamy et al. (2004)	1
119P/Parker-Hartley	Lamy et al. (2004) Lamy et al. (2004)	1
120P/Mueller 1	Lamy et al. (2004)	1
121P/Shoemaker-Holt 2	Lamy et al. (2004)	1
123P/West-Hartley	Lamy et al. (2004)	i
124P/Mrkos	Lamy et al. (2004)	1
125P/Spacewatch	Lamy et al. (2004)	1
126P/IRAS	Lamy et al. (2004)	1
128P/Shoemaker-Holt 1	Lamy et al. (2004)	1
129P/Shoemaker-Levy 3	Lamy et al. (2004)	1
130P/McNaught-Hughes	Lamy et al. (2004)	1
131P/Mueller 2	Lamy et al. (2004)	1
132P/Helin-Roman-Alu 2	Lamy et al. (2004)	1
134P/Koval-Vávrová	Lamy et al. (2004)	1
135P/Shoemaker-Levy 8	Lamy et al. (2004)	1
136P/Mueller 3	Lamy et al. (2004)	1
137P/Shoemaker-Levy 2	Lamy et al. (2004)	1
138P/Shoemaker-Levy 7	Lamy et al. (2004)	1
139P/Väisälä-Oterma	Lamy et al. (2004)	1

Table A.2. Continued

141P/Machholz2 Lamy et al. (2004) 143P/KM Jewitt (2002b) 144P/Kushida Lamy et al. (2004) 144P/Kushida-Muramatsu Lamy et al. (2004) 148P/Anderson-LINEAR Lamy et al. (2004) 152P/Helin-Lawrence Lamy et al. (2004) 154P/Brewington Lamy et al. (2004) 166P/NEAT Jewitt (2009b) 204P/Wild 2 Meech et al. (2004) 1172-Aneas Fornasier et al. (2007) 1173-Anchises Fornasier et al. (2007) 1647-Menelaus Fornasier et al. (2007) 1871-Astyanax Fornasier et al. (2007) 1993 RO Boehnhardt et al. (2001) 1994 ES2 Green et al. (1997) 1994 EV3 Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000)	5 1 1 1 1 1 1 1 1 1 2 1 1 2 1 1 3 2 1
143P/KM	5 1 1 1 1 1 1 1 2 1 1 2 1 1 1 1 3 3 3
Jewitt et al. (2003) Lamy et al. (2004) 144P/Kushida Lamy et al. (2004) 147P/Kushida-Muramatsu Lamy et al. (2004) 148P/Anderson-LINEAR Lamy et al. (2004) 152P/Helin-Lawrence Lamy et al. (2004) 154P/Brewington Lamy et al. (2004) 166P/NEAT Jewitt (2009b) 204P/Wild 2 Meech et al. (2004) 1172-Aneas Fornasier et al. (2007) 1173-Anchises Fornasier et al. (2007) 1647-Menelaus Fornasier et al. (2007) 1871-Astyanax Fornasier et al. (2007) 1893 RO Boehnhardt et al. (2001) Tegler & Romanishin (2000) 1994 ES ₂ Green et al. (1997) 1994 EV ₃ Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	1 1 1 1 1 2 1 2 1 2 1 3 2 1
Lamy et al. (2004) 144P/Kushida Lamy et al. (2004) 147P/Kushida-Muramatsu Lamy et al. (2004) 148P/Anderson-LINEAR Lamy et al. (2004) 152P/Helin-Lawrence Lamy et al. (2004) 154P/Brewington Lamy et al. (2004) 166P/NEAT Jewitt (2009b) 204P/Wild 2 Meech et al. (2004) 1172-Aneas Fornasier et al. (2007) 1173-Anchises Fornasier et al. (2007) 1647-Menelaus Fornasier et al. (2007) 1871-Astyanax Fornasier et al. (2007) 1993 RO Boehnhardt et al. (2001) Tegler & Romanishin (2000) 1994 ES ₂ Green et al. (1997) 1994 EV ₃ Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	1 1 1 1 1 2 1 2 1 2 1 3 2 1
144P/Kushida Lamy et al. (2004) 147P/Kushida-Muramatsu Lamy et al. (2004) 148P/Anderson-LINEAR Lamy et al. (2004) 152P/Helin-Lawrence Lamy et al. (2004) 154P/Brewington Lamy et al. (2004) 166P/NEAT Jewitt (2009b) 204P/Wild 2 Meech et al. (2004) 1172-Aneas Fornasier et al. (2007) 1647-Menelaus Fornasier et al. (2007) 1871-Astyanax Fornasier et al. (2007) 1993 RO Boehnhardt et al. (2001) 1994 ES2 Green et al. (1997) 1994 EV3 Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	1 1 1 1 1 2 1 2 1 2 1 3 2 1
147P/Kushida-Muramatsu Lamy et al. (2004) 148P/Anderson-LINEAR Lamy et al. (2004) 152P/Helin-Lawrence Lamy et al. (2004) 154P/Brewington Lamy et al. (2004) 166P/NEAT Jewitt (2009b) 204P/Wild 2 Meech et al. (2004) 1172-Aneas Fornasier et al. (2007) 1173-Anchises Fornasier et al. (2007) 1647-Menelaus Fornasier et al. (2007) 1871-Astyanax Fornasier et al. (2007) 1993 RO Boehnhardt et al. (2001) 1994 ES2 Green et al. (1997) 1994 EV3 Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	1 1 1 1 2 1 2 1 2 1 1 3
148P/Anderson-LINEAR Lamy et al. (2004) 152P/Helin-Lawrence Lamy et al. (2004) 154P/Brewington Lamy et al. (2004) 166P/NEAT Jewitt (2009b) 204P/Wild 2 Meech et al. (2004) 1172-Aneas Fornasier et al. (2007) 1173-Anchises Fornasier et al. (2007) 1647-Menelaus Fornasier et al. (2007) 1871-Astyanax Fornasier et al. (2007) 1993 RO Boehnhardt et al. (2001) Tegler & Romanishin (2000) Green et al. (1997) 1994 EV3 Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	1 1 2 1 2 1 1 1 3
154P/Brewington 166P/NEAT Jewitt (2009b) 204P/Wild 2 Meech et al. (2004) 1172-Aneas Fornasier et al. (2007) 1173-Anchises Fornasier et al. (2007) 1647-Menelaus Fornasier et al. (2007) 1871-Astyanax Fornasier et al. (2007) 1993 RO Boehnhardt et al. (2001) Tegler & Romanishin (2000) 1994 ES ₂ Green et al. (1997) 1994 EV ₃ Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	1 2 1 2 1 1 1 3
166P/NEAT Jewitt (2009b) 204P/Wild 2 Meech et al. (2004) 1172-Aneas Fornasier et al. (2007) 1173-Anchises Fornasier et al. (2007) 1647-Menelaus Fornasier et al. (2007) 1871-Astyanax Fornasier et al. (2007) 1993 RO Boehnhardt et al. (2001) Tegler & Romanishin (2000) Tegler & Romanishin (2000) 1994 ES2 Green et al. (1997) 1994 EV3 Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	2 1 2 1 1 3 2 1
204P/Wild 2 1172-Aneas Fornasier et al. (2007) 1173-Anchises Fornasier et al. (2007) 1647-Menelaus Fornasier et al. (2007) 1871-Astyanax Fornasier et al. (2007) 1993 RO Boehnhardt et al. (2001) Tegler & Romanishin (2000) 1994 ES ₂ Green et al. (1997) 1994 EV ₃ Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	1 2 1 1 3 2 1
1172-Aneas Fornasier et al. (2007) 1173-Anchises Fornasier et al. (2007) 1647-Menelaus Fornasier et al. (2007) 1871-Astyanax Fornasier et al. (2007) 1993 RO Boehnhardt et al. (2001) Tegler & Romanishin (2000) 1994 ES ₂ Green et al. (1997) 1994 EV ₃ Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	2 1 1 3 2 1
1173-Anchises 1647-Menelaus 1647-Menelaus 1871-Astyanax 1993 RO 1994 ES ₂ 1994 EV ₃ 1994 TA Fornasier et al. (2001) Green et al. (2001) Gril-Hutton & Licandro (2001) Boehnhardt et al. (2002) Tegler & Romanishin (2000) Green et al. (2001) Gril-Hutton & Licandro (2001) Boehnhardt et al. (2002) Tegler & Romanishin (2000) Jewitt & Luu (2001)	1 1 3 2 1
1647-Menelaus Fornasier et al. (2007) 1871-Astyanax Fornasier et al. (2007) 1993 RO Boehnhardt et al. (2001) Tegler & Romanishin (2000) 1994 ES ₂ Green et al. (1997) 1994 EV ₃ Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	1 3 2 1
1871-Astyanax Fornasier et al. (2007) 1993 RO Boehnhardt et al. (2001) Tegler & Romanishin (2000) 1994 ES ₂ Green et al. (1997) 1994 EV ₃ Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	3 2 1
1993 RO Boehnhardt et al. (2001) Tegler & Romanishin (2000) 1994 ES ₂ Green et al. (1997) 1994 EV ₃ Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	2 1
Tegler & Romanishin (2000) 1994 ES ₂ Green et al. (1997) 1994 EV ₃ Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	1
1994 EV ₃ Boehnhardt et al. (2001) Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	
Gil-Hutton & Licandro (2001) Boehnhardt et al. (2002) Tegler & Romanishin (2000) Jewitt & Luu (2001)	4
Boehnhardt et al. (2002) 1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	4
1994 TA Tegler & Romanishin (2000) Jewitt & Luu (2001)	4
Jewitt & Luu (2001)	
	2
	2
	2
	5
	4
1995 HM ₅ Romanishin & Tegler (1999)	
Gil-Hutton & Licandro (2001)	
Boehnhardt et al. (2001)	
Benecchi et al. (2011)	
Barucci et al. (2000)	0
6 ,	8 2
<u> </u>	1
1996 RQ ₂₀ Romanishin & Tegler (1999)	•
Jewitt & Luu (2001)	
Boehnhardt et al. (2001)	
Delsanti et al. (2001)	
Benecchi et al. (2011)	_
8 ()	7
1996 RR ₂₀ Jewitt & Luu (2001) Tegler & Romanishin (2000)	
Boehnhardt et al. (2002)	
	4
	1
1996 TK ₆₆ Tegler & Romanishin (2000)	
Jewitt & Luu (2001)	
Doressoundiram et al. (2002)	
	4
1996 TS ₆₆ Jewitt & Luu (1998)	
Romanishin & Tegler (1999)	
Davies et al. (2000) Jewitt & Luu (2001)	
	7
1997 CT ₂₉ Tegler & Romanishin (2000)	•
Benecchi et al. (2011)	
Barucci et al. (2000)	4
1997 CV ₂₉ Tegler & Romanishin (2003)	
	2
->>	1
1997 QH ₄ Tegler & Romanishin (2000)	
Jewitt & Luu (2001) Delsanti et al. (2001)	
	4
	1

Table A.2. Continued

Object	References	Nr.
1997 RT ₅	Gladman et al. (1998)	
	Boehnhardt et al. (2002)	4
1997 RX ₉	Benecchi et al. (2011) Gladman et al. (1998)	4 1
1997 SZ ₁₀	Tegler & Romanishin (2000)	1
1998 FS ₁₄₄	Tegler & Romanishin (2000) Tegler & Romanishin (2003)	1
1990 15 144	Benecchi et al. (2011)	
	Barucci et al. (2000)	4
1998 KG ₆₂	Gil-Hutton & Licandro (2001)	
	Boehnhardt et al. (2002)	
	Benecchi et al. (2011)	
	Doressoundiram et al. (2001)	5
1998 KS ₆₅	Tegler & Romanishin (2003)	_
1000 1717	Benecchi et al. (2011)	2
1998 KY ₆₁	Benecchi et al. (2011)	1
1998 UR ₄₃	Gil-Hutton & Licandro (2001)	
	Delsanti et al. (2001) Benecchi et al. (2011)	4
1998 UU ₄₃	Benecchi et al. (2011) Benecchi et al. (2011)	2
1998 WS ₃₁	Benecchi et al. (2011)	2
1550 (1531	Peixinho et al. (2004)	2
1998 WU ₂₄	Davies et al. (2001)	1
1998 WV ₂₄	Tegler & Romanishin (2000)	•
24	Benecchi et al. (2011)	2
1998 WV ₃₁	Delsanti et al. (2001)	
	Peixinho et al. (2004)	2
1998 WW ₃₁	Benecchi et al. (2009)	1
1998 WW_{31+B}	Benecchi et al. (2009)	1
1998 WX ₂₄	Tegler & Romanishin (2000)	_
1000 WW	Benecchi et al. (2011)	2
1998 WX ₃₁	Delsanti et al. (2001)	2
1000 WW	Benecchi et al. (2011)	3 2
1998 WY ₂₄ 1998 WZ ₃₁	Benecchi et al. (2011) Benecchi et al. (2011)	2
1996 W Z ₃₁	Peixinho et al. (2004)	3
1998 XY ₉₅	Boehnhardt et al. (2001)	3
1770 71195	Benecchi et al. (2011)	2
1999 CB ₁₁₉	Peixinho et al. (2004)	1
1999 CD ₁₅₈	Delsanti et al. (2004)	
	Doressoundiram et al. (2002)	
	Delsanti et al. (2006)	
	Benecchi et al. (2011)	5
1999 CF ₁₁₉	Delsanti et al. (2001)	
	Benecchi et al. (2011)	3
1999 CH ₁₁₉	Benecchi et al. (2011)	1
1999 CJ ₁₁₉	Benecchi et al. (2011)	1
1999 CL ₁₁₉	Peixinho et al. (2004)	2
1999 CQ ₁₃₃	Benecchi et al. (2011) Benecchi et al. (2011)	1
1999 CX ₁₃₁	Benecchi et al. (2011) Benecchi et al. (2011)	1
1777 CA ₁₃₁	Peixinho et al. (2004)	3
1999 HJ ₁₂	Benecchi et al. (2011)	2
1999 HS ₁₁	Tegler & Romanishin (2003)	_
	Peixinho et al. (2004)	
	Doressoundiram et al. (2001)	3
1999 HV ₁₁	Tegler & Romanishin (2003)	1
1999 OD ₄	Benecchi et al. (2011)	1
1999 OE ₄	Peixinho et al. (2004)	
1000 011	Benecchi et al. (2011)	4
1999 OH ₄	Benecchi et al. (2011)	_
1000 01	Snodgrass et al. (2010)	2
1999 OJ ₄	Peixinho et al. (2004)	
	Benecchi et al. (2011)	<i>E</i>
1000 ()	Benecchi et al. (2009)	5
1999 OJ _{4+B} 1999 OM ₄	Benecchi et al. (2009) Boehnhardt et al. (2002)	1 1
	DOMERNIANU ELAL UZUUZI	

Table A.2. Continued

Object	References	Nr.
1999 RT ₂₁₄	Benecchi et al. (2009)	1
1999 RX ₂₁₄	Peixinho et al. (2004)	
1000 PW	Benecchi et al. (2011)	3
1999 RY ₂₁₄	Peixinho et al. (2004)	2
1999 TR ₁₁	Tegler & Romanishin (2000) Benecchi et al. (2011)	2
1999 XY ₁₄₃	Benecchi et al. (2011) Benecchi et al. (2011)	1
2000 AF ₂₅₅	Benecchi et al. (2011) Benecchi et al. (2011)	1
2000 CE ₁₀₅	Benecchi et al. (2011)	2
2000 CF ₁₀₅	Tegler et al. (2003)	2
2000 CI 103	Benecchi et al. (2011)	
	Benecchi et al. (2009)	6
2000 CF_{105+B}	Benecchi et al. (2009)	1
2000 CG ₁₀₅	Benecchi et al. (2011)	
	Snodgrass et al. (2010)	3
2000 CK ₁₀₅	Benecchi et al. (2011)	2
2000 CL ₁₀₄	Boehnhardt et al. (2002)	
	Benecchi et al. (2011)	
	Peixinho et al. (2004)	4
2000 CN ₁₀₅	Peixinho et al. (2004)	
	Jewitt et al. (2007)	4
2000 CO ₁₀₅	Benecchi et al. (2011)	2
2000 CP ₁₀₄	Benecchi et al. (2011)	2
2000 CQ ₁₀₅	Benecchi et al. (2011)	
	Peixinho et al. (2004)	
	Tegler et al. (2003)	_
2000 CO	Jewitt et al. (2007)	5
2000 CQ ₁₁₄	Benecchi et al. (2011)	2
2000 CQ _{114+B}	Benecchi et al. (2009)	3 1
2000 CQ _{114+B} 2000 FS ₅₃	Benecchi et al. (2009) Tegler & Romanishin (2003)	1
2000 1 353	Benecchi et al. (2011)	2
2000 FV ₅₃	Peixinho et al. (2011)	1
2000 FZ ₅₃	Peixinho et al. (2004)	2
2000 GV ₁₄₆	Benecchi et al. (2011)	1
2000 KK ₄	McBride et al. (2003)	-
•	Tegler & Romanishin (2003)	
	Benecchi et al. (2011)	3
2000 KL ₄	Benecchi et al. (2011)	1
$2000 \; \mathrm{OU}_{69}$	Peixinho et al. (2004)	
	Benecchi et al. (2011)	3
2000 PD ₃₀	Benecchi et al. (2011)	2
2000 PE ₃₀	Delsanti et al. (2006)	
	Doressoundiram et al. (2007)	
	Benecchi et al. (2011)	
	Sheppard (2010)	_
2000 PM	Doressoundiram et al. (2001)	7
2000 PH ₃₀	Benecchi et al. (2011)	1
2000 QL ₂₅₁	Benecchi et al. (2009)	1
2000 QL _{251+B}	Benecchi et al. (2009)	1
2001 FM	Benecchi et al. (2009)	1 1
2001 FM ₁₉₄ 2001 FU ₁₇₂	Tegler et al. (2003)	1
2001 FU ₁₇₂ 2001 KA ₇₇	Snodgrass et al. (2010) Doressoundiram et al. (2002)	1
2001 KA77	Peixinho et al. (2004)	
	Doressoundiram et al. (2005b)	4
2001 KB ₇₇	Peixinho et al. (2004)	6
2001 KD ₇₇ 2001 KD ₇₇	Doressoundiram et al. (2002)	J
	Peixinho et al. (2004)	
	Doressoundiram et al. (2007)	
	Benecchi et al. (2011)	9
2001 KG ₇₇	Tegler et al. (2003)	ĺ
2001 KY ₇₆	Benecchi et al. (2011)	1
2001 OG ₁₀₉	Benecchi et al. (2011)	1
2001 OK ₁₀₈	Benecchi et al. (2011)	1
2001 QC ₂₉₈	Benecchi et al. (2011)	
	Jewitt et al. (2007)	
	Benecchi et al. (2009)	3

Table A.2. Continued

Object	References	Nr.
2001 QD ₂₉₈	Doressoundiram et al. (2005b)	1
2001 QF ₂₉₈	Doressoundiram et al. (2007)	
	Fornasier et al. (2004)	
	Delsanti et al. (2006)	0
2001 OV	Doressoundiram et al. (2005b)	8
2001 QX_{322}	Tegler et al. (2003) Benecchi et al. (2011)	
	Jewitt et al. (2007)	3
2001 XR ₂₅₄	Benecchi et al. (2009)	1
2001 XR _{254+B}	Benecchi et al. (2009)	1
2001 XU ₂₅₄	Benecchi et al. (2011)	1
2001 XZ ₂₅₅	Tegler et al. (2003)	1
2002 CB ₂₄₉	Peixinho et al. (2004)	1
2002 DH ₅	Peixinho et al. (2004)	1
2002 GB ₃₂	Sheppard (2010)	1
2002 PP ₁₄₉	Jewitt et al. (2007)	1
2003 FZ ₁₂₉	Sheppard (2010)	1
2003 GH ₅₅	Jewitt et al. (2007)	1
2003 HB ₅₇	Sheppard (2010)	1
2003 HX ₅₆	Snodgrass et al. (2010)	1
2003 QA ₉₂	Romanishin et al. (2010)	1
2003 QD ₁₁₂ 2003 QK ₉₁	Jewitt (2009b) Sheppard (2010)	1 1
2003 QK ₉₁ 2003 QQ ₉₁	Romanishin et al. (2010)	1
2003 QW ₁₁₁	Benecchi et al. (2009)	1
2003 QW _{111+B}	Benecchi et al. (2009)	1
2003 QW ₉₀	DeMeo et al. (2009)	2
2003 QY ₉₀	Benecchi et al. (2009)	1
2003 QY_{90+B}	Benecchi et al. (2009)	1
2003 SQ ₃₁₇	Snodgrass et al. (2010)	1
2003 TH ₅₈	Snodgrass et al. (2010)	1
2003 TJ ₅₈	Benecchi et al. (2009)	1
2003 TJ _{58+B}	Benecchi et al. (2009)	1
2003 UY ₂₉₁	Sheppard (2010)	1
2003 UZ ₁₁₇	Jewitt et al. (2007)	
2002 3/1	DeMeo et al. (2009)	4
2003 YL ₁₇₉	Romanishin et al. (2010)	1
2004 OJ ₁₄ 2004 PB ₁₀₈	Sheppard (2010)	1 1
2004 PB _{108+B}	Benecchi et al. (2009) Benecchi et al. (2009)	1
2004 PT _{108+B} 2004 PT ₁₀₇	Snodgrass et al. (2010)	1
2004 PY ₄₂	Jewitt (2009b)	1
2004 VN ₁₁₂	Sheppard (2010)	1
2004 XR ₁₉₀	Sheppard (2010)	1
2005 CB ₇₉	Snodgrass et al. (2010)	1
2005 EO ₂₉₇	Sheppard (2010)	1
2005 EO ₃₀₄	Benecchi et al. (2009)	1
2005 EO _{304+B}	Benecchi et al. (2009)	1
2005 GE ₁₈₇	Snodgrass et al. (2010)	1
2005 PU ₂₁	Sheppard (2010)	1
2005 SD_{278}	Sheppard (2010)	1
2006 SQ ₃₇₂	Sheppard (2010)	1
2007 TG ₄₂₂	Sheppard (2010)	1
2007 VJ ₃₀₅	Sheppard (2010)	1
2008 KV ₄₂	Sheppard (2010)	1
2008 OG ₁₉ 2008 YB ₃	Sheppard (2010)	1 1
2008 YB ₃ 2060-Chiron	Sheppard (2010) Hartmann et al. (1981)	1
2000-Cinion	Davies et al. (1998)	
	Parker et al. (1997)	
	Jewitt (2002b)	
	Bauer et al. (2003)	
	Doressoundiram et al. (2007)	
	Green et al. (1997)	34
2223-Sarpedon	Fornasier et al. (2007)	1
2357-Phereclos	Fornasier et al. (2007)	1
3548-Eurybates	Fornasier et al. (2007)	1
	Fornasier et al. (2007)	2
4035-1986 WD 4829-Sergestus	Fornasier et al. (2007)	2

Table A.2. Continued

Object	References	Nr.
5130-Ilioneus 5145-Pholus	Fornasier et al. (2007)	1
5145-Photus	Binzel (1992) Mueller et al. (1992)	
	Davies et al. (1993)	
	Davies et al. (1998)	
	Romanishin & Tegler (1999)	
	Weintraub et al. (1997)	
	Bauer et al. (2003)	
	Doressoundiram et al. (2007)	
	Buie & Bus (1992) Davies (2000)	
	Fink et al. (1992)	
	Green et al. (1997)	
	Tegler & Romanishin (1998)	41
5244-Amphilochos	Fornasier et al. (2007)	1
5258-1989 AU ₁	Fornasier et al. (2007)	1
5511-Cloanthus	Fornasier et al. (2007)	1
6545-1986 TR ₆	Fornasier et al. (2007)	1
6998-Tithonus	Fornasier et al. (2007)	1
7066-Nessus	Davies et al. (1998)	
	Romanishin & Tegler (1999) Bauer et al. (2003)	
	Davies (2000)	
	Tegler & Romanishin (1998)	17
7352-1994 CO	Fornasier et al. (2007)	1
8405-Asbolus	Weintraub et al. (1997)	_
	Brown & Luu (1997)	
	Davies et al. (1998)	
	Bauer et al. (2003)	
	Romon-Martin et al. (2002)	
	Davies (2000)	
	Rabinowitz et al. (2007)	43
9030-1989 UX ₅	Tegler & Romanishin (1998) Fornasier et al. (2007)	1
9430-Erichthonio	Fornasier et al. (2007)	1
9818-Eurymachos	Fornasier et al. (2007)	1
10199-Chariklo	Davies et al. (1998)	-
	Jewitt & Kalas (1998)	
	Romanishin & Tegler (1999)	
	N. McBride & Foster (1999)	
	Jewitt & Luu (2001)	
	Peixinho et al. (2001)	
	Bauer et al. (2003)	
	DeMeo et al. (2009) Davies (2000)	
	Tegler & Romanishin (1998)	64
10370-Hylonome	Davies et al. (1998)	0-1
100,011,101101110	Romanishin & Tegler (1999)	
	Bauer et al. (2003)	
	Doressoundiram et al. (2002)	
	Delsanti et al. (2006)	
	Davies (2000)	
11000 1004 66	Tegler & Romanishin (1998)	25
11089-1994 CS ₈	Fornasier et al. (2007)	1
11351-1997 TS ₂₅ 11488-1988 RM ₁₁	Fornasier et al. (2007) Fornasier et al. (2007)	$\frac{1}{2}$
11488-1988 KM ₁₁ 11663-1997 GO ₂₄	Fornasier et al. (2007)	1
12917-1998 TG ₁₆	Fornasier et al. (2007)	1
	Dotto et al. (2006)	2
12921-1998 WZ ₅	Fornasier et al. (2007)	$\frac{1}{2}$
13463-Antiphos	Fornasier et al. (2007)	
	Dotto et al. (2006)	2
	Fornasier et al. (2007)	1
13862-1999 XT ₁₆₀	E : (1/2007)	2
14707-2000 CC ₂₀	Fornasier et al. (2007)	2
	Fornasier et al. (2007)	
14707-2000 CC ₂₀ 15094-1999 WB ₂	Fornasier et al. (2007) Dotto et al. (2006)	3
14707-2000 CC ₂₀	Fornasier et al. (2007)	

Table A.2. Continued

Object	References	Nr.
15760-1992 QB ₁	Jewitt & Luu (2001)	
	Tegler & Romanishin (2000)	
	Boehnhardt et al. (2001)	
	Benecchi et al. (2011) Romanishin et al. (1997)	5
15788-1993 SB	Gil-Hutton & Licandro (2001)	3
13700 1773 BB	Jewitt & Luu (2001)	
	Tegler & Romanishin (2000)	
	McBride et al. (2003)	
	Delsanti et al. (2001)	
	Benecchi et al. (2011)	6
15789-1993 SC	Davies et al. (1997)	
	Romanishin & Tegler (1999)	
	Jewitt & Luu (1998)	
	Davies et al. (2000)	
	Jewitt & Luu (2001) Benecchi et al. (2011)	
	Luu & Jewitt (1996)	
	Romanishin et al. (1997)	
	Tegler & Romanishin (1998)	17
15807-1994 GV ₉	Gil-Hutton & Licandro (2001)	1
15809-1994 JS	Benecchi et al. (2011)	2
15810-1994 JR ₁	Green et al. (1997)	
	Romanishin & Tegler (1999)	
	Benecchi et al. (2011)	
	Barucci et al. (1999)	_
15000 1004 FD	Tegler & Romanishin (1998)	9
15820-1994 TB	Davies et al. (2000)	
	Jewitt & Luu (2001) Delsanti et al. (2001)	
	Delsanti et al. (2001) Delsanti et al. (2004)	
	Delsanti et al. (2004) Delsanti et al. (2006)	
	Doressoundiram et al. (2007)	
	Barucci et al. (1999)	
	Davies (2000)	
	Romanishin et al. (1997)	
	Tegler & Romanishin (1998)	15
15836-1995 DA ₂	Green et al. (1997)	
15054 1006 FN	Benecchi et al. (2011)	6
15874-1996 TL ₆₆	Jewitt et al. (2007)	
	Jewitt & Luu (1998)	
	Romanishin & Tegler (1999) Boehnhardt et al. (2001)	
	Davies et al. (2000)	
	Jewitt & Luu (2001)	
	Benecchi et al. (2011)	
	Doressoundiram et al. (2007)	
	Barucci et al. (1999)	
	Davies (2000)	
	Tegler & Romanishin (1998)	16
15875-1996 TP ₆₆	Jewitt & Luu (1998)	
	Romanishin & Tegler (1999)	
	Boehnhardt et al. (2001)	
	Davies et al. (2000) Jewitt & Luu (2001)	
	Barucci et al. (1999)	7
15883-1997 CR ₂₉	Jewitt & Luu (2001)	,
	Doressoundiram et al. (2001)	
	Benecchi et al. (2011)	3
15977-1998 MA ₁₁	Fornasier et al. (2007)	1
16684-1994 JQ ₁	Green et al. (1997)	
	Gil-Hutton & Licandro (2001)	
	Boehnhardt et al. (2002)	
	Tegler & Romanishin (2003)	~
17/1/C 1000 P.P.	Benecchi et al. (2011)	8
17416-1988 RR ₁₀	Fornasier et al. (2007)	1 1
18060-1999 XJ ₁₅₆	Fornasier et al. (2007)	1

Table A.2. Continued

Object	References	Nr.
18137-2000 OU ₃₀	Fornasier et al. (2007)	1
18268-Dardanos	Fornasier et al. (2007)	1
18493-Demoleon	Fornasier et al. (2007)	2
18940-2000 QV ₄₉	Fornasier et al. (2007)	1
19255-1994 VK ₈	Tegler & Romanishin (2000)	
	Doressoundiram et al. (2001)	4
10200 1007 57	Benecchi et al. (2011)	4
19299-1996 SZ ₄	Jewitt & Luu (2001)	
	Tegler & Romanishin (2000)	
	McBride et al. (2003) Boehnhardt et al. (2002)	5
19308-1996 TO ₆₆	Jewitt et al. (2007)	3
19308-1990 1066	Jewitt & Luu (1998)	
	Romanishin & Tegler (1999)	
	Hainaut et al. (2000)	
	Gil-Hutton & Licandro (2001)	
	Davies et al. (2000)	
	Jewitt & Luu (2001)	
	Boehnhardt et al. (2001)	
	Sheppard (2010)	
	Barucci et al. (1999)	
	Davies (2000)	
	Tegler & Romanishin (1998)	16
19521-Chaos	Davies et al. (2000)	
	Tegler & Romanishin (2000)	
	Boehnhardt et al. (2001)	
	Delsanti et al. (2004)	
	Doressoundiram et al. (2002)	
	Benecchi et al. (2011)	
	Doressoundiram et al. (2007)	
	Barucci et al. (2000)	13
20000-Varuna	Doressoundiram et al. (2002)	
	Hainaut (2001)	
	Jewitt & Sheppard (2002)	
	McBride et al. (2003)	
	Doressoundiram et al. (2007)	
	Jewitt et al. (2007)	
*****	Rabinowitz et al. (2007)	8
20108-1995 QZ ₉	Gil-Hutton & Licandro (2001)	
	Tegler & Romanishin (2000)	
20161 1006 FD	Benecchi et al. (2011)	3
20161-1996 TR ₆₆	Benecchi et al. (2011)	1
20738-1999 XG ₁₉₁	Fornasier et al. (2007)	2
22540 E : 1	Dotto et al. (2006)	2
23549-Epicles	Fornasier et al. (2007)	1
23694-1997 KZ ₃	Formasier et al. (2007)	1
24233-1999 XD ₉₄	Formasier et al. (2007)	1 1
24341-2000 AJ ₈₇ 24380-2000 AA ₁₆₀	Fornasier et al. (2007)	1
24390-2000 AA ₁₆₀ 24390-2000 AD ₁₇₇	Fornasier et al. (2007) Fornasier et al. (2007)	1
24390-2000 AD ₁₇₇	Dotto et al. (2007)	2
24420-2000 BU ₂₂	Fornasier et al. (2007)	1
24426-2000 CR ₁₂	Fornasier et al. (2007) Fornasier et al. (2007)	1
24444-2000 OP ₃₂	Fornasier et al. (2007)	1
24452-2000 QU ₁₆₇	Fornasier et al. (2007) Fornasier et al. (2007)	1
24467-2000 SS ₁₆₅	Fornasier et al. (2007)	1
24835-1995 SM ₅₅	Gil-Hutton & Licandro (2001)	1
2-1033-1773 BIVISS	Boehnhardt et al. (2001)	
	Delsanti et al. (2001)	
	McBride et al. (2001)	
	Delsanti et al. (2004)	
	Doressoundiram et al. (2002)	
	Benecchi et al 170111	
	Benecchi et al. (2011) Jewitt et al. (2007)	
	Jewitt et al. (2007) Doressoundiram et al. (2007)	

Table A.2. Continued

Object	References	Nr.
24952-1997 QJ ₄	Gil-Hutton & Licandro (2001)	
	Jewitt & Luu (2001)	
	Delsanti et al. (2001) Boehnhardt et al. (2002)	4
24978-1998 HJ ₁₅₁	Tegler & Romanishin (2003)	7
24576 1556 113151	Benecchi et al. (2011)	2
25347-1999 RQ ₁₁₆	Fornasier et al. (2007)	1
26181-1996 GQ ₂₁	McBride et al. (2003)	
	Boehnhardt et al. (2002)	
	Delsanti et al. (2006)	
	Doressoundiram et al. (2003)	_
2/200 1000 914	Doressoundiram et al. (2007)	8
26308-1998 SM ₁₆₅	Tegler & Romanishin (2000)	
	McBride et al. (2003)	
	Delsanti et al. (2004) Doressoundiram et al. (2007)	
	Jewitt et al. (2007)	
	Benecchi et al. (2009)	8
26308-1998 SM _{165+B}	Benecchi et al. (2009)	1
26375-1999 DE ₉	Jewitt & Luu (2001)	•
	Delsanti et al. (2001)	
	Doressoundiram et al. (2002)	
	McBride et al. (2003)	
	Delsanti et al. (2006)	
	Doressoundiram et al. (2003)	
	Tegler et al. (2003)	
	Benecchi et al. (2011)	
	Doressoundiram et al. (2007)	
	DeMeo et al. (2009)	12
28958-2001 CQ ₄₂	Rabinowitz et al. (2007) Fornasier et al. (2007)	13 1
28978-Ixion	Doressoundiram et al. (2002)	1
26) / 6-1/1011	Doressoundiram et al. (2002) Doressoundiram et al. (2007)	
	DeMeo et al. (2009)	
	Rabinowitz et al. (2007)	8
29981-1999 TD ₁₀	Delsanti et al. (2001)	
	McBride et al. (2003)	
	Doressoundiram et al. (2002)	
	Doressoundiram et al. (2007)	
	Consolmagno et al. (2000)	
20.000 H; I	Rabinowitz et al. (2007)	18
30698-Hippokoon	Fornasier et al. (2007)	3
31820-1999 RT ₁₈₆ 31821-1999 RK ₂₂₅	Fornasier et al. (2007)	1 1
31824-Elatus	Fornasier et al. (2007) Peixinho et al. (2001)	1
31024-Liatus	Delsanti et al. (2004)	
	Doressoundiram et al. (2002)	
	Doressoundiram et al. (2007)	
	Gutiérrez et al. (2001)	14
32430-2000 RQ ₈₃	Fornasier et al. (2007)	1
32532-Thereus	Farnham & Davies (2003)	
	Tegler et al. (2003)	
	Barucci et al. (2002)	
	Bauer et al. (2003)	
	Doressoundiram et al. (2007)	
	DeMeo et al. (2009)	221
32615 2001 OU	Rabinowitz et al. (2007)	221 1
32615-2001 QU ₂₇₇ 32794-1989 UE ₅	Fornasier et al. (2007) Fornasier et al. (2007)	1
32929-1995 QY ₉	Gil-Hutton & Licandro (2001)	1
52727 1773 Q 19	Davies et al. (2000)	
	Barucci et al. (1999)	5
33001-1997 CU ₂₉	Tegler & Romanishin (2000)	5
27	Jewitt & Luu (2001)	
	Doressoundiram et al. (2001)	
	Benecchi et al. (2011)	
	Barucci et al. (2000)	

Table A.2. Continued

Object	References	Nr.
33128-1998 BU ₄₈	Delsanti et al. (2004)	
	Doressoundiram et al. (2002)	
	Delsanti et al. (2006)	
	Bauer et al. (2003)	9
33340-1998 VG ₄₄	Boehnhardt et al. (2001)	
	Doressoundiram et al. (2001)	
	McBride et al. (2003)	
	Doressoundiram et al. (2002)	
	Delsanti et al. (2006) Benecchi et al. (2011)	
	Doressoundiram et al. (2007)	12
34785-2001 RG ₈₇	Fornasier et al. (2007)	12
35671-1998 SN ₁₆₅	Jewitt & Luu (2001)	1
33071-1776 SIN ₁₆₅	Gil-Hutton & Licandro (2001)	
	Delsanti et al. (2001)	
	McBride et al. (2003)	
	Fornasier et al. (2004)	
	Doressoundiram et al. (2001)	6
38083-Rhadamanth	Boehnhardt et al. (2002)	1
38084-1999 HB ₁₂	Benecchi et al. (2011)	
12	Peixinho et al. (2004)	
	Doressoundiram et al. (2001)	3
38628-Huya	Ferrin et al. (2001)	
•	Jewitt & Luu (2001)	
	Schaefer & Rabinowitz (2002)	
	Boehnhardt et al. (2002)	
	McBride et al. (2003)	
	Doressoundiram et al. (2007)	
	Doressoundiram et al. (2001)	
	Rabinowitz et al. (2007)	58
39285-2001 BP ₇₅	Fornasier et al. (2007)	1
40314-1999 KR ₁₆	Jewitt & Luu (2001)	
	Delsanti et al. (2006)	
	Boehnhardt et al. (2002)	_
42201 2001 LID	Jewitt et al. (2007)	5
42301-2001 UR ₁₆₃	Doressoundiram et al. (2007) Doressoundiram et al. (2005b)	
	Benecchi et al. (2003b)	6
42355-Typhon	Peixinho et al. (2004)	U
42333-Typhon	Tegler et al. (2004)	
	Doressoundiram et al. (2007)	
	Jewitt et al. (2007)	
	DeMeo et al. (2009)	
	Benecchi et al. (2009)	
	Rabinowitz et al. (2007)	12
42355-Typhon+B	Benecchi et al. (2009)	1
43212-2000 AL ₁₁₃	Fornasier et al. (2007)	1
44594-1999 OX ₃	Tegler & Romanishin (2000)	
	Bauer et al. (2003)	
	Delsanti et al. (2001)	
	McBride et al. (2003)	
	Boehnhardt et al. (2002)	
	Doressoundiram et al. (2002)	
	Peixinho et al. (2004)	
	Doressoundiram et al. (2005b)	
	Doressoundiram et al. (2007)	
	Delsanti et al. (2006)	
	Jewitt et al. (2007)	
	Sheppard (2010) Doressoundiram et al. (2001)	30
45802-2000 PV	Benecchi et al. (2011)	2
45802-2000 PV ₂₉	Deficeelli et al. (2011)	

Table A.2. Continued

Object	References	Nr
47171-1999 TC ₃₆	Boehnhardt et al. (2001)	
	Delsanti et al. (2001)	
	McBride et al. (2003)	
	Delsanti et al. (2004)	
	Dotto et al. (2003)	
	Tegler et al. (2003)	
	Doressoundiram et al. (2007)	
	Protopapa et al. (2009)	
	DeMeo et al. (2009)	
	Benecchi et al. (2009)	
	Doressoundiram et al. (2001)	
	Rabinowitz et al. (2007)	23
47171-1999 TC _{36+B}	Benecchi et al. (2009)	1
47932-2000 GN ₁₇₁	McBride et al. (2003)	
	Boehnhardt et al. (2002)	
	Doressoundiram et al. (2007)	
	DeMeo et al. (2009)	
	Rabinowitz et al. (2007)	13
47967-2000 SL ₂₉₈	Fornasier et al. (2007)	1
48249-2001 SY ₃₄₅	Fornasier et al. (2007)	1
48252-2001 TL ₂₁₂	Fornasier et al. (2007)	1
48639-1995 TL ₈	Delsanti et al. (2004)	
	Doressoundiram et al. (2002)	
	Benecchi et al. (2011)	
	Sheppard (2010)	4
49036-Pelion	Tegler & Romanishin (2000)	
	Boehnhardt et al. (2002)	
	Doressoundiram et al. (2002)	3
50000-Quaoar	Fornasier et al. (2004)	
	Tegler et al. (2003)	
	DeMeo et al. (2009)	
	Rabinowitz et al. (2007)	4
51359-2000 SC ₁₇	Fornasier et al. (2007)	1
52747-1998 HM ₁₅₁	Tegler & Romanishin (2003)	1
52872-Okyrhoe	Bauer et al. (2003)	
•	Delsanti et al. (2001)	
	Dotto et al. (2003)	
	Delsanti et al. (2006)	
	Doressoundiram et al. (2007)	
	Doressoundiram et al. (2001)	18
52975-Cyllarus	Boehnhardt et al. (2001)	
,	Delsanti et al. (2004)	
	Bauer et al. (2003)	
	Doressoundiram et al. (2002)	
	Tegler et al. (2003)	9
53469-2000 AX ₈	Fornasier et al. (2007)	1
54520-2000 PJ ₃₀	Benecchi et al. (2011)	1
54598-Bienor	Delsanti et al. (2004)	
	Doressoundiram et al. (2002)	
	Tegler et al. (2003)	
	Dotto et al. (2003)	
	Bauer et al. (2003)	
	Doressoundiram et al. (2007)	
	DeMeo et al. (2009)	
	Rabinowitz et al. (2007)	15
55565-2002 AW ₁₉₇	Doressoundiram et al. (2005a)	13
33303 2002 1111 197	Fornasier et al. (2004)	
	Jewitt et al. (2007)	
	DeMeo et al. (2009)	
	Rabinowitz et al. (2007)	9
55576-Amycus	Peixinho et al. (2004)	,
55570-2 miyeus	Bauer et al. (2004)	
	Fornasier et al. (2004)	
	Doressoundiram et al. (2005a)	
	Doressoundiram et al. (2005a) Doressoundiram et al. (2007)	19
55636 2002 TV		19
55636-2002 TX ₃₀₀	Tegler et al. (2003)	
	Doressoundiram et al. (2005b)	
	Jewitt et al. (2007)	
	Ortiz et al. (2004)	
	Rabinowitz et al. (2008) Rabinowitz et al. (2007)	8

Table A.2. Continued

Object	References	Nr.
55637-2002 UX ₂₅	Jewitt et al. (2007)	
	Doressoundiram et al. (2007)	
	DeMeo et al. (2009)	
	Rabinowitz et al. (2007)	8
55638-2002 VE ₉₅	Rabinowitz et al. (2007)	_
	Barucci et al. (2006)	2
56968-2000 SA ₉₂	Fornasier et al. (2007)	1
58534-Logos	Jewitt & Luu (2001)	
	Boehnhardt et al. (2001)	
	Gil-Hutton & Licandro (2001)	
	Benecchi et al. (2011)	
	Barucci et al. (2000)	7
59524 Lagas I D	Benecchi et al. (2009)	7
58534-Logos+B	Benecchi et al. (2009)	1
59358-1999 CL ₁₅₈	Doressoundiram et al. (2002)	1
60454-2000 CH ₁₀₅	Peixinho et al. (2004)	2
60458 2000 CM	Benecchi et al. (2011)	2
60458-2000 CM ₁₁₄	Tegler et al. (2003) Benecchi et al. (2009)	2
60458-2000 CM _{114+B}		2 1
60558-Echeclus	Benecchi et al. (2009)	1
00530-ECHECIUS	Boehnhardt et al. (2002) Delsanti et al. (2006)	
	. ,	
	Bauer et al. (2003)	
	Jewitt (2009b)	8
60608-2000 EE ₁₇₃	DeMeo et al. (2009) Boehnhardt et al. (2002)	٥
50008-2000 EE ₁₇₃	Benecchi et al. (2002)	
	Tegler et al. (2003)	3
60620-2000 FD ₈	Boehnhardt et al. (2002)	3
00020-2000 FD ₈	Benecchi et al. (2001)	
	Peixinho et al. (2011)	3
60621-2000 FE ₈	Doressoundiram et al. (2002)	3
00021-2000 FE ₈	Benecchi et al. (2011)	
	Tegler et al. (2011)	3
63252-2001 BL ₄₁	Bauer et al. (2003)	3
03232-2001 BL ₄₁	Doressoundiram et al. (2003)	
	Peixinho et al. (2004)	
	Tegler et al. (2003)	10
65150-2002 CA ₁₂₆	Fornasier et al. (2007)	1
65225-2002 EK ₄₄	Fornasier et al. (2007)	1
65489-Ceto	Tegler et al. (2003)	•
0540) Ceto	Jewitt et al. (2007)	
	Benecchi et al. (2009)	3
65489-Ceto+B	Benecchi et al. (2009)	1
66452-1999 OF ₄	Peixinho et al. (2004)	
00.02 1999 014	Benecchi et al. (2011)	3
66652-Borasisi	Delsanti et al. (2001)	5
00032 Borasisi	McBride et al. (2003)	
	Delsanti et al. (2006)	
	Benecchi et al. (2011)	
	Benecchi et al. (2009)	
	Doressoundiram et al. (2001)	7
66652-Borasisi+B	Benecchi et al. (2009)	1
69986-1998 WW ₂₄	Doressoundiram et al. (2002)	•
	Peixinho et al. (2004)	3
69987-1998 WA ₂₅	Benecchi et al. (2011)	1
69988-1998 WA ₃₁	Benecchi et al. (2011)	1
07700 1770 111131	Peixinho et al. (2004)	2
69990-1998 WU ₃₁	Peixinho et al. (2004)	1
73480-2002 PN ₃₄	Tegler et al. (2004)	1
75 100 2002 11134	Doressoundiram et al. (2007)	
	Rabinowitz et al. (2007)	10
76804-2000 QE	Fornasier et al. (2007)	10
/U0U4-2UUU QE	romasier et al. (2007)	1

Table A.2. Continued

Object	References	Nr.
79360-1997 CS ₂₉	Romanishin & Tegler (1999)	
	Jewitt & Luu (2001)	
	Davies et al. (2000)	
	Boehnhardt et al. (2001)	
	Delsanti et al. (2006)	
	Benecchi et al. (2011) Jewitt et al. (2007)	
	Barucci et al. (2007)	
	Benecchi et al. (2009)	
	Tegler & Romanishin (1998)	15
79360-1997 CS _{29+B}	Benecchi et al. (2009)	1
79978-1999 CC ₁₅₈	Delsanti et al. (2001)	
150	Doressoundiram et al. (2002)	2
79983-1999 DF ₉	Doressoundiram et al. (2002)	1
80806-2000 CM ₁₀₅	Benecchi et al. (2011)	
	Benecchi et al. (2009)	2
82075-2000 YW ₁₃₄	Peixinho et al. (2004)	
	Benecchi et al. (2011)	
	Doressoundiram et al. (2005b)	
	Doressoundiram et al. (2007)	
	Jewitt et al. (2007)	
	Sheppard (2010)	1.4
82155-2001 F7:	Benecchi et al. (2009)	14
82155-2001 FZ ₁₇₃	Peixinho et al. (2004) Tegler et al. (2003)	
	Doressoundiram et al. (2007)	
	Jewitt et al. (2007)	9
82158-2001 FP ₁₈₅	Peixinho et al. (2004)	
200111183	Doressoundiram et al. (2005b)	
	Tegler et al. (2003)	
	Jewitt et al. (2007)	7
33982-Crantor	Peixinho et al. (2004)	
	Bauer et al. (2003)	
	Tegler et al. (2003)	
	Doressoundiram et al. (2005a)	
	Fornasier et al. (2004)	
	Doressoundiram et al. (2007)	
0.4522 2002 TG	DeMeo et al. (2009)	17
84522-2002 TC ₃₀₂	Sheppard (2010)	2
84700 2002 VIV	Rabinowitz et al. (2008)	2
84709-2002 VW ₁₂₀	Fornasier et al. (2007)	1 1
85627-1998 HP ₁₅₁ 85633-1998 KR ₆₅	Benecchi et al. (2011) Boehnhardt et al. (2002)	1
55555-1776 IXIX65	Tegler & Romanishin (2003)	
	Benecchi et al. (2011)	4
36047-1999 OY ₃	Tegler & Romanishin (2000)	7
	Boehnhardt et al. (2002)	
	Doressoundiram et al. (2002)	
	Benecchi et al. (2011)	
	Jewitt et al. (2007)	5
36177-1999 RY ₂₁₅	Boehnhardt et al. (2002)	
	Benecchi et al. (2011)	
	Doressoundiram et al. (2001)	3
87269-2000 OO ₆₇	Tegler et al. (2003)	
	Benecchi et al. (2011)	
7777 2000 07	Sheppard (2010)	3
37555-2000 QB ₂₄₃	Peixinho et al. (2004)	_
2020 2001 KE	Doressoundiram et al. (2007)	6
88269-2001 KF ₇₇	Tegler et al. (2003)	1
88611-Teharonhia	Benecchi et al. (2009)	1
88611-Teharonhia+B	Benecchi et al. (2009)	1
90377-Sedna	Barucci et al. (2005b)	
	Sheppard (2010)	3
90482-Orcus	Rabinowitz et al. (2007) Rabinowitz et al. (2004)	3
70+02-Ofcus	de Bergh et al. (2004)	
	Rabinowitz et al. (2007)	6
90568-2004 GV ₉	DeMeo et al. (2009)	U
70300-2004 G v 9		2
2000 200 r G 19	Rabinowitz et al. (2008)	2

Table A.2. Continued

Object	References	Nr.
91133-1998 HK ₁₅₁	Boehnhardt et al. (2001)	
	McBride et al. (2003)	
	Doressoundiram et al. (2002)	
	Benecchi et al. (2011)	
	Doressoundiram et al. (2001)	6
91205-1998 US ₄₃	Peixinho et al. (2004)	2
91554-1999 RZ ₂₁₅	Boehnhardt et al. (2002)	1
95626-2002 GZ ₃₂	Doressoundiram et al. (2005b)	
	Bauer et al. (2003)	
	Tegler et al. (2003)	
	Fornasier et al. (2004)	
	Doressoundiram et al. (2007)	
	Rabinowitz et al. (2007)	18
99328-2001 UY ₁₂₃	Fornasier et al. (2007)	1
105685-2000 SC ₅₁	Fornasier et al. (2007)	1
111113-2001 VK ₈₅	Fornasier et al. (2007)	1
118228-1996 TQ ₆₆	Romanishin & Tegler (1999)	
	Davies et al. (2000)	
	Gil-Hutton & Licandro (2001)	
	Jewitt & Luu (2001)	
	Benecchi et al. (2011)	
	Tegler & Romanishin (1998)	7
118379-1999 HC ₁₂	Boehnhardt et al. (2002)	
	Benecchi et al. (2011)	3
118702-2000 OM ₆₇	Sheppard (2010)	1
119068-2001 KC ₇₇	Tegler et al. (2003)	1
119070-2001 KP ₇₇	Benecchi et al. (2011)	1
119315-2001 SQ ₇₃	Tegler et al. (2003)	1
119951-2002 KX ₁₄	DeMeo et al. (2009)	
	Rabinowitz et al. (2007)	
	Romanishin et al. (2010)	3
119979-2002 WC ₁₉	Benecchi et al. (2009)	1
120061-2003 CO ₁	Tegler et al. (2003)	
-	Doressoundiram et al. (2007)	11
120132-2003 FY ₁₂₈	DeMeo et al. (2009)	
120	Sheppard (2010)	3
120178-2003 OP ₃₂	Rabinowitz et al. (2008)	1
120347-Salacia	Benecchi et al. (2009)	1
120347-Salacia+B	Benecchi et al. (2009)	1
120348-2004 TY ₃₆₄	Rabinowitz et al. (2007)	1
120453-1988 RE ₁₂	Fornasier et al. (2007)	1
121725-1999 XX ₁₄₃	Bauer et al. (2003)	
145	Doressoundiram et al. (2002)	
	Peixinho et al. (2004)	10
123509-2000 WK ₁₈₃	Benecchi et al. (2009)	1
123509-2000 WK _{183+B}	Benecchi et al. (2009)	1
124729-2001 SB ₁₇₃	Fornasier et al. (2007)	1
127546-2002 XU ₉₃	Sheppard (2010)	1
129772-1999 HR ₁₁	Doressoundiram et al. (2001)	1
130391-2000 JG ₈₁	Benecchi et al. (2011)	1
131695-2001 XS ₂₅₄	Benecchi et al. (2011)	1
134340-Pluto	Buratti et al. (2003)	1
134340-1 luto	Jewitt & Luu (2001)	
		4
124240 Pluto - P	Buie et al. (2006)	4 1
134340-Pluto+B 134340-Pluto+C	Buie et al. (2006)	_
	Buie et al. (2006)	1
134340-Pluto+D	Buie et al. (2006)	1
134860-2000 OJ ₆₇	Doressoundiram et al. (2002)	
	Delsanti et al. (2006)	
	Benecchi et al. (2011)	-
124040 2006 27	Benecchi et al. (2009)	5
134860-2000 OJ _{67+B}	Benecchi et al. (2009)	1
135182-2001 QT ₃₂₂	Romanishin et al. (2010)	1
136108-Haumea	Jewitt et al. (2007)	
	Rabinowitz et al. (2007)	
	Trujillo et al. (2007)	3
136199-Eris	Carraro et al. (2006)	
	DeMeo et al. (2009)	

Table A.2. Continued

Object	References	Nr.
136472-Makemake	Rabinowitz et al. (2007)	1
137294-1999 RE ₂₁₅	Boehnhardt et al. (2007)	1
13/274-1777 NE ₂₁₅	Benecchi et al. (2011)	3
137295-1999 RB ₂₁₆	Boehnhardt et al. (2002)	1
138537-2000 OK ₆₇	Delsanti et al. (2001)	1
130337 2000 01107	Delsanti et al. (2004)	
	Doressoundiram et al. (2002)	
	Benecchi et al. (2011)	5
144897-2004 UX ₁₀	Romanishin et al. (2010)	1
145451-2005 RM ₄₃	DeMeo et al. (2009)	-
110101 2000 14443	Rabinowitz et al. (2008)	2
145452-2005 RN ₄₃	DeMeo et al. (2009)	2
145453-2005 RR ₄₃	DeMeo et al. (2009)	_
110 100 2000 Tele43	Rabinowitz et al. (2008)	3
145480-2005 TB ₁₉₀	Sheppard (2010)	1
148209-2000 CR ₁₀₅	Tegler et al. (2003)	1
140207 2000 CR 105	Benecchi et al. (2011)	
	Sheppard (2010)	3
148780-Altjira	Benecchi et al. (2009)	1
148780-Altjira+B	Benecchi et al. (2009)	1
150642-2001 CZ ₃₁	Delsanti et al. (2006)	1
163135-2002 CT ₂₂	Fornasier et al. (2007)	1
163216-2002 EN ₆₈	Fornasier et al. (2007)	1
168700-2000 GE ₁₄₇	Benecchi et al. (2011)	1
168703-2000 GP ₁₈₃	Doressoundiram et al. (2002)	1
100703 2000 G1 183	Benecchi et al. (2011)	
	Peixinho et al. (2004)	5
181708-1993 FW	Green et al. (1997)	5
101,00 13,00 1	Tegler & Romanishin (2003)	
	Benecchi et al. (2011)	
	Barucci et al. (2000)	
	Romanishin et al. (1997)	8
181855-1998 WT ₃₁	Peixinho et al. (2004)	O
101033 1550 11131	Snodgrass et al. (2010)	3
181867-1999 CV ₁₁₈	Peixinho et al. (2004)	1
181871-1999 CO ₁₅₃	Benecchi et al. (2011)	1
181874-1999 HW ₁₁	Benecchi et al. (2011)	1
1010/11/333 11/11	Sheppard (2010)	2
181902-1999 RD ₂₁₅	Boehnhardt et al. (2002)	-
101302 1333 165213	Benecchi et al. (2011)	2
182397-2001 QW ₂₉₇	Sheppard (2010)	1
182933-2002 GZ ₃₁	Benecchi et al. (2009)	1
192388-1996 RD ₂₉	Fornasier et al. (2007)	1
192929-2000 AT ₄₄	Fornasier et al. (2007)	1
208996-2003 AZ ₈₄	Fornasier et al. (2004)	•
200770-2003 AL84	Doressoundiram et al. (2007)	
	DeMeo et al. (2009)	
	Rabinowitz et al. (2008)	9
	1000110 WILL OF U. (2000)	