

Effective algebraic integration in bounded genus

Jorge Vitório Pereira and Roberto Svaldi

ABSTRACT

We introduce and study birational invariants for foliations on projective surfaces built from the adjoint linear series of positive powers of the canonical bundle of the foliation. We apply the results in order to investigate the effective algebraic integration of foliations on the projective plane. In particular, we describe the Zariski closure of the set $\Sigma_{d,g}$ of foliations on \mathbb{P}^2 of degree d admitting rational first integrals with fibers having geometric genus bounded by g .

Contents

1	Introduction	1
2	Kodaira dimension of foliations	5
3	Effective threshold and adjoint dimension	10
4	Singularities	13
5	Non-isotrivial fibrations	15
6	Classification via adjoint dimension	19
7	Variation in moduli	22
8	Foliations with rational first integrals	25
	References	29

1. Introduction

1.1 Effective algebraic integration

It seems fair to say that the simplest class of algebraic ordinary differential equations consists of the class of equations having all its solutions algebraic. In general, given an explicit differential equation, it is a difficult problem to decide whether or not it belongs to this distinguished class. Perhaps the first positive result on the subject is Schwarz's list of parameters for which Gauss' hypergeometric equation has an algebraic solution [Sch73].

Motivated by this remarkable result, a lot of activity on the study of algebraic solutions of linear differential equations took place in the XIXth century leading to a fairly good understanding of the problem for homogeneous linear differential equations. Among the works dealing with this question one can

2010 *Mathematics Subject Classification* 37F75; 14E99

Keywords: Holomorphic foliations; effective algebraic integration; degree of invariant algebraic curves.

This collaboration initiated while both authors were visiting James M^cKernan at UCSD, and continued during a visit of the second author to IMPA. We are grateful to both institutions for the favorable working conditions. The first author is partially supported by Cnpq and FAPERJ. The second author was partially supported by NSF research grant no: 1200656 and no: 1265263. During the final revision of this work he was supported by funding from the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC Grant agreement no. 307119.

find contributions by Fuchs, Gordan, Jordan, Halphen, and Klein just to name a few. At that time, the community seemed to believe that it would be possible to decide whether or not all solutions of a given linear differential equations are algebraic, see for instance the concluding remarks¹ of [Kle56, Section 3, Chapter V].

By the end of XIXth century mathematicians like Painlevé, Autonne, and Poincaré [Hen91, Hen97] started to study the next case, that is, polynomial differential equations of first order and of first degree. In modern language, they studied foliations on the projective plane with special emphasis on the existence of methods/algorithms to decide whether or not all leaves are algebraic. We will call this general line of enquiry effective algebraic integration. The results obtained at that time relied on strong assumptions on the nature of the singularities of the foliations and were not considered definitive as one can learn from the Introduction² of [Hen97]. For a modern account of some of these classical results see [GZ97] and [Per03, Chapter 7].

The results of the XIXth century on the effective integration of linear differential equations were revisited in the course of the XXth century. It was then made clear that a full solution for the problem was not available, rather, the problem was reduced to a similar one for rank one linear differential equations over curves. More precisely, in order to be able to decide whether or not a homogeneous linear differential equation $P(x, y, y', y'', y''', \dots, y^{(n)}) = 0$ has all its solutions algebraic it suffices to be able to solve the following problem: given an element u belonging to an algebraic extension of the field $\mathbb{C}(x)$, decide if u is the logarithmic derivative of an element v also belonging to an algebraic extension of $\mathbb{C}(x)$. Some authors expressed doubts on the possibility of solving this problem. For instance, in [Har71, page 51] one can find the view of Hardy³ on the subject.

Despite the skepticism of Hardy and others (cf. [Ris70]), in the late 1960's Risch (loc. cit.) showed that this problem, in its turn, can be reduced to the following one: given an explicit divisor on an explicit algebraic curve C , decide whether or not such divisor is of finite order in the Jacobian of C . Risch proved that this problem can be solved by restricting the data modulo two distinct primes and using the resulting bounds in positive characteristic to devise an explicit bound in characteristic zero. For a detailed account on the case of second order homogeneous differential equations see [BD79]. The interested reader can find more about the history of effective algebraic integration of linear differential equations in [vdPS03, page 124], [Gra08, Chapter III], and the references therein.

The corresponding problem for (non-linear) differential equations of the first order and of the first degree is still wide open and received considerably less attention. After being dormant for a good while, the interest towards it has been revived by experts in foliation theory who considered the problem of bounding the degree of algebraic leaves of foliations on \mathbb{P}^2 , see for instance [CLN91, Car94, CC97, EK03] and references therein. The influence of arithmetic on the subject was rediscovered by Lins Neto [LN02] who determined algebraic families (pencils) of foliations on the projective plane with fixed number and analytical type of singularities and with algebraic leaves of arbitrarily large degree.

¹ “Thus is the problem, which we formulated at the beginning of this paragraph [present all linear homogenous differential equations of the second order with rational coefficients: $y'' + py' + qy = 0$ which possess altogether algebraic solutions], fully solved.”

² “Je me suis occupé de nouveau de la même question dans ces derniers temps, dan l’ espoir que je parviendrais à généraliser les résultats obtenus. Cet espoir a été déçu. J’ai obtenu cependant quelques résultats partiels, que je prends la liberté de publier, estimant qu’on pourra s’en servir plus tard pour obtenir, par un nouvel effort, une solution plus satisfaisante du problème.”

³ “But no method has been devised as yet by which we can always determine in a finite number of steps whether a given elliptic integral is pseudo-elliptic, and integrate it if it is, and there is reason to suppose that no such method can be given.”

1.2 Degenerations of planar foliations admitting a rational first integrals

This work investigates the problem of effective algebraic integration for foliations on projective surfaces. In order to focus the discussion and clarify the framework in which we are going to carry it, we introduce the following conjecture.

CONJECTURE 1.1. *The Zariski closure in $\mathbb{P}H^0(\mathbb{P}^2, T_{\mathbb{P}^2}(d-1))$ of the set of foliations of degree d on \mathbb{P}^2 which admit a rational integral consists of transversely projective foliations.*

This conjecture is inspired by a remark made by Painlevé⁴ ([Pai97, pp. 216–217]) in his Stockholm’s lectures. Knowledge of a transversely projective structure for a given foliation, in view of their recent description [CP14, LPT16], would allow to reduce the problem to either the determination of periods of differential forms – when, after passing to a ramified covering, the foliation is defined by a closed rational 1-form – or to the algebraic integrability of Riccati equations.

The main results of this paper provide evidence in favor of this conjecture and are obtained using birational techniques. More precisely, we use basic results on adjoint linear series, the birational classification of foliated surfaces according to their Kodaira dimension [McQ08, Bru04, Men00] to obtain a variant of the classification which we now proceed to explain.

1.3 Adjoint dimension of foliations

The works of the Italian school of algebraic geometry in the beginning of the XXth century showed how much of the geometry of a smooth projective surface X can be determined by the order of growth of the function

$$n \mapsto h^0(X, K_X^{\otimes n}).$$

Whenever this function grows slower than a quadratic polynomial, one has a rather precise description of the surface (the so called Enriques-Kodaira classification). A similar classification is also available in dimension three thanks to the works of the modern school of birational geometry, and there is also a similar picture in arbitrary dimensions conditional on the so-called Abundance Conjecture.

In the case of foliations on surfaces, McQuillan, Brunella and Mendes obtained a very precise classification – analogue to the Enriques-Kodaira classification – in terms of the Kodaira dimension of the foliation. As in the case of surfaces, the Kodaira dimension of a foliation \mathcal{F} , $\text{kod}(\mathcal{F})$, measures the growth of the function $h^0(X, K_{\mathcal{F}}^{\otimes n})$ where $K_{\mathcal{F}}$ is the bundle of holomorphic 1-forms along the leaves of the foliation.

As the terminology suggests the canonical bundle together with its dual are the most obvious naturally determined line bundles on a variety. Combined with the fact that the integers $h^0(X, K_X^{\otimes n})$ ($n > 0$) are birational invariants for smooth projective varieties, its study is rather natural if one wants to understand varieties birationally. For foliations of arbitrary dimension/codimension, besides the canonical bundle, one also has another naturally attached line bundle: the determinant of the conormal bundle. If \mathcal{F} is a foliation on a projective surface X with canonical singularities then it turns out that for arbitrary $n, m \geq 0$ the integers $h^0(X, K_{\mathcal{F}}^{\otimes n} \otimes N_{\mathcal{F}}^{\otimes m})$ are birational invariants. Most of the results obtained in this paper steam from this simple observation. We define the adjoint dimension of a foliation according to the order of growth of the function $h^0(X, K_{\mathcal{F}}^{\otimes n} \otimes N_{\mathcal{F}}^{\otimes m})$, see Section 3.

⁴“J’ajoute qu’on ne peut espérer résoudre d’un coup qui consiste à limiter n . L’énoncé vers lequel il faut tendre doit avoir la forme suivante: “On sait reconnaître si l’intégrale d’une équation $F(y', y, x) = 0$ donnée est algébrique ou ramener l’équation aux quadratures.” Dans ce dernier cas, la question reviendrait à reconnaître si une certaine intégrale abélienne (de première ou de troisième espèce) n’a que deux ou une périodes.”

Building on the classification of foliations on surfaces according to their Kodaira dimension, in Section 6 we present a classification in function of the adjoint dimension. The results we obtain are summarized in Table 1. The outcome of the classification provides a framework well-suited to deal with families of foliations (Section 7) mainly due to the fact that it is more flexible with respect to the type of singularities which are allowed (Section 4). The classification in terms of the adjoint dimension also reflects distinct cases of the problem of effective algebraic integration (Section 8).

adj	kod	Description
$-\infty$	$-\infty$	Rational fibration
	0	Finite quotient of Riccati foliation generated by global vector field
	1	Riccati foliation
0	0	Finite quotient of linear foliation on a torus
1	0	Finite quotient of $E \times C \rightarrow C$, $g(C) \geq 2$
	1	Finite quotient of $E \times C \rightarrow E$, $g(C) \geq 2$
	1	Turbulent foliation
	1	Non-isotrivial elliptic fibration
2	$-\infty$	Irreducible quotient of $\mathbb{H} \times \mathbb{H} \rightarrow \mathbb{H}$
	1	Finite quotient of $C_1 \times C_2 \rightarrow C_1$, $g(C_i) \geq 2$
	2	General type

TABLE 1. Classification of foliations according to their adjoint/Kodaira dimensions.

1.4 Plan of the paper and statement of main results

The bulk of the paper starts by reviewing the classification of foliations with respect to their Kodaira dimension in Section 2. Then we introduce new birational invariants for foliations on surfaces, notably the effective threshold and the adjoint dimension, in Section 3. Section 4 is devoted to the study of a variation of the concept of canonical singularities, the so-called ε -canonical singularities. We prove in Corollary 4.10 that, for $\varepsilon > 0$, this concept is stable for small perturbations of the singularity of the foliation. This fact will be particularly important in the study of families of foliations carried out in Section 7.

Section 5 is devoted to the proof of the boundedness of non-isotrivial fibrations of bounded genus in families, see Theorem 5.4. In the particular case of \mathbb{P}^2 , the result reads as follows.

THEOREM A. *Let \mathcal{F} be a foliation on \mathbb{P}^2 . Assume that \mathcal{F} is birationally equivalent to a non-isotrivial fibration of genus $g \geq 2$. Then the degree of the general leaf of \mathcal{F} is bounded by*

$$\left(4 \left(42(2g-2)\right)!\right)^2 (4g-4) \deg(\mathcal{F}).$$

Theorem A refines the main result of [Per02] where it was established the existence of a bound for the degree of the general leaf depending on its genus and on the first $k > 0$ for which the linear system $|K_{\mathcal{F}}^{\otimes k}|$ defines a rational map with two dimensional image. The existence of universal k working for every non-isotrivial fibration of genus g was not known then - and is still not known at present time - hence the existence of a bound depending only on the degree of the foliation and on the genus was unclear. In comparison to [Per02] the proof of the result above has two new ingredients. The first is a bound on multiplicities of irreducible components of fibers of relatively minimal non-isotrivial fibrations of genus $g \geq 2$, cf. Proposition 5.3. The second new ingredient is the use of standard results on adjoint linear series (recalled in Section 5.1) in order to obtain effective $(n, m) \in \mathbb{N}^2$ such that the rational map

defined by $|K_{\mathcal{F}}^{\otimes n} \otimes K_X^{\otimes m}|$ has two dimensional image. By imposing further assumptions on the nature of the singularities of a foliation on \mathbb{P}^2 we obtain significantly better bounds (sub-linear on g), refining a classical result of Poincaré, cf. Theorem 5.6.

In Section 6 we carry out the classification of foliations on surfaces according to the adjoint dimension, see Table 1. The proof strongly relies on the classification of foliations according to the Kodaira dimension, but it does need to dwell with its subtlest point: the classification of non-abundant foliations. A nice corollary of the classification is a cohomological characterization of rational fibrations, which is a weak analogue of Castelnuovo's Criterion for the rationality of surfaces, cf. [Bea96, Thm. V.1].

THEOREM B. *Let \mathcal{F} be a foliation with at worst canonical singularities on a smooth projective surface X . The foliation \mathcal{F} is a rational fibration if, and only if, $h^0(X, K_{\mathcal{F}}^{\otimes n} \otimes N_{\mathcal{F}}^{\otimes m}) = 0$ for every $n \geq 1$ and every $m > 0$.*

Section 7 investigates families of foliations. There it is shown that the set of effective thresholds in a family does not accumulate at zero (Theorem 7.6). More important, it prepares the ground for the proof of the most compelling evidence we have so far in favor of Conjecture 1.1.

THEOREM C. *The Zariski closure in $\mathbb{P}(H^0(\mathbb{P}^2, T_{\mathbb{P}^2}(d-1)))$ of the set of degree d foliations admitting a rational first integral with general fiber of genus $\leq g$ is formed by transversely projective foliations.*

Its proof is presented in Section 8 and relies on Theorem A, on the birational classification of foliations, and on basic properties of families of foliations.

ACKNOWLEDGMENTS. *The authors wish to thank Calum Spicer for many valuable conversations, Carlo Gasbarri for suggesting an improvement of the bounds in Theorem A, and the anonymous referees for suggesting many improvements in the presentation of this work.*

2. Kodaira dimension of foliations

We start things off by reviewing the birational classification of foliations on surfaces following [McQ08] and [Bru04]. No new results are presented in this section. We have only included proofs of a few key properties of the Zariski decomposition of the canonical bundle of a foliation which will be used in the sequel.

2.1 Singularities of foliations

DEFINITION 2.1. *Let \mathcal{F} be a foliation on X and let $\pi : Y \rightarrow X$ be a birational morphism. Denote by \mathcal{G} the pull-back of \mathcal{F} under π . If E is an exceptional divisor of π then the discrepancy of \mathcal{F} along E is*

$$a(\mathcal{F}, E) = \text{ord}_E(K_{\mathcal{G}} - \pi^* K_{\mathcal{F}}).$$

DEFINITION 2.2. *Let \mathcal{F} be a foliation on X . A point $x \in X$ is canonical for \mathcal{F} if and only if $a(\mathcal{F}, E) \geq 0$ for every divisor E over x . A point $x \in X$ is log canonical for \mathcal{F} if and only $a(\mathcal{F}, E) \geq -1$ for every divisor E over x .*

EXAMPLE 2.3. *Consider the pencil of foliations on $X = \mathbb{P}^2$ defined by the vector fields $sx \frac{\partial}{\partial x} + ty \frac{\partial}{\partial y}$ where $(s : t) \in \mathbb{P}^1$. If $s \cdot t \cdot (s - t) \neq 0$ then $\mathcal{F}_{(s:t)}$ is a foliation with trivial canonical bundle and three singularities at the points $(0 : 0 : 1)$, $(0 : 1 : 0)$, and $(1 : 0 : 0)$. For $(s : t) \notin \mathbb{P}^1(\mathbb{Q})$ the three singularities are canonical. For $(s : t) \in \mathbb{P}^1(\mathbb{Q}) - \{(0 : 1), (1 : 0), (1 : 1)\}$, two of the singularities are log canonical but not canonical, while the third singularity is canonical. Finally, when $s \cdot t \cdot (s - t) = 0$, the vector field will have one of the coordinate axis as a line of singularities. The corresponding foliation will have canonical bundle $\mathcal{O}_{\mathbb{P}^2}(-1)$ and only one singularity which is log canonical but not canonical.*

Any foliation on a projective surface is birationally equivalent to a foliation having at worst canonical singularities thanks to the following which is essentially due to Seidenberg.

THEOREM 2.4. *Let \mathcal{F} be a foliation on a smooth projective surface X . Then there exists a finite composition of blow-ups $\pi : Y \rightarrow X$ such that all the singularities of $\pi^*\mathcal{F}$ are canonical.*

2.2 Kodaira dimension

DEFINITION 2.5. *Let \mathcal{F} be a foliation with at worst canonical singularities on a smooth projective surface X . The Kodaira dimension of \mathcal{F} , $\text{kod}(\mathcal{F})$, is by definition*

$$\text{kod}(\mathcal{F}) := \text{kod}(K_{\mathcal{F}}) = \max_{m \in \mathbb{N}} \{\dim \phi_m(X)\},$$

where $\phi_m : X \dashrightarrow \mathbb{P}(H^0(X, K_{\mathcal{F}}^{\otimes m})^*)$ and we adopt the convention that $\dim \phi_m(X) = -\infty$ when $h^0(X, K_{\mathcal{F}}^{\otimes m}) = 0$. (and it is not possible to define the associated map).

The numerical Kodaira dimension of \mathcal{F} , $\nu(\mathcal{F})$, is defined to be the numerical dimension of $K_{\mathcal{F}}$, that is:

- $\nu(\mathcal{F}) = -\infty$ if $K_{\mathcal{F}}$ is not pseudoeffective, while
- if $K_{\mathcal{F}}$ is pseudoeffective with Zariski decomposition $K_{\mathcal{F}} = P + N$ then $\nu(\mathcal{F}) = 0$ if P is numerically zero, $\nu(\mathcal{F}) = 1$ if $P \neq 0$ but $P^2 = 0$, and $\nu(\mathcal{F}) = 2$ if $P^2 > 0$.

The classification of foliations with negative numerical Kodaira dimension is stated in the next result is due to Miyaoka.

THEOREM 2.6. *Let \mathcal{F} be a foliation on a projective surface X . If $K_{\mathcal{F}}$ is not pseudoeffective then \mathcal{F} is birationally equivalent to a \mathbb{P}^1 -bundle over a curve.*

2.3 Relatively minimal models

DEFINITION 2.7. *Let \mathcal{F} be a foliation with canonical singularities on a smooth projective surface X . An irreducible curve $C \subset X$ is called \mathcal{F} -exceptional if $K_X \cdot C = -1$ (i.e. $C \simeq \mathbb{P}^1$ and $C^2 = -1$) and the contraction of C gives rise to a foliation with canonical singularities.*

DEFINITION 2.8. *Let \mathcal{F} be a foliation with canonical singularities on a smooth projective surface X . A relatively minimal model for \mathcal{F} is the datum of a foliation \mathcal{G} with canonical singularities and without \mathcal{G} -exceptional curves on a smooth projective surface Y which is birationally equivalent to \mathcal{F} . We say that \mathcal{G} is a minimal model if for any birational map $\pi : Z \dashrightarrow Y$ and any foliation \mathcal{H} on Z with canonical singularities such that $\pi_*\mathcal{H} = \mathcal{G}$, π is a birational morphism.*

The definitions above and the next result are essentially due to Brunella [Bru99]. The only minor difference is that in the original definition of \mathcal{F} -exceptional curve Brunella only considered reduced singularities instead of canonical singularities. Nonetheless, his proof works also in this slightly more general situation.

THEOREM 2.9. *Let \mathcal{F} be a foliation with at worst canonical singularities on a smooth surface X . There exists a birational morphism $\pi : X \rightarrow Y$ such that $\pi_*\mathcal{F}$ is a relatively minimal model for \mathcal{F} . Moreover, $\pi_*\mathcal{F}$ is a minimal model for \mathcal{F} unless \mathcal{F} is birationally equivalent to a rational fibration, a Riccati foliation, or Brunella's special foliation \mathcal{H} .*

The reader will find the explicit construction of the foliation \mathcal{H} from the theorem in the paper just cited.


REMARK 2.10. *The above theorem highlights one of the differences between the birational classification of projective surfaces and that of foliations on surfaces: while surfaces of non-negative Kodaira dimension always have a unique minimal model, there are foliations of Kodaira dimension zero and one which do not have unique minimal models.*

2.4 Zariski decomposition and nef models

If \mathcal{L} is a pseudoeffective line bundle on a smooth projective surface then \mathcal{L} is \mathbb{Q} -linearly equivalent to $P_{\mathcal{L}} + N_{\mathcal{L}}$ where $P_{\mathcal{L}}$ is a nef \mathbb{Q} -divisor and $N_{\mathcal{L}}$ is a contractible effective \mathbb{Q} -divisor satisfying $P_{\mathcal{L}} \cdot N_{\mathcal{L}} = 0$. This is the so-called Zariski decomposition of \mathcal{L} . We will denote by $i(\mathcal{F})$ the index of $K_{\mathcal{F}}$, i.e., the minimum of the set $\{n \in \mathbb{N} \mid nN \text{ has integral coefficients}\}$.

THEOREM 2.11. [Bru04, Chapter 8, Theorem 1] *Let \mathcal{F} be a relatively minimal foliation on a smooth projective surface X . If $K_{\mathcal{F}}$ is pseudoeffective and $P + N$ is its Zariski decomposition then the support of N is a disjoint union of Hirzebruch-Jung strings.*

A Hirzebruch-Jung string is a chain of smooth rational curves of self-intersection at most -2 . At one end of the chain, the handle of the Hirzebruch-Jung string, the foliation has only one singularity. Every other curve in the chain contains two singularities of the foliation. There is only one singularity of \mathcal{F} on the Hirzebruch-Jung string which does not coincide with a singularity of its support. There exists a unique leaf of \mathcal{F} not contained in the Hirzebruch-Jung string that passes through this singularity. Such curve is called the tail of the Hirzebruch-Jung string.



DEFINITION 2.12. *Let \mathcal{F} be a relatively minimal foliation with pseudoeffective $K_{\mathcal{F}}$ on a smooth projective surface X . The order of a maximal Hirzebruch-Jung string contained in the support of N is the determinant of the negative of the intersection matrix of its support.*

The following proposition shows that the order and the index are closely related.

PROPOSITION 2.13. *Notation as in the definition above. The following assertions hold true.*

- (i) *The order of a maximal Hirzebruch-Jung string J contained in the support of N coincides with the smallest $o \in \mathbb{N}$ such that the coefficients of N corresponding to curves in J belong to $\frac{1}{o}\mathbb{N}$.*
- (ii) *The contraction of a Hirzebruch-Jung string of order o is locally isomorphic to the quotient of a smooth foliation on $(\mathbb{C}^2, 0)$ by the cyclic group generated by an automorphism of the form $(x, y) \mapsto (\xi_o \cdot x, \xi_o^a \cdot y)$ where ξ_o is a primitive root of unity of order o and a is a natural number relatively prime to o .*

Proof. The statement is local so we may very well assume that the support of N is connected. Let us write $N = \sum_{i=1}^k a_i E_i$ where E_i are the irreducible components of N . We denote by E_1 the handle of the Hirzebruch-Jung string while the other curves are numbered following the order in which they appear in the chain.

Let $A = (E_i \cdot E_j)_{i,j}$ be the intersection matrix of the Hirzebruch-Jung string and let $o = \det(-A)$ be the order of the Hirzebruch-Jung string. To determine the coefficients a_1, \dots, a_k we have to solve the linear system $(-A) \cdot (a_1, a_2, \dots, a_k)^T = (1, 0, \dots, 0)^T$. Therefore the coefficients a_i certainly lie in $\frac{1}{o}\mathbb{N}$. To see that o is the minimal number with such property it suffices to notice that $a_k = 1/o$, cf. [McQ08, proof of Proposition III.1.4]. This proves item (1). Item (2) is [McQ08, Reinterpretation III.2.bis.3.a] \square

In the Lemma below, we collect some properties of tails of Hirzebruch-Jung strings for later use.

LEMMA 2.14. *Let \mathcal{F} be a relatively minimal foliation with pseudoeffective canonical bundle on a smooth projective surface X . Let T be an irreducible invariant curve not contained in the support of N and let o_1, \dots, o_k be the orders of Hirzebruch-Jung strings intersecting T . Then the following assertions hold true.*

- (i) *The intersection of the positive part of the Zariski decomposition of $K_{\mathcal{F}}$ with T is given by the formula*

$$P \cdot T = K_{\mathcal{F}} \cdot T - \sum_{i=1}^k \frac{1}{o_i}.$$

- (ii) *If \mathcal{F} admits a holomorphic first integral $f : U \rightarrow \mathbb{C}$ defined on a \mathcal{F} -invariant neighborhood of T which vanishes along T then the vanishing order along T is a multiple of the least common multiple of o_1, \dots, o_k .*

Proof. Item (1) is [McQ08, Remark III.1.3.a]. To verify item (2) let us work locally on a neighborhood V of a Hirzebruch-Jung string intersecting T . Let $\pi : V \rightarrow W$ be the contraction of the Hirzebruch-Jung string we are considering and o be its order. Perhaps after restricting V to a smaller neighborhood we can assume that W is the quotient of a neighborhood \tilde{V} of the origin in \mathbb{C}^2 by a cyclic group generated by $\varphi(x, y) = (\xi_o \cdot x, \xi_o^a \cdot y)$ according to Proposition 2.13. We can also assume that the pull-back \mathcal{G} of $\pi_*(\mathcal{F}|_V)$ to \tilde{V} is the foliation defined by the level sets of the coordinate function y . The pull-back of $\pi_*(f|_V)$ to \tilde{V} is a holomorphic function g constant along the leaves of \mathcal{G} . The φ -invariance of g implies that $g(x, y) = h(y^o)$ for some one-variable holomorphic function h . Item (2) follows. \square

DEFINITION 2.15. *Let \mathcal{F} be a relatively minimal foliation on a smooth surface X with pseudoeffective canonical divisor. The nef model of \mathcal{F} is the foliation obtained by contracting the negative part of the Zariski decomposition of $K_{\mathcal{F}}$.*

2.5 Canonical models

DEFINITION 2.16. *A foliation \mathcal{F} on a normal projective surface X is called a canonical model if $K_{\mathcal{F}}$ is nef and $K_{\mathcal{F}} \cdot C = 0$ implies $C^2 \geq 0$ for every irreducible curve $C \subset X$.*


THEOREM 2.17. [McQ08, Theorem 3.3.2] *Let \mathcal{F} be relatively minimal foliation with $K_{\mathcal{F}}$ pseudoeffective on a smooth surface X . Then there exists a morphism $\pi : X \rightarrow Y$ from X to a normal projective surface Y such that $\mathcal{G} = \pi_*\mathcal{F}$ is a canonical model. The singular points of Y and the corresponding exceptional fibers of π are of one of the following forms.*

- (i) *The singular point is a cyclic quotient singularity and the exceptional divisor over it is a chain of rational curves of self-intersection at most -2*




The foliation around the singular point is the quotient of a smooth foliation; or the quotient of a canonical foliation singularity on a (germ of) smooth surface;

- (ii) *The singular point is dihedral quotient singularity and the exceptional divisor over it has the following dual graph:*



The foliation around the singularity is again the quotient of a smooth foliation or of a canonical singularity on a (germ of) smooth surface.

- (iii) The singular point is an elliptic Gorenstein singularity and the exceptional divisor is a cycle of smooth rational curves each of self-intersection at most -2 ; or a unique nodal rational curve of negative self-intersection



The foliation around the singular point is isomorphic to a cusp of a Hilbert modular foliation (cf. [McQ08, Theorem IV.2.2]). Moreover, the canonical bundle of the foliation on the canonical model is never \mathbb{Q} -Cartier.

When compared with the theory for projective surfaces, item (3) of the above Theorem is quite surprising. The fact that the canonical bundle is never \mathbb{Q} -Cartier is a clear obstruction to the base point freeness of $|K_{\mathcal{F}}^{\otimes n}|$ and for the finite generation of the canonical algebra of the foliation. It turns out that this is the only obstruction, cf. [McQ08, Corollary IV.2.3].

2.6 Kodaira dimension zero

THEOREM 2.18. [McQ08, Fact IV.3.3] *Let \mathcal{F} be a relatively minimal foliation on a smooth projective surface X with $\nu(\mathcal{F}) = 0$. Let $\pi : X \rightarrow Z$ be the contraction of the negative part of $K_{\mathcal{F}}$, i.e. $\pi_*\mathcal{F}$ is a nef model for \mathcal{F} . Then there exists a smooth projective surface Y and a quasi-étale cyclic covering $p : Y \rightarrow Z$ of degree $i(\mathcal{F})$ such that $p^*\pi_*\mathcal{F}$ is a foliation with trivial canonical bundle. In particular, $\text{kod}(\mathcal{F}) = 0$.*

The resulting surface Y belongs to the following list:

- (i) Product of a hyperbolic curve and an elliptic curve;
- (ii) Abelian surfaces;
- (iii) Projective bundle over an elliptic curve;
- (iv) Rational surface.

Consequently the klt surface Z has Kodaira dimension 1, 0, or $-\infty$ according to whether Y fits in case (1), (2), or (3)/(4). One can also determine the possibilities for the index of \mathcal{F} . This is done in [Per02]. There it is shown that

$$i(\mathcal{F}) \in \{1, 2, 3, 4, 5, 6, 8, 10, 12\} \quad (2.1)$$

when \mathcal{F} has Kodaira dimension zero.

2.7 Kodaira dimension one

The classification of foliations of Kodaira dimension one is essentially due to Mendes, see [Men00, Theorem 3.3.1].

THEOREM 2.19. *Let \mathcal{F} be a relatively minimal foliation on a smooth projective surface X . Assume that $\text{kod}(\mathcal{F}) = 1$ and let $f : X \rightarrow C$ be the Iitaka fibration of $K_{\mathcal{F}}$. If \mathcal{F} coincides with the foliation defined by f then f is non-isotrivial elliptic fibration. Otherwise \mathcal{F} is completely transverse to a general fiber F of f and we have the following possibilities:*

- (i) *The genus of F is zero and \mathcal{F} is a Riccati foliation; or*
- (ii) *The genus of F is one and \mathcal{F} is a turbulent foliation; or*
- (iii) *The genus of F is at least two and \mathcal{F} is an isotrivial fibration of genus at least two.*

2.8 Non-abundant foliations

The most striking difference between the birational classification of projective surfaces and the classification of rank one foliations in dimension two is the existence of foliations having canonical bundle with numerical dimension one and negative Kodaira dimension. This phenomenon is restricted to a rather special class of foliations as pointed out by the next result.

THEOREM 2.20. *Let \mathcal{F} be a relatively minimal foliation on a smooth projective surface X . If the numerical dimension of \mathcal{F} does not coincide with the Kodaira dimension of \mathcal{F} then*

- (i) $\nu(\mathcal{F}) = 1$,
- (ii) $\text{kod}(\mathcal{F}) = -\infty$,
- (iii) *X is the minimal desingularization of the Baily-Borel compactification of an irreducible quotient of $\mathbb{H} \times \mathbb{H}$, and*
- (iv) *\mathcal{F} is induced by one of the two natural fibrations on $\mathbb{H} \times \mathbb{H}$.*

Arguably this result constitutes the hardest part of the classification of foliations. The known proofs of this result rely heavily on Brunella's plurisubharmonic variation of the Poincaré metric and were obtained by Brunella and McQuillan in a collaborative effort.

In Section 6 we will carry out a classification of foliations in terms of another birational invariant. This new classification relies heavily on the original classification of surface foliations according to their Kodaira dimension; nonetheless, it does not need the full power of it. In particular, all that we need to know about non-abundant foliations is contained in the following Lemma.

LEMMA 2.21. *Let \mathcal{F} be a relatively minimal foliation with $\nu(\mathcal{F}) = 1$ and $\text{kod}(\mathcal{F}) = -\infty$. Then $h^1(X, \mathcal{O}_X) = 0$ and $P \cdot N_{\mathcal{F}}^* = P \cdot K_X > 0$ where P is the positive part of the Zariski decomposition of $K_{\mathcal{F}}$.*

Proof. If $h^1(X, \mathcal{O}_X) = h^0(X, \Omega_X^1) \neq 0$ then the restriction of a holomorphic 1-form to the leaves of \mathcal{F} either vanishes identically or gives rise to a non-zero section of $K_{\mathcal{F}}$. Thus if $\text{kod}(\mathcal{F}) = -\infty$ we obtain that \mathcal{F} factors through the Albanese map of X and is a fibration. Hence $\text{kod}(\mathcal{F}) \geq 0$ contrary to our assumptions. Thus $h^1(X, \mathcal{O}_X) = 0$.

Since $h^1(X, \mathcal{O}_X) = 0$ we obtain that $\chi(\mathcal{O}_X) \geq 1$. Let $\mathcal{L} = \mathcal{O}_X(mP)$ where m is a sufficiently divisible positive integer. By Riemann-Roch,

$$\chi(\mathcal{L}) = \chi(\mathcal{O}_X) + 1/2(m^2 P^2 - mP \cdot K_X)$$

If $P \cdot K_X < 0$ then $\chi(\mathcal{L}) > 0$. Thus $h^0(X, \mathcal{L}) + h^2(X, \mathcal{L}) > 0$. But if m is sufficiently large then $K_X \otimes \mathcal{L}^*$ is not pseudoeffective and consequently $h^2(X, \mathcal{L}) = h^0(X, K_X \otimes \mathcal{L}^*) = 0$. It follows that $h^0(X, K_{\mathcal{F}}^{\otimes m}) = h^0(X, \mathcal{L}) > 0$, contradicting $\text{kod}(\mathcal{F}) = -\infty$. \square

3. Effective threshold and adjoint dimension

In this section we define the effective threshold and the adjoint dimension of a foliation on a smooth projective surface and prove their birational invariance.

3.1 Effective threshold

DEFINITION 3.1. Let \mathcal{F} be a foliation with canonical singularities on a smooth projective surface X . If the canonical bundle of \mathcal{F} is pseudoeffective then we define the effective threshold of \mathcal{F} , $\text{eff}(\mathcal{F})$, as the largest $\varepsilon \in \mathbb{R}_{\geq 0} \cup \{\infty\}$ such that $K_{\mathcal{F}} + \varepsilon N_{\mathcal{F}}^*$ is pseudoeffective. If $K_{\mathcal{F}}$ is not pseudoeffective, then we set $\text{eff}(\mathcal{F}) = -\infty$.

EXAMPLE 3.2. Let \mathcal{F} be a very general foliation on \mathbb{P}^2 of degree d . It is well known that \mathcal{F} has reduced, and in particular canonical, singularities. Recall that the degree of \mathcal{F} is defined as the number of tangencies between \mathcal{F} and a general line. In this case $K_{\mathcal{F}} = \mathcal{O}_{\mathbb{P}^2}(d-1)$ and $N_{\mathcal{F}}^* = \mathcal{O}_{\mathbb{P}^2}(-d-2)$. If $d = 0$ then $K_{\mathcal{F}}$ is not pseudoeffective. If instead $d \geq 1$ then $K_{\mathcal{F}}$ is pseudoeffective and

$$\text{eff}(\mathcal{F}) = \frac{d-1}{d+2}.$$

The reader should notice that $\text{eff}(\mathcal{F}) < 1$ for every foliation on \mathbb{P}^2 .

This is by no means a coincidence since $K_X = K_{\mathcal{F}} + N_{\mathcal{F}}^*$ and foliations on a surface X of negative Kodaira dimension will always have $\text{eff}(\mathcal{F}) < 1$ as K_X is not pseudoeffective. If instead X has non-negative Kodaira dimension then K_X is pseudoeffective and consequently $\text{eff}(\mathcal{F}) \geq 1$ for every foliation on X .

Similarly, one sees that $\text{eff}(\mathcal{F}) = \infty$ if and only if both $K_{\mathcal{F}}$ and $N_{\mathcal{F}}^*$ are pseudoeffective. Foliations with pseudoeffective conormal bundle have recently been classified by Touzet, [Tou16]. They fit in one of the following descriptions:

- (i) after a finite étale cover \mathcal{F} is defined by a closed holomorphic 1-form; or
- (ii) there exists a morphism from X to a quotient of a polydisc \mathbb{D}^m by an irreducible lattice and \mathcal{F} is the pull-back of one of the m tautological foliations on the polydisc. In particular \mathcal{F} is transversely hyperbolic.

Notice that the dimension of the ambient manifold is not necessarily equal to the dimension of the polydisc.

REMARK 3.3. Using the identity $K_X = K_{\mathcal{F}} + N_{\mathcal{F}}^*$ we can write

$$K_{\mathcal{F}} + \varepsilon N_{\mathcal{F}}^* = (1 - \varepsilon)(K_{\mathcal{F}} + \frac{\varepsilon}{1 - \varepsilon} K_X),$$

when $\varepsilon \neq 1$.

When $\text{eff}(\mathcal{F})$ is small, we will often work with $K_{\mathcal{F}} + \varepsilon K_X$ as that is more convenient.

3.2 Adjoint dimension

DEFINITION 3.4. Let \mathcal{F} be a foliation with canonical singularities on a projective surface X . Consider the pluricanonical maps

$$\varphi_{m,n} : X \dashrightarrow \mathbb{P}H^0(X, K_{\mathcal{F}}^{\otimes m} \otimes N_{\mathcal{F}}^{*\otimes n})^*$$

for $m \geq 1, n \geq 1$. The adjoint dimension of \mathcal{F} , denoted $\text{adj}(\mathcal{F})$, is the maximal dimension of the image of these maps. If $h^0(X, K_{\mathcal{F}}^{\otimes m} \otimes N_{\mathcal{F}}^{*\otimes n}) = 0$ for every $m, n \geq 1$ then we set $\text{adj}(\mathcal{F}) = -\infty$.

DEFINITION 3.5. Let \mathcal{F} be a foliation with canonical singularities on a projective surface X . The numerical adjoint dimension of \mathcal{F} , $\text{adj}_{\text{num}}(\mathcal{F})$, is equal to $-\infty$ if $\text{eff}(\mathcal{F}) \leq 0$ and equal to the maximal numerical dimension of $K_{\mathcal{F}} + \varepsilon N_{\mathcal{F}}^*$ for $\varepsilon \in (0, \text{eff}(\mathcal{F}))$ otherwise.

Of course $\text{adj}(\mathcal{F}) \leq \text{adj}_{\text{num}}(\mathcal{F})$.

3.3 Birational invariance

The significance of the concepts of effective threshold and of (numerical) adjoint dimension for the purpose of the birational classification of foliations on surfaces is assured by the next proposition.

PROPOSITION 3.6. *Let (X, \mathcal{F}) and (Y, \mathcal{G}) be two birationally equivalent foliations. If \mathcal{F} and \mathcal{G} have at worst canonical singularities then $\text{eff}(\mathcal{F}) = \text{eff}(\mathcal{G})$, $\text{adj}(\mathcal{F}) = \text{adj}(\mathcal{G})$ and $\text{adj}_{\text{num}}(\mathcal{F}) = \text{adj}_{\text{num}}(\mathcal{G})$. Furthermore, $h^0(X, K_{\mathcal{F}}^{\otimes n} \otimes N_{\mathcal{F}}^{*\otimes m}) = h^0(Y, K_{\mathcal{G}}^{\otimes n} \otimes N_{\mathcal{G}}^{*\otimes m})$ for every $n, m \geq 0$.*

Proof. Since we can choose a foliation (Z, \mathcal{H}) on a smooth projective surface Z dominating both (X, \mathcal{F}) and (Y, \mathcal{G}) , there is no loss of generality in assuming the existence of a birational morphism $\pi : (X, \mathcal{F}) \rightarrow (Y, \mathcal{G})$. In fact, we can even assume (and will) that π is the blow-up of a point $p \in Y$. Let E be the exceptional divisor.

We will first prove that $\text{eff}(\mathcal{F}) = \text{eff}(\mathcal{G})$. First notice that $K_{\mathcal{G}} + \varepsilon N_{\mathcal{G}}^* = \pi_*(K_{\mathcal{F}} + \varepsilon N_{\mathcal{F}}^*)$. Therefore if $K_{\mathcal{F}} + \varepsilon N_{\mathcal{F}}^*$ is pseudoeffective then the same holds true for $K_{\mathcal{G}} + \varepsilon N_{\mathcal{G}}^*$. This shows that $\text{eff}(\mathcal{G}) \geq \text{eff}(\mathcal{F})$. To prove the converse inequality, we will need to use that \mathcal{G} has canonical singularities. Since π is the blow-up of a point by assumption, we have that $K_{\mathcal{F}} - \pi^* K_{\mathcal{G}} = aE$ for some $a \in \{0, 1\}$. Since $K_X - \pi^* K_Y = E$ we also have that $N_{\mathcal{F}}^* - \pi^* N_{\mathcal{G}}^* = (1 - a)E$, and consequently

$$K_{\mathcal{F}} + \varepsilon N_{\mathcal{F}}^* = \pi^*(K_{\mathcal{G}} + \varepsilon N_{\mathcal{G}}^*) + (a + \varepsilon(1 - a))E. \quad (3.1)$$

Therefore, if $K_{\mathcal{G}} + \varepsilon N_{\mathcal{G}}^*$ is pseudoeffective then the same holds true for $K_{\mathcal{F}} + \varepsilon N_{\mathcal{F}}^*$. We conclude that $\text{eff}(\mathcal{G}) \leq \text{eff}(\mathcal{F})$ and the equality between the effective thresholds follow.

Let us now prove that $\text{adj}_{\text{num}}(\mathcal{F}) = \text{adj}_{\text{num}}(\mathcal{G})$. We have just seen that $\text{eff}(\mathcal{F}) = \text{eff}(\mathcal{G})$. Then, $\text{adj}_{\text{num}}(\mathcal{F}) = -\infty$ if and only if $\text{adj}_{\text{num}}(\mathcal{G}) = -\infty$, by definition. Let us consider $\varepsilon \in (0, \text{eff}(\mathcal{F}))$ and let

$$\begin{aligned} K_{\mathcal{F}} + \varepsilon N_{\mathcal{F}}^* &= P_{\mathcal{F}, \varepsilon} + N_{\mathcal{F}, \varepsilon}, \\ K_{\mathcal{G}} + \varepsilon N_{\mathcal{G}}^* &= P_{\mathcal{G}, \varepsilon} + N_{\mathcal{G}, \varepsilon} \end{aligned}$$

be the Zariski decompositions of the divisors under scrutiny. By [Pro03, Theorem 2.2] and (3.1), it follows that for any $\varepsilon \in (0, \text{eff}(\mathcal{F}))$, $P_{\mathcal{F}, \varepsilon} = \pi^* P_{\mathcal{G}, \varepsilon}$ and $N_{\mathcal{F}, \varepsilon} = \pi^* N_{\mathcal{G}, \varepsilon} + a + \varepsilon(1 - a)$. Since by definition $\text{adj}_{\text{num}}(\mathcal{F}) = \max \nu(P_{\mathcal{F}, \varepsilon}) \mid \varepsilon \in (0, \text{eff}(\mathcal{F}))$ (and analogously for $\text{adj}_{\text{num}}(\mathcal{G})$), we have verified the desired equality.

To conclude the proof of the proposition it suffices to verify that $h^0(X, K_{\mathcal{F}}^{\otimes n} \otimes N_{\mathcal{F}}^{*\otimes m}) = h^0(Y, K_{\mathcal{G}}^{\otimes n} \otimes N_{\mathcal{G}}^{*\otimes m})$ for every $n, m \geq 0$. Once these equalities are proved, the equality $\text{adj}(\mathcal{F}) = \text{adj}(\mathcal{G})$ follows. Let us fix $n, m \geq 0$. From the isomorphism $K_{\mathcal{F}}^{\otimes n} \otimes N_{\mathcal{F}}^{*\otimes m} = \pi^*(K_{\mathcal{G}}^{\otimes n} \otimes N_{\mathcal{G}}^{*\otimes m}) \otimes \mathcal{O}_X((na + m(1 - a))E)$ we deduce the short exact sequence

$$0 \rightarrow \pi^*(K_{\mathcal{G}}^{\otimes n} \otimes N_{\mathcal{G}}^{*\otimes m}) \rightarrow K_{\mathcal{F}}^{\otimes n} \otimes N_{\mathcal{F}}^{*\otimes m} \rightarrow \mathcal{O}_E((na + m(1 - a))E) \rightarrow 0.$$

Since $h^0(E, \mathcal{O}_E((na + (1 - a))E)) = 0$, we obtain the sought identity.

□

3.4 Convention

For an arbitrary foliation \mathcal{F} on a smooth projective surface X we define the adjoint dimension, the numerical adjoint dimension and the effective threshold as the corresponding quantity for any foliation \mathcal{G} with canonical singularities birationally equivalent to \mathcal{F} .

4. Singularities

4.1 Adjoint discrepancy and ε -canonical singularities

DEFINITION 4.1. Let \mathcal{F} be a foliation on X and let $\pi : Y \rightarrow X$ be a birational morphism. Denote by \mathcal{G} the pull-back of \mathcal{F} under π . If E is an exceptional divisor of π then the adjoint discrepancy of \mathcal{F} along E is the function

$$a(\mathcal{F}, E) : [0, \infty) \longrightarrow \mathbb{R} \\ t \longmapsto \text{ord}_E(K_{\mathcal{G}} + tN_{\mathcal{G}}^* - (\pi^*K_{\mathcal{F}} + t\pi^*N_{\mathcal{F}}^*)).$$

DEFINITION 4.2. Let \mathcal{F} be a foliation on X and $\varepsilon \geq 0$ a real number. A point $x \in X$ is ε -canonical if and only if the adjoint discrepancy of \mathcal{F} along any divisor E over x satisfies $a(\mathcal{F}, E)(t) \geq 0$ for every $t \geq \varepsilon$. The foliation \mathcal{F} is said to have ε -canonical singularities if every point $x \in X$ is ε -canonical. The smallest ε for which $x \in X$ is ε -canonical will be called the canonical threshold of \mathcal{F} at x .

PROPOSITION 4.3. Let (X, \mathcal{F}) and (Y, \mathcal{G}) be two foliations on smooth projective surfaces. Assume that \mathcal{F} and \mathcal{G} are birationally equivalent. If both \mathcal{F} and \mathcal{G} have ε -canonical singularities, then for any pair of integers n, m satisfying $m/n \geq \varepsilon$ we have that

$$h^0(X, K_{\mathcal{F}}^{\otimes n} \otimes N_{\mathcal{F}}^{*\otimes m}) = h^0(Y, K_{\mathcal{G}}^{\otimes n} \otimes N_{\mathcal{G}}^{*\otimes m}).$$

In particular, if $\text{eff}(\mathcal{F}) \geq \varepsilon$ then $\text{eff}(\mathcal{F}) = \text{eff}(\mathcal{G})$.

Proof. The proof is completely analogue to the proof of Proposition 3.6. \square

REMARK 4.4. We point out that ε' -canonical singularities are ε -canonical for every $\varepsilon \geq \varepsilon'$. In particular, canonical singularities are ε -canonical singularities for every $\varepsilon \geq 0$. Also note that the canonical threshold of a log canonical singularity is at most $1/2$, i.e. log canonical singularities are ε -canonical for every $\varepsilon \geq 1/2$. This is a straightforward consequence of the simple fact that for every divisor E exceptional over X extracted on a smooth birational surface $\pi : Y \rightarrow X$ then $\text{ord}_E(K_Y - \pi^*K_X) \in \mathbb{Z}_{>0}$.

NOTATION 4.5. If $p, q \geq 1$ are relatively prime integers then we will write

$$\frac{p}{q} = [u_0, u_1, \dots, u_n] = u_0 + \frac{1}{u_1 + \frac{1}{\dots + \frac{1}{u_n}}}$$

for the continued fraction presentation of their quotient.

If $p, q \geq 1$ are relatively prime positive integers then vector fields of the form $v = px \frac{\partial}{\partial x} + qy \frac{\partial}{\partial y}$ are tangent to the fibers of rational map $(x, y) \mapsto y^p/x^q$. As a simple computation shows, see for instance [MM80, Appendix 1, Theorem 3], the minimal reduction of singularities of v coincides with the minimal resolution of indeterminacies of the rational map above by blow-ups on closed points. In particular, there exists a unique irreducible component of the exceptional divisor which is not invariant by the resulting foliation.

DEFINITION 4.6. Let $p, q \geq 1$ be relatively prime positive integers and consider the germ of foliation on $X = (\mathbb{C}^2, 0)$ defined by $v = px \frac{\partial}{\partial x} + qy \frac{\partial}{\partial y}$. Let $\pi : Y \rightarrow X$ be the minimal reduction of singularities of \mathcal{F} , let \mathcal{G} be the transformed foliation $\pi^*\mathcal{F}$, and let E be the irreducible component of the exceptional divisor which is not \mathcal{G} invariant. We will denote the order of $K_Y - \pi^*K_X$ along E by $\varphi(p, q)$.

LEMMA 4.7. Notations as in Definition 4.6. If we write $p/q = [u_0, u_1, \dots, u_n]$ then the following assertions hold true.

- (i) π is the composition of exactly $\sum_{i=0}^n u_i$ blow-ups; and
- (ii) the order of $K_Y - \pi^*K_X$ along E satisfies $\varphi(p, q) \geq \sum_{i=0}^n u_i$.

Proof. The key observation is that the reduction of singularities of v follows step-by-step Euclid's algorithm for the computation of $\gcd(p, q)$.

Assume that $p \geq q$ and write p/q as a continued fraction $[u_0, u_1, \dots, u_n]$. The proof will be by induction on the number $N = \sum_{i=1}^n u_i$.

If $p = q = 1$ then clearly $N = 1$ and the result is obvious in this case. Assume $p > q$ and consider the blow-up $s: Z \rightarrow X$ of the origin with exceptional divisor E_0 . Over the exceptional divisor we will find two singularities with eigenvalues $(p - q, q)$ and $(p, q - p)$. Since we are assuming that $p > q$ then the pair $(p, q - p)$ corresponds to a canonical singularity while the pair $(p - q, q)$ corresponds to a non-canonical singularity. Observe that

$$\frac{p - q}{q} = [u_0 - 1, u_1, \dots, u_n]$$

Assuming that the result is true for $N - 1$ then the first part of the statement follows.

To verify item (2), notice that $K_Z = s^*K_X + E_0$. If $r: Y \rightarrow Z$ is the minimal desingularization of $s^*\mathcal{F}$ then by induction hypothesis $\text{ord}_E(K_Y - r^*K_Z) \geq N - 1$. Since $\pi = s \circ r$, we can write

$$\begin{aligned} \text{ord}_E(K_Y - \pi^*K_X) &= \text{ord}_E(K_Y - r^*(K_Z - E_0)) = \\ &= \text{ord}_E(K_Y - r^*K_Z) + \text{ord}_E(r^*E_0) \geq N. \end{aligned}$$

Then the Lemma follows by induction. \square

REMARK 4.8. *The inequality in part (2) of the Lemma becomes an equality only for singularities with eigenvalues of the form $(1, q)$. If p and q are both strictly greater than one, at some intermediate step we will be forced to blow-up at the intersection of two exceptional divisors and one will get a greater order at the end. For instance, if $p/q = [u_0, u_1]$ then order of $K_Y - \pi^*K_X$ along the last exceptional divisor is $\varphi(p, q) = (u_1 + 1)u_0 - 1$.*

As a consequence of the above description we are able to characterize ε -canonical singularities for small values of $\varepsilon > 0$.

PROPOSITION 4.9. *Let \mathcal{F} be a germ of foliation on $(\mathbb{C}^2, 0)$. If the canonical threshold of \mathcal{F} at 0 is strictly less than $1/4$ then 0 is a log-canonical singularity.*

Proof. Let v be a generator of $T_{\mathcal{F}}$. Assume first that the linear part of v is zero. If $\pi: Y \rightarrow (\mathbb{C}^2, 0)$ is the blow-up of the origin, $\mathcal{G} = \pi^*\mathcal{F}$ and E is the exceptional divisor then $K_{\mathcal{G}} = \pi^*K_{\mathcal{F}} - aE$, where $a \geq 1$. On the other hand $N_{\mathcal{G}}^* = \pi^*N_{\mathcal{F}}^* + (a + 1)E$. Therefore, if $\varepsilon < 1/2$ then the origin is not ε -canonical.

Assume now that the linear part of v is non-zero but nilpotent. We will use the description of the resolution process of this kind of singularities presented in [Bru04, Chapter 1, proof of Theorem 1]. If we blow-up the origin then we obtain only one singularity over the exceptional divisor which is invariant by the transformed foliation. This new singularity can have zero linear part or non-zero but nilpotent linear part. Let us analyze the two possibilities. Start with the case where the linear part is zero and let $\pi: Y \rightarrow (\mathbb{C}^2, 0)$ be the composition of the two obvious blow-ups. As before we will set $\mathcal{G} = \pi^*\mathcal{F}$ and will let E_1, E_2 be the two irreducible components of the exceptional divisor of π with E_2 corresponding to the last blow-up. Notice that $K_{\mathcal{G}} = \pi^*K_{\mathcal{F}} - aE_2$ for some $a \geq 1$ and $N_{\mathcal{G}}^* = \pi^*N_{\mathcal{F}}^* + E_1 + (a + 2)E_2$. Hence if $\varepsilon < 1/3$ then 0 is not an ε -canonical singularity. Let us now deal with the second possibility. If the blow-up of a nilpotent singularity with non-zero linear part leads to another nilpotent singularity with non-zero linear part then one further blow-up gives rise to a singularity with trivial linear part. Let now

$\pi : Y \rightarrow (\mathbb{C}^2, 0)$ be the composition of the three obvious blow-ups, and let E_1, E_2, E_3 be the irreducible components of the exceptional divisor numbered according to the order of appearance. If we set $\mathcal{G} = \pi^* \mathcal{F}$ then $K_{\mathcal{G}} = \pi^* K_{\mathcal{F}} - aE_3$ for some $a \geq 1$ and $N_{\mathcal{G}}^* = \pi^* N_{\mathcal{F}}^* + E_1 + 2E_2 + (a+3)E_3$. Thus if $\varepsilon < 1/4$ then 0 is not a ε -canonical singularity.

Therefore if $\varepsilon < 1/4$ then the linear part of v is non-nilpotent and we can apply [McQ08, Fact I.1.8] to conclude that 0 is a log-canonical singularity of \mathcal{F} . \square

COROLLARY 4.10. *Let \mathcal{F} be a germ of foliation on $(\mathbb{C}^2, 0)$ defined by a germ of vector field v . If $0 < \varepsilon < 1/4$ then 0 is a ε -canonical singularity of \mathcal{F} if and only if the linear part of v is non-nilpotent and one of the following holds:*

- (i) *the singularity of v is canonical; or*
- (ii) *the singularity of v is not canonical, v is analytically conjugated to $px \frac{\partial}{\partial x} + qy \frac{\partial}{\partial y}$ with p, q relatively prime positive integers, and $\varphi(p, q) \geq \frac{1-\varepsilon}{\varepsilon}$.*

Proof. Proposition 4.9 implies that the linear part of v is non-nilpotent. If 0 is not a canonical singularity then by [McQ08, Fact I.1.9] we know that v is analytically conjugated to $px \frac{\partial}{\partial x} + qy \frac{\partial}{\partial y}$ for suitable relatively prime positive integers p, q . If $\pi : Y \rightarrow X = (\mathbb{C}^2, 0)$ is the minimal reduction of singularities of \mathcal{F} , E denotes the last exceptional divisor and $\mathcal{G} = \pi^* \mathcal{F}$ then $K_{\mathcal{G}} = \pi^* K_{\mathcal{F}} - E$. Therefore the adjoint discrepancy of \mathcal{F} along E is (cf. Remark 3.3)

$$\begin{aligned} a(\mathcal{F}, E)(t) &= (1-t) \operatorname{ord}_E(K_{\mathcal{G}} + \frac{t}{1-t} K_Y - \pi^*(K_{\mathcal{F}} + \frac{t}{1-t} K_X)) = \\ &= (1-t)(-1 + \frac{t}{1-t} \varphi(p, q)). \end{aligned}$$

Since the adjoint discrepancy is clearly non-negative along all the other divisors in the minimal resolution it follows that 0 is an ε -canonical singularity if and only if $\varphi(p, q) \geq \frac{1-\varepsilon}{\varepsilon}$. \square

4.2 Example: log canonical foliations on the projective plane

For a foliation \mathcal{F} on the projective plane with log-canonical singularities one can easily verify the following assertions.

- (i) If $d = \deg(\mathcal{F}) \geq 4$ then $\operatorname{eff}(\mathcal{F}) = \frac{d-1}{d+2}$.
- (ii) If $d = \deg(\mathcal{F}) = 3$ then $\operatorname{eff}(\mathcal{F}) = 2/5$ unless \mathcal{F} has radial singularities.
- (iii) If $d = \deg(\mathcal{F}) = 2$ then $\operatorname{eff}(\mathcal{F}) = 1/4$ unless \mathcal{F} has radial singularities or dicritical singularities of type $(1, 2)$.

One could try to pursue a case-by-case analysis in order to provide an explicit lower bound for the positive effective thresholds of foliations of degree two and three with log-canonical singularities. We will show later in Section 7 that the positive effective thresholds of foliations varying in an algebraic family do not accumulate at zero. Unfortunately, our proof is not effective and, a priori, the bound might depend on the family.

5. Non-isotrivial fibrations

5.1 Producing sections

Our original motivation to introduce and study the adjoint dimension of foliations lies on our poor understanding of the linear systems $|K_{\mathcal{F}}^{\otimes n}|$. When \mathcal{F} is a foliation of general type we are not aware of lower bounds on n such that $|K_{\mathcal{F}}^{\otimes n}|$ is not empty. For the linear systems $|K_{\mathcal{F}}^{\otimes n} \otimes K_X^{\otimes m}|$ the situation

is considerably better. We can apply the current knowledge on adjoint linear systems to obtain effective bounds on n, m such that $|K_{\mathcal{F}}^{\otimes m} \otimes K_X^{\otimes n}|$ defines a rational map with two dimensional image.

PROPOSITION 5.1. *Let \mathcal{F} be a foliation with canonical singularities on a smooth projective surface. If $\text{kod}(\mathcal{F}) = 2$ then the linear system $|K_X + 4i(\mathcal{F})K_{\mathcal{F}}|$ defines a rational map with two dimensional image.*

Proof. This is an immediate consequence of [Rei88, Cor. 2]. \square

The proposition above is certainly not optimal. The real question underlying the whole issue here is whether or not one can provide universal bounds which do not depend on the index of the foliation. The reader will find a more precise formulation of this question in Problem 6.8.

5.2 Bound for the index of hyperbolic fibrations

In order to use the results above to provide explicit bounds for the degree of leaves of non-isotrivial hyperbolic fibrations we need to obtain bounds for the index of the foliation.

LEMMA 5.2. *Let \mathcal{F} be a relatively minimal foliation on a smooth projective surface X . Assume $K_{\mathcal{F}}$ is defined by a fibration $f : X \rightarrow C$ and that the general fiber of f has genus at least two. If T is an irreducible curve invariant by \mathcal{F} which intersects the support of the negative part of $K_{\mathcal{F}}$ and it is not contained in it (i.e. T is a tail) then one of the following holds:*

- (i) $P \cdot T = 0$ and T intersects exactly two connected components of the support of N , both of them of order 2; or
- (ii) $P \cdot T \geq \frac{1}{42}$.

Proof. It follows from Lemma 2.14 that

$$P \cdot T = K_{\mathcal{F}} \cdot T - \sum_{i=1}^k \frac{1}{o_i} = -\chi(T) + s + k - \sum_{i=1}^k \frac{1}{o_i} \quad (5.1)$$

where s is the number of singularities of \mathcal{F} on T which are not contained in the support of N , [Bru04, Chapter 2, Prop. 3].

Assume $P \cdot T = 0$. If $s = 0$ then we have the following possibilities for k and $o = (o_1, \dots, o_k)$: $k = 3$ and $o = (2, 4, 4)$; or $k = 3$ and $o = (3, 3, 3)$; or $k = 3$ and $o = (2, 3, 6)$; or $k = 4$ and $o = (2, 2, 2, 2)$. In all cases the whole fiber F containing T is the union of k Hirzebruch-Jung strings joined by a single common tail T and $\chi(F) = \chi_{\text{orb}}(\tilde{T}) = 0$. Since $\chi(F) < 0$ by assumption, we get that $P \cdot T > 0$ contradicting our assumption. The only remaining possibility is $s = 1$, $k = 2$ and $o = (2, 2)$. Item (1) follows.

If $P \cdot T > 0$ then it is an elementary and well known fact that the lower bound for (5.1) is equal to $1/42$ and is attained by $s = 0$, $k = 3$, and $o = (2, 3, 7)$. \square

PROPOSITION 5.3. *Let \mathcal{F} be a relatively minimal foliation on a smooth projective surface X . Assume \mathcal{F} is defined by a fibration $f : X \rightarrow C$ and that the general fiber of f has genus $g \geq 2$. Then*

$$i(\mathcal{F}) \leq (42(2g - 2))!.$$

Proof. Let $F = \sum m_i C_i$ be a fiber of f and let $K_{\mathcal{F}} = P + N$ be the Zariski decomposition of $K_{\mathcal{F}}$. If C_i is a tail then, according to Lemma 5.2, either the Hirzebruch-Jung strings intersecting it have order two or $P \cdot C_i \geq 1/42$. In the later case we get that $m_i \leq 42(2g - 2)$ since $P \cdot F = K_{\mathcal{F}} \cdot F = -\chi(F) = 2g - 2$. Moreover, Lemma 2.14 item (b) implies that the least common multiple of the orders of the Hirzebruch-Jung strings intersecting C_i divides $m_i \leq 42(2g - 2)$. The Proposition is then proved. \square

5.3 Boundedness of fibers of non-isotrivial fibrations of a given genus

Theorem A of the Introduction will follow rather easily from the more general result below.

THEOREM 5.4. *Let \mathcal{F} be a foliation with canonical singularities on a smooth projective surface X . Suppose that \mathcal{F} is a fibration with general fiber F of genus g . If $\text{kod}(\mathcal{F}) = 2$ then for every nef divisor H we have*

$$F \cdot H \leq M(K_X + 4i(\mathcal{F})K_{\mathcal{F}}) \cdot H,$$

where $M = M(g)$ satisfies the following inequality

$$M \leq 2(4i(\mathcal{F}) + 1)(2g - 2) \leq (4(42(2g - 2))! + 1)(4g - 4).$$

Proof. When \mathcal{F} is induced by a fibration and $\text{kod}(\mathcal{F}) = 2$ then the fibration is non isotrivial and the genus of the general fibre is ≥ 2 , see [Ser92, Theorem 2.1] and the discussion before that theorem.

For simplicity we will write $L = K_X + (4i(\mathcal{F}))K_{\mathcal{F}}$. Let F denote a general leaf of \mathcal{F} . If $m \geq 1$ is an integer then $mL|_F = m(4i(\mathcal{F}) + 1)K_F$. On the one hand, Riemann-Roch Theorem implies that

$$h^0(F, \mathcal{O}_F(mL|_F)) = m(4i(\mathcal{F}) + 1)(2g - 2) - g + 1.$$

On the other hand, since according to Proposition 5.1 the linear system $|L|$ defines a rational map with two dimensional image, $h^0(X, \mathcal{O}_X(mL)) \geq \binom{m+2}{2}$.

If we take $m = 2(4i(\mathcal{F}) + 1)(2g(F) - 2)$ then

$$\begin{aligned} h^0(X, \mathcal{O}_X(mL - F)) &\geq h^0(X, \mathcal{O}_X(mL)) - h^0(F, \mathcal{O}_F(m(4i(\mathcal{F}) + 1)K_F)) \geq \\ &\geq \binom{m+2}{2} - m(4i(\mathcal{F}) + 1)(2g - 2) + g - 1 = g. \end{aligned}$$

In particular, $|mL - F|$ is non-empty. As H is an arbitrary nef divisor on X then $(mL - F) \cdot H \geq 0$, which concludes the first part of the proof. The inequality in the last part of the statement is then just a consequence of Proposition 5.3. \square

5.4 Proof of Theorem A

Let \mathcal{F} be a foliation on \mathbb{P}^2 birationally equivalent to a non-isotrivial fibration of genus $g \geq 2$. Notice that its canonical bundle is isomorphic to $\mathcal{O}_{\mathbb{P}^2}(\deg(\mathcal{F}) - 1)$. Let $\pi : X \rightarrow \mathbb{P}^2$ be a birational morphism such that all the singularities of $\mathcal{G} = \pi^*\mathcal{F}$ are canonical and also such that on this model the foliation is induced by an actual fibration, that is, there exists a morphism $g : X \rightarrow \mathbb{P}^1$ and $\mathcal{G} = \ker dg$.

If we take $H = \pi^*\mathcal{O}_{\mathbb{P}^2}(1)$ then the degree of an algebraic leaf L of \mathcal{F} is given by

$$\deg(L) = H \cdot \pi^*L = H \cdot \hat{L},$$

where \hat{L} is the strict transform of L . We can thus apply Theorem 5.4 and Proposition 5.3 to deduce that

$$\begin{aligned} \deg(L) &\leq (4(42(2g - 2))! + 1)(4g - 4)(K_X + 4i(\mathcal{G})K_{\mathcal{G}}) \cdot H \\ &= (4(42(2g - 2))! + 1)(4g - 4)(K_{\mathbb{P}^2} + 4i(\mathcal{G})K_{\mathcal{F}}) \cdot \pi_*H \\ &= (4(42(2g - 2))! + 1)(4g - 4)(-3 + 4(42(2g - 2))!(\deg(\mathcal{F}) - 1)) \\ &\leq (4(42(2g - 2))!)^2 (4g - 4) \deg(\mathcal{F}), \end{aligned}$$

This concludes the proof of Theorem A. \square

5.5 Log canonical foliations on \mathbb{P}^2 of high degree

The bounds appearing in Theorem 5.4 are ridiculously large and far from optimal. Proposition 5.5 below combined with the results presented in Section 7 (notably Theorem 7.6) indicates that the dependence of

M on g in Theorem 5.4 should be at worst linear on g . The results of [McQ15] also indicate the existence of such linear bounds which are not universal but depend on the family of foliations in question.

PROPOSITION 5.5. *Let \mathcal{F} be a foliation with canonical singularities on a projective surface X . Assume that \mathcal{F} is a fibration with general fiber F of geometric genus $g \geq 2$ and that $H^0(X, K_{\mathcal{F}}^{\otimes a} \otimes N_{\mathcal{F}}^{*\otimes b})$ admits three algebraically independent sections for some $a > 0$ and $b \geq 0$. Then for every nef divisor H we have*

$$F \cdot H \leq 2a(2g - 2)(aK_{\mathcal{F}} + bN_{\mathcal{F}}^*) \cdot H.$$

Proof. For simplicity we will write $\mathcal{L} = aK_{\mathcal{F}} + bN_{\mathcal{F}}^*$. Let F denote a general leaf of \mathcal{F} . If $m \geq 1$ is an integer then $mL|_F = amK_F$. On the one hand, by Riemann-Roch Theorem

$$h^0(F, \mathcal{O}_F(mL)|_F) = ma(2g - 2) - g + 1.$$

On the other hand, our assumption on $H^0(X, \mathcal{O}_X(L))$ implies that $h^0(X, \mathcal{O}_X(mL)) \geq \binom{m+2}{2}$. If we take $m = 2a(2g - 2)$ then

$$\begin{aligned} h^0(X, \mathcal{O}_X(mL)) - h^0(F, \mathcal{O}_F(mL)|_F) &\geq \binom{2a(2g - 2) + 2}{2} - 2a^2(2g - 2)^2 + (g - 1) \\ &= 6a(g - 1) + g > 0. \end{aligned}$$

In particular, $|2a(2g - 2)L - F|$ is non-empty.

If H is an arbitrary nef divisor on X then $(2a(2g - 2)L - F) \cdot H \geq 0$, which concludes the proof. \square

In the case of foliations of the projective plane with log canonical singularities and of degree greater or equal to 5, we can actually obtain bounds that are better than linear using a simple variation of the argument used to prove Proposition 5.5.

THEOREM 5.6. *Let \mathcal{F} be a foliation on \mathbb{P}^2 of degree $d \geq 5$. Assume that \mathcal{F} has log canonical singularities and admits a rational first integral with general fiber of geometric genus $g \geq 2$. If F is a general leaf of \mathcal{F} then*

$$\deg(F) \leq \left\lceil \frac{4(2g - 2)}{(d - 4)^2} \right\rceil (d - 4).$$

Proof. Since the singularities of \mathcal{F} are ε -canonical for $\varepsilon = 1/2$ (see Remark 4.4) we have that the dimension of the vector spaces $H^0(\mathbb{P}^2, K_{\mathcal{F}}^{\otimes 2m} \otimes N_{\mathcal{F}}^{*\otimes m})$, $m > 0$ is unaltered after replacing \mathcal{F} by a model with at worst canonical singularities.

Let F be a general fiber of the rational first integral of \mathcal{F} and consider the real-valued function

$$f(m) = \binom{m(d - 4) + 2}{2} - 2m(2g - 2) + g - 1.$$

Its values on positive integers correspond to the difference $h^0(\mathbb{P}^2, K_{\mathcal{F}}^{\otimes 2m} \otimes N_{\mathcal{F}}^{*\otimes m}) - h^0(\tilde{F}, K_{\tilde{F}}^{\otimes 2m})$, where \tilde{F} is the normalization of F . Since $f(4(2g - 2)/(d - 4)^2) = (dg + 8g - 12)/(d - 4)$ which is clearly positive and moreover the derivative of f satisfies $f'(4(2g - 2)/(d - 4)^2) = (3/2)d + 4g - 10 > 0$, it follows that if m is the smallest integer greater than $4(2g - 2)/(d - 4)^2$ then there exists a section of $K_{\mathcal{F}}^{\otimes 2m} \otimes N_{\mathcal{F}}^{*\otimes m} \simeq \mathcal{O}_{\mathbb{P}^2}(m(d - 4))$ vanishing identically on F . The Theorem follows. \square

As already mentioned in the Introduction, this Theorem 5.6 refines a classical result of Poincaré, see [Hen91, pages 169 and 176] and [Per03, Chapter 7, Corollary 14].

6. Classification via adjoint dimension

In this section we apply the results recalled in Section 2 to obtain a classification of foliations on surfaces according to their adjoint dimension.

6.1 K_X -negative extremal rays

Recall that for a smooth projective surface X the K_X -negative extremal rays are spanned by numerical classes of rational curves of self-intersection either $-1, 0$ or 1 . The first case corresponds to the exceptional divisor of the blow-up of a smooth point, the second to a smooth fiber of a \mathbb{P}^1 -bundle, while the last one is just the class of a line in \mathbb{P}^2 .

LEMMA 6.1. *Let \mathcal{F} be a relatively minimal foliation with pseudoeffective $K_{\mathcal{F}}$ on a smooth projective surface X , and let $K_{\mathcal{F}} = P + N$ be the Zariski decomposition of $K_{\mathcal{F}}$. Assume there exists a K_X -negative extremal curve $C \subset X$ and $P \cdot C = 0$. Then the Kodaira dimension of \mathcal{F} is either 0 or 1. Moreover, if $\text{kod}(\mathcal{F}) = 1$, then the image of C in the canonical model $\pi : X \rightarrow Z$ of \mathcal{F} is proportional to $\pi_* K_{\mathcal{F}}$.*

Proof. If C is an extremal ray with $C^2 \geq 1$ then Hodge index theorem implies that P is numerically zero. Theorem 2.18 implies $\text{kod}(\mathcal{F}) = 0$.

If instead $C^2 = 0$ then P is numerically equivalent to a non-negative multiple of C and we deduce that either $\nu(\mathcal{F}) = 0$ or $\nu(\mathcal{F}) = 1$. The case $\nu(\mathcal{F}) = 0$ follows as before. If $\nu(\mathcal{F}) = 1$ and since P is numerically equivalent to an effective divisor, we can apply Theorem 2.20 and Lemma 2.21 to deduce that $\text{kod}(\mathcal{F}) = 1$.

From now on assume that $C^2 = -1$ and let $\pi : X \rightarrow Y$ be the contraction of \mathcal{F} into its canonical model. If C is not contracted by π then write $\pi^* \pi_* C = C + \sum a_i E_i$ where $a_i > 0$ and the E_i are π -exceptional divisors. Thus $\pi_* P \cdot \pi_* C = P \cdot \pi^* \pi_* C = P \cdot C$ since P is the pull-back of a nef divisor from Y and hence π -exceptional curves intersect P trivially. As we are assuming $P \cdot C = 0$ we deduce from Hodge index Theorem that either P is numerically trivial, or that $\pi_* C^2 = 0$ and $\pi_* P$ is numerically equivalent to a positive multiple of $\pi_* C$. Hence $\nu(\mathcal{F}) \in \{0, 1\}$. As before, we obtain that in both cases $\nu(\mathcal{F}) = \text{kod}(\mathcal{F})$.

Suppose now that C is contracted by π . In this case C is \mathcal{F} -invariant according to Theorem 2.17. Since $C^2 = -1$ and \mathcal{F} is relatively minimal we have that $Z(\mathcal{F}, C) \geq 3$. Notice that $K_{\mathcal{F}} \cdot C = -2 + Z(\mathcal{F}, C)$ and, as we are assuming $P \cdot C = 0$, according to Lemma 2.14 we also have that $K_{\mathcal{F}} \cdot C = \sum_{i=1}^k 1/o_i$ where o_i are the orders of the Hirzebruch-Jung strings intersecting C . Then we must have $k = 2$ and $o_1 = o_2 = 2$; or $k = 3$ and $(o_1, o_2, o_3) \in \{(2, 3, 6), (3, 3, 3)\}$; or $k = 4$ and $(o_1, o_2, o_3, o_4) = (2, 2, 2, 2)$. If we contract the Hirzebruch-Jung strings intersecting C , we obtain that the direct image of C has self-intersection ≥ 0 , cf. [McQ08, Remark III.2.2]. Thus C cannot be contracted by π contrary to our assumption. \square

6.2 Kodaira dimension zero

LEMMA 6.2. *Let \mathcal{F} be a relatively minimal foliation with pseudoeffective $K_{\mathcal{F}}$ on a smooth projective surface X . If $\pi : X \rightarrow Z$ is the contraction of the negative part of $K_{\mathcal{F}}$ (i.e. $\pi_* \mathcal{F}$ is a nef model of \mathcal{F}) and we write $K_X + \Delta = \pi^* K_Z$ then $i(\mathcal{F})N - \Delta$ is effective.*

Proof. If E_1, \dots, E_k are the exceptional divisors of π then Δ is defined by the relations

$$\Delta \cdot E_i = -K_X \cdot E_i = 2 + E_i^2.$$

Notice that $2 + E_i^2 \leq 0$ for every i , while $2 + E_i^2 \geq (E_1 + \dots + E_k) \cdot E_i$ for every i and the latter inequality is strict when E_i is either a handle or a tail in a Hirzebruch-Jung string. Therefore by [KM98,

Corollary 4.2] the coefficients of Δ lie in $[0, 1)$. Since N is effective the lemma follows. \square

PROPOSITION 6.3. *Let \mathcal{F} be a relatively minimal foliation of Kodaira dimension zero on a smooth projective surface X . If $\pi : X \rightarrow Z$ is the contraction of the negative part of the Zariski decomposition of $K_{\mathcal{F}}$ and (X, Δ) is the pair satisfying $K_X + \Delta = \pi^* K_Z$ then the adjoint dimension and the numerical adjoint dimension of \mathcal{F} coincide with the Kodaira dimension of (X, Δ) . Moreover, when $\text{adj}(\mathcal{F}) \geq 0$ then $\text{eff}(\mathcal{F}) \geq \frac{1}{i(\mathcal{F})+1} \geq \frac{1}{13}$.*

Proof. Let $K_{\mathcal{F}} = P + N$ be the Zariski decomposition of $K_{\mathcal{F}}$. Since we are assuming that \mathcal{F} has Kodaira dimension zero we have that $P \sim_{\mathbb{Q}} 0$. Let $\pi : X \rightarrow Z$ be the contraction of the support of N and notice that we can write

$$K_{\mathcal{F}} + \varepsilon K_X = \varepsilon \pi^* K_Z + (N - \varepsilon \Delta).$$

Assume that ε is rational and satisfies $\varepsilon < 1/i(\mathcal{F})$. Lemma 6.2 implies that $(N - \varepsilon \Delta)$ is effective. Hence for any k sufficiently divisible, $h^0(X, k(\varepsilon \pi^* K_Z + (N - \varepsilon \Delta))) \geq h^0(X, k\varepsilon \pi^* K_Z) = h^0(Z, k\varepsilon K_Z)$. Since every irreducible component E of the support of $(N - \varepsilon \Delta)$ is π -exceptional we also have the opposite inequality. This shows that the Kodaira dimension of Z is equal to the adjoint dimension of \mathcal{F} .

To verify that the adjoint dimension and the numerical adjoint dimension of \mathcal{F} coincide first observe that every irreducible component E of $N - \varepsilon \Delta$ satisfies $\pi^* K_Z \cdot E = 0$. Therefore the numerical dimension of $K_{\mathcal{F}} + \varepsilon K_X$ coincides with the numerical dimension of K_Z . As the numerical dimension of K_Z and the Kodaira dimension of (X, Δ) coincide, the Proposition follows.

Finally, the last part of the statement follows from the fact that for a relatively minimal foliation \mathcal{F} of Kodaira dimension 0, $i(\mathcal{F}) \leq 12$, see (2.1). \square

6.3 Kodaira dimension one

PROPOSITION 6.4. *Let \mathcal{F} be a relatively minimal foliation of Kodaira dimension one on a smooth projective surface X . Let g be the genus of a general fiber of the Iitaka fibration of \mathcal{F} . If $g = 0$ then $\text{adj}(\mathcal{F}) = \text{adj}_{\text{num}}(\mathcal{F}) = -\infty$. Otherwise*

$$\text{adj}(\mathcal{F}) = \text{adj}_{\text{num}}(\mathcal{F}) = \min\{g, 2\} \text{ and } \text{eff}(\mathcal{F}) \geq \frac{1}{4i(\mathcal{F}) + 1}.$$

Proof. Let $f : X \rightarrow B$ be the Iitaka fibration of \mathcal{F} . Assume first that $g = 0$. Then for a general fiber F of f we have that $K_{\mathcal{F}} \cdot F = 0$ and $K_X \cdot F = -2$. Hence $K_{\mathcal{F}} + \varepsilon K_X$ is not pseudoeffective for every $\varepsilon > 0$. It follows that $\text{adj}(\mathcal{F}) = \text{adj}_{\text{num}}(\mathcal{F}) = -\infty$.

Assume now that $g \geq 1$. Let $K_{\mathcal{F}} = P + N$ be the Zariski decomposition of $K_{\mathcal{F}}$ and let $\pi : X \rightarrow Z$ be the contraction of the negative part of $K_{\mathcal{F}}$. Denote by \mathcal{G} the direct image of \mathcal{F} . We claim that $4i(\mathcal{F})K_{\mathcal{G}} + K_Z$ is nef. Suppose not, and let D be an effective divisor such that $(4i(\mathcal{F})K_{\mathcal{G}} + K_Z) \cdot D < 0$. By the Cone Theorem we can numerically decompose D as a sum $\sum a_i C_i + R$ where R is a pseudoeffective divisor and satisfies $K_Z \cdot R \geq 0$; C_i are K_Z -negative extremal rays satisfying $0 < -K_Z \cdot C_i \leq 4$ and $a_i \in \mathbb{R}_{>0}$. Therefore, there exists a K_Z -negative extremal ray C such that $(4i(\mathcal{F})K_{\mathcal{G}} + K_Z) \cdot C < 0$. If $K_{\mathcal{G}} \cdot C = 0$ then Lemma 6.1 implies that C is numerically proportional to $K_{\mathcal{G}}$. Infact, denoting by \tilde{C} the strict transform of C on X , we have that $P \cdot \tilde{C} = \pi^{K_{\mathcal{G}}} \cdot \tilde{C} = 0$. Consequently C is proportional to a general fiber of $f \circ \pi^{-1}$ and must intersect K_Z non-negatively. Thus $K_{\mathcal{G}} \cdot C > 0$. Since $i(F)K_{\mathcal{G}}$ is Cartier we deduce that $4i(\mathcal{F})K_{\mathcal{G}} \cdot C \geq 4$. It follows that also in this case $(4i(\mathcal{F})K_{\mathcal{G}} + K_Z) \cdot C \geq 0$. We conclude that $4i(\mathcal{F})K_{\mathcal{G}} + K_Z$ is nef. Consequently we obtain that

$$K_{\mathcal{F}} + \frac{1}{4i(\mathcal{F})} K_X = \pi^* \left(K_{\mathcal{G}} + \frac{1}{4i(\mathcal{F})} K_Z \right) + \left(N - \frac{1}{4i(\mathcal{F})} \Delta \right) \quad (6.1)$$

where Δ is defined by $K_X + \Delta = \pi^* K_Z$. Since the singularities of Z are klt, it follows that $N - \frac{1}{4i(\mathcal{F})}\Delta$ is effective and that $K_{\mathcal{F}} + \frac{1}{4i(\mathcal{F})}K_X$ is pseudoeffective. Thus $\text{eff}(\mathcal{F}) \geq \frac{1}{4i(\mathcal{F})+1}$.

It remains to determine the adjoint dimension of \mathcal{F} . For that, notice that (6.1) is the Zariski decomposition of $K_{\mathcal{F}} + \frac{1}{4i(\mathcal{F})}K_X$. When $g = 1$, since K_X is trivial when restricted to the general fiber of f it follows that the positive part $\pi^* \left(K_G + \frac{1}{4i(\mathcal{F})}K_Z \right)$ is numerically proportional to a general fiber and also that there exists an effective \mathbb{Q} -divisor D on B such that $\pi^* \left(K_G + \frac{1}{4i(\mathcal{F})}K_Z \right) = f^*B$. Hence $\text{adj}_{\text{num}}(\mathcal{F}) = \text{adj}(\mathcal{F}) = 1$.

To prove the claim for $g \geq 2$ it suffices to verify that $\pi^* (K_G + \varepsilon K_Z)^2 > 0$ for ε sufficiently small. If this were not the case then $K_G \cdot K_Z = 0$ and $K_Z \cdot K_Z = 0$. The Hodge index theorem would imply that $\pi^* K_Z$ is proportional to a general fiber f . But this is not possible since $\pi^* K_Z \cdot F = 2g - 2 > 0$ for any fiber F of f . \square

6.4 Kodaira dimension two and non-abundant foliations

LEMMA 6.5. *Let \mathcal{F} be a relatively minimal foliation with canonical singularities which is not a fibration by rational curves. Let $K_{\mathcal{F}} = P + N$ be the Zariski decomposition of $K_{\mathcal{F}}$. If $\text{kod}(\mathcal{F}) \notin \{0, 1\}$ then $P + \frac{1}{3i(\mathcal{F})}K_X$ is nef.*

Proof. Aiming at a contradiction, let C be a curve such that $(P + 1/3i(\mathcal{F})K_X) \cdot C < 0$. As in the proof of Proposition 6.3 we can assume that C is a K_X -negative extremal curve and therefore $K_X \cdot C \in \{-3, -2, -1\}$. By Lemma 6.1, $P \cdot C > 0$. Hence

$$-K_X \cdot C_i > 3i(\mathcal{F})(P \cdot C_i) \geq 3$$

gives the sought contradiction. \square

PROPOSITION 6.6. *Let \mathcal{F} be a relatively minimal foliation with canonical singularities and pseudoeffective canonical bundle. If $\text{kod}(\mathcal{F}) \notin \{0, 1\}$ then $\text{adj}_{\text{num}}(\mathcal{F}) = \text{adj}(\mathcal{F}) = 2$.*

Proof. Let $K_{\mathcal{F}} = P + N$ be the Zariski decomposition of $K_{\mathcal{F}}$. Since $\text{kod}(\mathcal{F}) \neq 0$ we have that $\nu(\mathcal{F}) \geq 1$. Lemma 6.5 implies that $P + \varepsilon K_X$ is nef for $0 < \varepsilon$ sufficiently small.

Assume by contradiction that \mathcal{F} is not of adjoint general type. Then $(P + \varepsilon K_X)^2$ must vanish identically. In fact, if $(P + \varepsilon K_X)^2$ does not vanish identically, then for $0 < \varepsilon$ sufficiently small $(P + \varepsilon K_X)^2 > 0$ by the nefness of $P + \varepsilon K_X$. That implies that $P + \varepsilon K_X$ is big $0 < \varepsilon$ sufficiently small and the same holds true for $K_{\mathcal{F}} + \varepsilon K_X$, as $N \geq 0$, reaching a contradiction. From this observation, it follows that $P^2 = P \cdot K_X = K_X^2 = 0$. Lemma 2.21 implies that $\text{kod}(\mathcal{F}) \geq 0$. But this is not possible by the assumptions in the statement of the proposition. Hence we reach the desired contradiction and the result follows. \square

6.5 Characterization of rational fibrations (Proof of Theorem B)

One immediate consequence of the classification of foliations according to their adjoint dimension is the characterization of rational fibrations stated in the Introduction as Theorem B.

THEOREM 6.7. *Let \mathcal{F} be a foliation with canonical singularities on a smooth projective surface X . Then \mathcal{F} is a rational fibration if and only if $h^0(X, K_{\mathcal{F}}^{\otimes m} \otimes N_{\mathcal{F}}^{\otimes n}) = 0$ for every $m > 0$ and every $n \geq 0$.*

Proof. If $\text{adj}(\mathcal{F}) \geq 0$ then $h^0(X, K_{\mathcal{F}}^{\otimes m} \otimes N_{\mathcal{F}}^{\otimes n}) \neq 0$ for some $m, n > 0$ by definition. If instead $\text{adj}(\mathcal{F}) = -\infty$ and \mathcal{F} is not a fibration by rational curves then by Theorem 2.6, $K_{\mathcal{F}}$ must be pseudoeffective and $\text{kod}(\mathcal{F}) = -\infty$. Moreover, we can assume that \mathcal{F} is relatively minimal by Proposition 3.6. But then Proposition 6.6 gives a contradiction, as it would imply that $\text{adj}(\mathcal{F}) = 2$. \square

For foliations on smooth surfaces of Kodaira dimension 0 or 1, $h^0(X, K_{\mathcal{F}}^{\otimes n}) > 0$ for some n between 1 and 12, see [Per05] and [CLV16, Section 4]. It is a simple matter to obtain effective non-vanishing of $h^0(X, K_{\mathcal{F}}^{\otimes n} \otimes N_{\mathcal{F}}^{\otimes m})$ for foliations \mathcal{F} of adjoint general type as functions of their index $i(\mathcal{F})$. This is what we did in the proof of 5.1 when $\kappa(\mathcal{F}) = 2$. The real question here is if one can do that regardless of the index of the the foliation.

PROBLEM 6.8. *Find universal bounds on $(n, m) \in \mathbb{Z}_{>0} \times \mathbb{Z}_{>0}$ in order to ensure the non-vanishing of $h^0(X, K_{\mathcal{F}}^{\otimes m} \otimes N_{\mathcal{F}}^{\otimes n})$ for foliations of adjoint general type.*

For bounded families of foliations, the results of Section 7 imply the existence of bounds depending on the family.

7. Variation in moduli

7.1 Families of foliations

We start by spelling out the definition of family of foliated surfaces.

DEFINITION 7.1. *Let $\pi : \mathcal{X} \rightarrow T$ be a family of smooth projective surfaces, i.e. \mathcal{X} and T are irreducible complex manifolds and π is a proper submersion with projective surfaces as fibers. A family of foliations parametrized by T is a foliation \mathcal{F} of dimension one on \mathcal{X} which is everywhere tangent to the fibers of π . If \mathcal{X}, T, π and \mathcal{F} are all algebraic then we say that \mathcal{F} is an algebraic family of foliations.*

For the sake of notation, we will indicate with \mathcal{X}_t the (schematic) fiber of π over a point $t \in T$ and with $\mathcal{F}_t, \mathcal{F}|_{\mathcal{X}_t}$.

Notice that in the definition above we do not impose any condition on the nature of singularities of \mathcal{F} , contrary to what is done in [Bru01] where the singularities are supposed to be reduced. Also when the dimension of T is at least two it may happen that some fibers of π are contained in the singular set of \mathcal{F} .

REMARK 7.2. *Although no assumption is made on the nature of the singularities, the singular set of \mathcal{F} has codimension at least two as the singular set of any foliation on a smooth manifold. Then there exists a non empty Zariski open subset $U \subset T$ such that $\text{sing}(\mathcal{F}_t) = \text{sing}(\mathcal{F}) \cap \mathcal{X}_t$ for every $t \in U$. In particular, $K_{\mathcal{F}_t} = (K_{\mathcal{F}})|_{\mathcal{X}_t}$, for every $t \in U$. Note also that for every $t \in U$, we have the equality $\det N_{\mathcal{F}|_{\mathcal{X}_t}}^* = N_{\mathcal{F}_t}^*$.*

It is useful to think of an algebraic family of foliations parametrized by T as a foliation defined over the function field $\mathbb{C}(T)$. Algebraic properties of a very general member \mathcal{F}_t of the family – like existence of invariant algebraic curves, rational first integrals, transversely projective structures – are displayed already when one considers the foliation as defined over $\mathbb{C}(T)$. Also the Kodaira dimension (resp. the adjoint dimension) of the foliation defined over $\mathbb{C}(T)$ coincides with the Kodaira dimension (adjoint dimension) of a very general member of the family.

7.2 Partial reduction of singularities for families

When the singularities of a family of foliations \mathcal{F} on smooth surfaces are reduced, Brunella [Bru01] showed that the Kodaira dimension of the foliations is constant in the family. Later, Cascini and Floris proved that for families of foliations with reduced singularities and for m sufficiently large, the m -th plurigenera $h^0(\mathcal{X}_t, mK_{\mathcal{F}_t})$ is constant. They also presented examples of families of foliations with reduced singularities which have non constant m -th plurigenera for small values of m , see [CF15, Section 3.4].

When we do not restrict to families of foliations with reduced singularities, then one of the sources of difficulties in applying birational techniques similar to those of [CF15] to understand the behavior of the plurigenera comes from the fact that canonical singularities are not stable in the Zariski topology: the set of foliations with at worst canonical singularities can fail to be Zariski open as the family of foliations on \mathbb{C}^2 parametrized by \mathbb{C} and defined by $x dy - t y dx$ shows. In this family the singularity at the origin is canonical if and only if $t \notin \mathbb{Q}_+$. Thus a very general foliation in the family has canonical singularities, but the set of foliations with non-canonical singularities is Zariski dense. This unpleasant situation can be avoided if instead one considers ε -canonical singularities for $\varepsilon > 0$.

LEMMA 7.3. *Let \mathcal{F} be an algebraic family of foliations parametrized by an algebraic variety T . If $0 < \varepsilon < 1/4$ then the subset of T corresponding to foliations with isolated and ε -canonical singularities is a Zariski open subset of T .*

Proof. This is a simple consequence of Corollary 4.10. If a singularity is not ε -canonical, $0 < \varepsilon < 1/4$, then either its linear part is nilpotent or the singularity is formally equivalent to one of the finitely many singularities of the form $p x \frac{\partial}{\partial x} + q y \frac{\partial}{\partial y}$ with p, q relatively prime positive integers satisfying $\varphi(p, q) < \frac{\varepsilon}{1-\varepsilon}$ (see Definition 4.6 for the meaning of φ). Since both conditions are clearly closed the lemma follows. \square

PROPOSITION 7.4. *Given an algebraic family of foliation \mathcal{F} parametrized by an algebraic variety T and a real number $\varepsilon > 0$, there exists a non-empty Zariski open subset $U \subset T$ and a family of foliations \mathcal{G} on $\mathcal{Y} \rightarrow U$ obtained from $\mathcal{F}|_U$ by a finite composition of blow-ups over (multi)-sections such that for every closed point $t \in U$, the foliation \mathcal{G}_t has at worst ε -canonical singularities.*

Proof. First replace T by a Zariski open subset in such a way that the singular scheme of \mathcal{F} becomes flat over T . In particular, every irreducible component of the singular set of \mathcal{F} is a multi-section of the projection to T .

We will say that an irreducible component Σ of the singular set of \mathcal{F} is reduced when, for a very general $t \in T$, the points in Σ_t are reduced singularities for \mathcal{F}_t .

If we interpret \mathcal{F} as a foliation $\mathcal{F}_{\overline{\mathbb{C}(T)}}$ over $\overline{\mathbb{C}(T)}$, the Zariski closure of the function field of T , then the reduced/non-reduced singularities of $\mathcal{F}_{\overline{\mathbb{C}(T)}}$ correspond, respectively, to irreducible components of $\text{sing}(\mathcal{F})$ having reduced/non-reduced singularities at the very general point.

The proof of Seidenberg's theorem presented in [MM80, Appendix 1] and [Bru04] works for arbitrary algebraically closed fields of characteristic zero. It consists of showing that the iterated blow-ups of non-reduced points will eventually lead to foliations with only reduced singularities. First one aims to obtain foliations with all singularities having non-trivial linear part. To achieve this one controls the multiplicity of the singularities of the foliation by means of Van den Essen formula which is established in [vdE79, pages 52-53] (see also [MM80, Appendix 1, formulas (2.1) and (2.2)]) for arbitrary algebraically closed fields of characteristic zero. Then to go from non-trivial linear part to non-nilpotent linear part, one carries out explicit algebraic computation computations, for details see [MM80, Appendix 1, Theorem 2]. Finally, to go from non-nilpotent linear part to quotient of eigenvalues not belonging to $\mathbb{Q}_{>0}$ one relies on Euclidean division algorithm [MM80, Appendix 1, Theorem 3].

Therefore blowing-up non-reduced multi-sections, restricting T to suitable Zariski open subsets, and repeating if necessary, will lead to the sought a family of foliations over a Zariski open subset of T with all irreducible components of its singular set reducible. We apply Lemma 7.3 to conclude. \square

7.3 Families of foliations of negative adjoint dimension

Foliations of negative adjoint dimension also behave better in families compared to foliations of negative Kodaira dimension.

LEMMA 7.5. *Let $(\pi : \mathcal{X} \rightarrow T, \mathcal{F})$ be an algebraic family of foliations. If for a very general closed point $t_0 \in T$ the foliation \mathcal{F}_{t_0} is reduced and has negative adjoint dimension then there exists a non-empty Zariski open subset $U \subset T$ such that for every closed point $t \in U$ the foliation \mathcal{F}_t has negative adjoint dimension.*

Proof. Assume first that for a very general point $t \in T$ the foliation \mathcal{F}_t has Kodaira dimension one. Since the adjoint dimension is negative, \mathcal{F}_t must be a Riccati foliation. It follows from [CLV16, Proposition 4.3] that for some $n \leq 42$ the linear system $|K_{\mathcal{F}_t}^{\otimes n}|$ is non-empty and defines the reference rational fibration. Moreover, the general fiber of the reference fibration intersects $K_{\mathcal{F}_t}$ trivially. After replacing T by a Zariski open subset we can assume that $\text{sing}(\mathcal{F}_t) = \text{sing}(\mathcal{F}) \cap \mathcal{X}_t$, see Remark 7.2. We can also assume that $\pi_* K_{\mathcal{F}}^{\otimes n}$ is locally free. Consider the rational map defined by pluricanonical sections

$$\mathcal{X} \dashrightarrow \mathbb{P}((\pi_* K_{\mathcal{F}}^{\otimes n})^*).$$

We can further restrict T in order to get an actual morphism. Let $P : \mathcal{X} \rightarrow B \subset \mathbb{P}((\pi_* K_{\mathcal{F}}^{\otimes n})^*)$ be such morphism. We do not claim that P has irreducible fibers, but since \mathcal{F}_t is a Riccati foliation of Kodaira dimension one for a very general t , it follows that very general fibers of P are smooth curves with rational connected components, completely transverse to \mathcal{F}_t , and intersecting $K_{\mathcal{F}}$ trivially. Hence, using for instance Ehresmann's fibration theorem, we can conclude that the same holds true for fibers of F over a Zariski open subset V of B . Let $U \subset T$ be a Zariski open subset contained in the image of V under the natural projection $B \rightarrow T$. We can apply [Bru04, Proposition 4.1] to deduce that for every $t \in U$ the foliation \mathcal{F}_t is a Riccati foliation and as such has negative adjoint dimension.

Assume now that for a very general point $t \in T$ the foliation \mathcal{F}_t has Kodaira dimension zero. We will make use of Lefschetz principle to deal with this case. Since the family \mathcal{F} is defined by finitely many equations, we can consider a finitely generated extension K of \mathbb{Q} over which everything in sight is defined. Embed K into \mathbb{C} and apply Theorem 2.18. We deduce that after restricting T to a Zariski open subset U and base changing the family \mathcal{F} through an étale covering $V \rightarrow U$ we obtain that the resulting family $\mathcal{X}' \rightarrow U$ is birationally equivalent to a finite quotient of a smooth family of foliations \mathcal{G} on $\mathcal{Z} \rightarrow V$ defined by global holomorphic vector fields. Since we are assuming that for a very general $t \in T$ the foliation has negative adjoint dimension it follows that the very general fiber of $\mathcal{Z} \rightarrow V$ is a surface of negative Kodaira dimension and the corresponding foliation is a Riccati foliation. It follows that for every $t \in U$, \mathcal{F}_t has negative adjoint dimension.

Finally, let us assume that a very general $t \in T$ the foliation \mathcal{F}_t is a rational fibration. Let $U \subset T$ be a Zariski open subset over which every irreducible component of the singular scheme of \mathcal{F} is flat. As we are assuming that \mathcal{F}_t is a foliation with reduced singularities tangent to a rational fibration for a very general $t \in T$, it follows that for a very general $t \in U$ the singularities of \mathcal{F}_t have invertible linear part and quotient of eigenvalues in $\mathbb{Q}_{<0}$. Flatness of the singular scheme of \mathcal{F} over U guarantees that the singularities of \mathcal{F}_t have invertible linear part for every $t \in U$. Continuity of the quotient of eigenvalues ensures that they are constant functions of $t \in U$. Thus, for every $t \in U$, the foliations \mathcal{F}_t have reduced singularities. It follows from [Bru01, Lemma 1] that \mathcal{F}_t is a rational fibration for every $t \in U$. \square

7.4 Boundedness of the effective threshold in families

We have now all the ingredients to prove the result mentioned at the end of Section 4.2.

THEOREM 7.6. *Let $(\pi : \mathcal{X} \rightarrow T, \mathcal{F})$ be an algebraic family of foliations. Then there exists $\delta > 0$ such that, for every $t \in T$, the following holds true: $\text{adj}(\mathcal{F}_t) = -\infty$ or $\text{eff}(\mathcal{F}_t) \geq \delta$. In other words, if $\text{eff}(\mathcal{F}_t) < \delta$ then $\text{adj}(\mathcal{F}_t) = -\infty$.*

Proof. The proof is by Noetherian induction. Therefore it suffices to show that the result is valid over a

Zariski open subset of T .

In particular, after replacing T by a Zariski open subset, we can assume that $\text{sing}(\mathcal{F}_t) = \text{sing}(\mathcal{F}) \cap \mathcal{X}_t$ for every $t \in T$.

Proposition 7.4 guarantees that there is no loss of generality in assuming that \mathcal{F}_t has canonical singularities for a very general $t \in T$.

If $\text{adj}(\mathcal{F}_t) \geq 0$ for a very general $t \in T$ then there exists $m, n > 0$ such that $h^0(\mathcal{X}_t, K_{\mathcal{F}_t}^{\otimes m} \otimes N_{\mathcal{F}_t}^*{}^{\otimes n}) > 0$ for a very general $t \in T$. Choose $\varepsilon > 0$ small enough and apply Proposition 7.4 to obtain a Zariski open $U \subset T$ such that \mathcal{F}_t has at worst ε -canonical singularities for every $t \in U$. Remark 7.2 guarantees that $K_{\mathcal{F}_t}^{\otimes m} \otimes N_{\mathcal{F}_t}^*{}^{\otimes n}$ is the restriction of a line bundle on \mathcal{X} to $\pi^{-1}(t)$. By semi-continuity it follows that $\text{eff}(\mathcal{F}_t) \geq \frac{n}{m}$ for every $t \in U$.

If instead $\text{adj}(\mathcal{F}_t) = -\infty$ for a very general $t \in T$ then Lemma 7.5 implies that the same holds true for every t in a Zariski open subset of T .

In any case, we have just proved that the result is true for the restriction of \mathcal{F} to a non-empty Zariski open subset U of T .

The proof continues by passing to irreducible components of $T \setminus U$. To be more precise let S be a resolution of singularities of one such irreducible component, and let $\mathcal{Y} \rightarrow S$ be the base to change of $\mathcal{X} \rightarrow T$ to $S \rightarrow T \setminus U \subset T$. If $v \in H^0(\mathcal{X}, T_{\mathcal{X}/T} \otimes K_{\mathcal{F}})$ is a twisted vector field defining \mathcal{F} then it induces an element $w \in H^0(\mathcal{Y}, T_{\mathcal{Y}/S} \otimes (K_{\mathcal{F}})_{\mathcal{Y}})$. If w is identically zero then the points of S do not correspond to foliations and we do nothing. Otherwise, we divide w by the divisorial components of its singular set in order to obtain the defining twisted vector field of a family of foliations over S . Since we are back to our original problem, but with a lower dimensional base, we are done by Noetherian induction. \square

8. Foliations with rational first integrals

8.1 Transversely affine and transversely projective foliations

A foliation on a projective surface X is called transversely affine if for any rational 1-form ω_0 defining \mathcal{F} , there exists a rational 1-form ω_1 such that

$$d\omega_0 = \omega_0 \wedge \omega_1 \quad \text{and} \quad d\omega_1 = 0.$$

Similarly, a foliation \mathcal{F} on X is called transversely projective if for any rational 1-form ω_0 defining \mathcal{F} there exists rational 1-forms ω_1 and ω_2 such that

$$d\omega_0 = \omega_0 \wedge \omega_1$$

$$d\omega_1 = 2\omega_0 \wedge \omega_2$$

$$d\omega_2 = \omega_1 \wedge \omega_2.$$

For a thorough discussion about transversely affine and transversely projective foliations of codimension one on projective manifolds the reader should consult [CP14] and [LPT16] respectively.

8.2 Statement of the main result

This section is devoted to the proof of the following result.

THEOREM 8.1. *Let $(\pi : \mathcal{X} \rightarrow T, \mathcal{F})$ be an algebraic family of foliations and $g \geq 0$ be an integer. Let $\Sigma_g \subset T$ be the Zariski closure of the set of parameters corresponding to foliations birationally equivalent to a fibration of geometric genus at most g . Then for every $t \in \Sigma_g$ the foliation \mathcal{F}_t is transversely projective.*

If one considers the universal family of degree d foliations on \mathbb{P}^2 then one promptly realizes that Theorem C is nothing but a particular case of this more general statement.

8.3 Example

Before dealing with the proof of Theorem 8.1 let us analyze the Zariski closure of the set of foliations admitting a rational first integral in a family derived from Gauss hypergeometric equation.

Whenever $c \notin \mathbb{Z}$, Gauss hypergeometric equation

$$z(1-z)w'' + (c - (a+b+1)z)w' - abw = 0,$$

admits as general solution in a neighborhood of the origin the function

$$\varphi(z) = C_1 F(a, b, c; z) + C_2 z^{1-c} F(a-c+1, b-c+1, 2-c; z),$$

where C_1, C_2 are arbitrary constants to be determined by boundary conditions and

$$F(a, b, c; z) = 1 + \sum \frac{(a)_n (b)_n}{(c)_n} z^n, \quad (p)_n := p(p+1)(p+2) \cdots (p+n-1).$$

The change of variable $y(z) = -d \log w(z)$ associates a Riccati equation/foliation to any second order differential equation. In this new coordinate the family of foliations induced by Gauss hypergeometric equation can be written as

$$\omega = z(1-z)dy - z(1-z)y^2 + (c - (a+b+1)z)y + abdz.$$

If $\varphi(z)$ is an arbitrary solution of Gauss hypergeometric equation then $y = -d \log \varphi(z)$ is a solution of the corresponding Riccati equation. If we choose $c \in \mathbb{Q} - \mathbb{Z}$, $a \in \mathbb{Z}_{<0}$, and $b = c - 1 + \beta$ where $\beta \in \mathbb{Z}_{<0}$ then it is clear from the explicit form of the solutions that all the leaves of the foliation corresponding to this choice of parameters are algebraic. It follows that the set of foliations in this family admitting a rational integral is Zariski dense.

On the one hand, for a very generic choice of parameters, the monodromy group of Gauss hypergeometric equation is Zariski dense in $\text{Aut}(\mathbb{P}^1)$, see for instance [Yos97, Chapter 4, § 5]. On the other hand, in the case of transversely affine Riccati foliations the monodromy group is not Zariski dense (it must be a solvable subgroup of $\text{Aut}(\mathbb{P}^1)$). Therefore for a very generic choice of parameters the foliation defined by ω is transversely projective, but not transversely affine.

To conclude, we point out that for the choice of parameters made above the foliations are birationally equivalent to fibrations by rational curves. Hence one cannot hope to replace transversely projective by transversely affine in the statement of Theorem 8.1.

8.4 Non-isotrivial fibrations

We now start the proof of Theorem 8.1. We first treat the case of foliations birationally equivalent to non-isotrivial fibrations.

PROPOSITION 8.2. *Let $g \geq 1$ be a natural number and let $(\pi : \mathcal{X} \rightarrow T, \mathcal{F})$ be an algebraic family of foliations. The Zariski closure in T of the set of parameters corresponding to foliations birationally equivalent to non-isotrivial fibrations of genus at most g consists of foliations admitting rational first integrals.*

Proof. According to [CP06, Proposition 2.1] it suffices to prove that the fibers of the non-isotrivial fibrations in the family belong to a bounded family of curves.

For $g = 1$ the boundedness is clear since the fibers of non-isotrivial elliptic fibration \mathcal{F}_t are contained

in zero sets of sections of $K_{\mathcal{F}_t}^{\otimes 12}$, see for instance [CLV16, Proposition 4.2]. The boundedness of fibers of non-isotrivial fibrations of genus $g \geq 2$ is guaranteed by Theorem A. \square

8.5 Isotrivial fibrations of adjoint general type

For isotrivial fibrations of adjoint general type the situation is better when compared to non-isotrivial fibrations as there is no need to bound the genus in order to obtain boundedness of the leaves.

PROPOSITION 8.3. *Let $(\pi : \mathcal{X} \rightarrow T, \mathcal{F})$ be an algebraic family of foliations. The Zariski closure in T of the set of parameters corresponding to foliations of adjoint general type birationally equivalent to isotrivial fibrations consists of foliations admitting rational first integrals.*

Proof. If \mathcal{F} is an isotrivial fibration of adjoint general type on a projective surface X then \mathcal{F} has Kodaira dimension one and the Iitaka fibration of $K_{\mathcal{F}}$ is an isotrivial fibration of genus $g \geq 2$. According to [CLV16, Proposition 4.10] there are at least two linearly independent sections σ_1, σ_2 of $K_{\mathcal{F}}^{\otimes k}$ for some $k \leq 42$. Consider the rational map $f = (\sigma_1 : \sigma_2) : X \dashrightarrow \mathbb{P}^1$ defined by them. The foliation \mathcal{G} defined by f coincides with the foliation defined by the Iitaka fibration of $K_{\mathcal{F}}$. Its normal bundle is of the form $N_{\mathcal{G}} = f^*T_{\mathbb{P}^1} \otimes \mathcal{O}_X(-\Delta) = K_{\mathcal{F}}^{\otimes 2k} \otimes \mathcal{O}_X(-\Delta)$ where Δ is an effective divisor. Since the leaves of \mathcal{F} are contained in fibers of the Iitaka fibration of $K_{\mathcal{G}}$, we repeat the argument to obtain the existence of a $k' \leq 42$ such that the leaves of \mathcal{F} are contained in zero set of sections of $K_X^{\otimes k'} \otimes K_{\mathcal{F}}^{\otimes 2k'k} \otimes \mathcal{O}_X(-k'\Delta)$. This suffices to prove the boundedness of the leaves of foliations in a family having adjoint general type and birationally equivalent to isotrivial fibrations. \square

8.6 First integrals and transverse structures

PROPOSITION 8.4. *Let \mathcal{F} be a foliation on a projective surface X . If $\text{adj}(\mathcal{F}) < 2$ then \mathcal{F} is a transversely projective foliation. Moreover, if $\text{adj}(\mathcal{F}) \in \{0, 1\}$ then \mathcal{F} is a transversely affine foliation.*

Proof. This is a straightforward consequence of the classification. If \mathcal{F} has adjoint dimension zero then it is birationally equivalent to a finite quotient of a foliation defined by a closed rational 1-form. Since the property of being transversely affine is invariant by dominant rational maps, \mathcal{F} is transversely affine. If \mathcal{F} has adjoint dimension one then \mathcal{F} is either a fibration (and therefore clearly transversely affine) or \mathcal{F} is a turbulent foliation which is well-known to be transversely affine (see for instance [Per03, Proposition 22]). Finally if \mathcal{F} has negative adjoint dimension then it is either a fibration, a Riccati foliation, or a finite quotient of a Riccati foliation. In any case we have that \mathcal{F} is a transversely projective foliation. \square

PROPOSITION 8.5. *Let $(\pi : \mathcal{X} \rightarrow T, \mathcal{F})$ be an algebraic family of foliations. If for a very general closed point $t_0 \in T$ the foliation \mathcal{F}_{t_0} is a transversely projective foliation then for every closed point $t \in T$ the foliation \mathcal{F}_t is a transversely projective foliation. Similarly, if for a very general closed point $t_0 \in T$ the foliation \mathcal{F}_{t_0} is a transversely affine foliation then for every closed point $t \in T$ the foliation \mathcal{F}_t is a transversely affine foliation.*

Proof. We can interpret the family of foliation as a single foliation defined over the function field $\mathbb{C}(T)$. By assumption, this foliation is transversely projective. Hence there exists a triplet $(\omega_0, \omega_1, \omega_2)$ of rational differential 1-forms with coefficients in $\overline{\mathbb{C}(T)}$, the algebraic closure of $\mathbb{C}(T)$, satisfying the equations

$$d\omega_0 = \omega_0 \wedge \omega_1$$

$$d\omega_1 = 2\omega_0 \wedge \omega_2$$

$$d\omega_2 = \omega_1 \wedge \omega_2.$$

and such that ω_0 is a 1-form differential form defined over $\mathbb{C}(T)$ which defines \mathcal{F} . According to [Cas02, Lemma 3.2] we can assume that ω_1, ω_2 are also defined over $\mathbb{C}(T)$ (no need to pass to the algebraic

closure). Therefore, over \mathbb{C} , we have the equations

$$\begin{aligned} d\omega_0 \wedge d\pi &= \omega_0 \wedge \omega_1 \wedge d\pi \\ d\omega_1 \wedge d\pi &= 2\omega_0 \wedge \omega_2 \wedge d\pi \\ d\omega_2 \wedge d\pi &= \omega_1 \wedge \omega_2 \wedge d\pi. \end{aligned}$$

If $t \in T$ is such that $\pi^{-1}(t)$ is not contained in the polar set of $(\omega_i)_\infty$ for $i = 0, 1, 2$ nor in the zero set of ω_0 then the restriction of the triple $(\omega_0, \omega_1, \omega_2)$ to the fiber over t defines a (singular) projective structure for the foliation \mathcal{F}_t on $X_t = \pi^{-1}(t)$.

Let us fix $t_0 \in T$ such that $X_0 = \pi^{-1}(t_0)$ is contained in the polar set of ω_i ($i = 0, 1, 2$) or in the zero set of ω_0 and let $f \in \pi^* \mathcal{O}_{T, t_0}$ be a rational function on X_0 corresponding to a generator of the maximal ideal of \mathcal{O}_{T, t_0} . Notice that we can replace the triplet $(\omega_0, \omega_1, \omega_2)$ by $(f^k \omega_0, \omega_1, f^{-k} \omega_2)$. Thus, there is no loss of generality in assuming that $\pi^{-1}(t_0)$ is not contained in $(\omega_0)_\infty \cup (\omega_0)_0$.

For $i = 0, 1, 2$, let a_i be the order of ω_i along X_0 and set $\alpha_i = \text{Res}_{X_0} f^{-a_i} \omega_i \wedge \frac{df}{f}$. As mentioned above we will assume that $a_0 = 0$ and, therefore, α_0 is just the restriction of ω_0 to the fiber X_0 .

If a_1 is negative then, comparing the orders along X_0 of $d\omega_0 \wedge df$ and of $\omega_0 \wedge \omega_1 \wedge df$, we deduce that $\alpha_0 \wedge \alpha_1 = 0$ and we can write $\alpha_0 = g\alpha_1$ for some rational function $g \in \mathbb{C}(X_0)$. Let $G \in \mathbb{C}(\mathcal{X})$ be a rational function on \mathcal{X} extending g . According to formula (14) of [CLNL⁺07] we can replace the triplet $(\omega_0, \omega_1, \omega_2)$ by the triplet

$$(\omega_0, \omega_1 - f^{-a_1} G \omega_0, \omega_2 + f^{-a_1} G \omega_1 + f^{-2a_1} G^2 \omega_0 - f^{-a_1} dG).$$

This increases a_1 . After a finite number of changes we may assume that $a_0 = 0$ and $a_1 \geq 0$.

Finally, if a_2 is negative and $a_1 > 0$ then α_0 is closed and it is clear that \mathcal{F}_{t_0} is transversely projective. If instead $a_2 < 0$ and $a_0 = a_1 = 0$ then comparing the orders along X_0 of $d\omega_1 \wedge df$ and $\omega_0 \wedge \omega_2 \wedge df$ we deduce that $\alpha_0 \wedge \alpha_2 = 0$. Thus we can write $\alpha_2 = h\alpha_0$ for a suitable rational function $h \in \mathbb{C}(X_0)$. From the equation $d\omega_2 \wedge df = \omega_1 \wedge \omega_2 \wedge df$ we deduce that $d\alpha_2 = \alpha_1 \wedge \alpha_2$. Combining these two identities we obtain

$$d(h\alpha_0) = \alpha_1 \wedge (h\alpha_0) \implies d\alpha_0 = \left(\alpha_1 - \frac{dh}{h}\right) \wedge \alpha_0.$$

Finally, comparing this identity with $d\alpha_0 = \alpha_0 \wedge \alpha_1$ (first equation) we obtain that $d\alpha_0 = -(1/2) \frac{dh}{h} \wedge \alpha_0$. Thus \mathcal{F}_{t_0} is transversely projective also in this case. \square

8.7 Proof of Theorem 8.1 (and of Theorem C)

Let $(\pi : \mathcal{X} \rightarrow T, \mathcal{F})$ be an algebraic family of foliations and $g \geq 0$ be an integer. We want to prove that the Zariski closure of $\Sigma_g \subset T$ (subset parametrizing foliations with rational first integral of genus at most g) corresponds to transversely projective foliations.

If a very general member of the family, say \mathcal{F}_t , is not of adjoint general type then Proposition 8.4 implies that \mathcal{F}_t is transversely projective. We can apply Proposition 8.5 to conclude that every foliation in the family is also transversely projective.

If instead a very general member is of adjoint general type then we will argue as in the proof of Theorem 7.6 to obtain a non-empty Zariski open subset of T such that every foliation parametrized by this subset is of adjoint general type.

Proposition 7.4 allows us to assume the existence of a non-empty Zariski open subset $U_0 \subset T$ that for a very general $t \in U_0$, the foliation \mathcal{F}_t has canonical singularities. Since \mathbb{C} is uncountable we also know that there exists $n, m > 0$ and an open subset $U_1 \subset T$ such that for every $t \in U_1$, the linear system $|K_{\mathcal{F}_t}^{\otimes m} \otimes N_{\mathcal{F}_t}^*|^{\otimes n}|$ defines a rational map with two dimensional image. Notice that there may

exist foliations in $U_0 \cap U_1$ which are not of adjoint general type because of the presence of non-canonical singularities. To remedy this we take $\varepsilon > 0$ sufficiently small in order to obtain from Lemma 7.3 a non-empty Zariski open $U_2 \subset T$ such that \mathcal{F}_t has ε -canonical singularities. Every foliation parametrized by non-empty Zariski open $U = U_0 \cap U_1 \cap U_2$ is of adjoint general type.

Propositions 8.2 and 8.3 imply that the Zariski closure in T of $\Sigma_g \cap U$ corresponds to foliations with rational first integrals. The Theorem follows by Noetherian induction. \square

REFERENCES

- BD79 Francesco Baldassarri and Bernard Dwork, *On second order linear differential equations with algebraic solutions*, Amer. J. Math. **101** (1979), no. 1, 42–76. MR 527825 (81d:34002)
- Bea96 Arnaud Beauville, *Complex algebraic surfaces*, second ed., London Mathematical Society Student Texts, vol. 34, Cambridge University Press, Cambridge, 1996, Translated from the 1978 French original by R. Barlow, with assistance from N. I. Shepherd-Barron and M. Reid. MR 1406314
- Bru99 Marco Brunella, *Minimal models of foliated algebraic surfaces*, Bull. Soc. Math. France **127** (1999), no. 2, 289–305. MR 1708643 (2000h:32045)
- Bru01 ———, *Invariance par déformations de la dimension de Kodaira d’un feuilletage sur une surface*, Essays on geometry and related topics, Vol. 1, 2, Monogr. Enseign. Math., vol. 38, Enseignement Math., Geneva, 2001, pp. 113–132. MR 1929324
- Bru04 ———, *Birational geometry of foliations*, Publicações Matemáticas do IMPA. [IMPA Mathematical Publications], Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 2004. MR 2114696 (2005k:32037)
- Car94 Manuel M. Carnicer, *The Poincaré problem in the nondicritical case*, Ann. of Math. (2) **140** (1994), no. 2, 289–294. MR 1298714
- Cas02 Guy Casale, *Suites de Godbillon-Vey et intégrales premières*, C. R. Math. Acad. Sci. Paris **335** (2002), no. 12, 1003–1006. MR 1955577
- CC97 Antonio Campillo and Manuel M. Carnicer, *Proximity inequalities and bounds for the degree of invariant curves by foliations of $\mathbf{P}_{\mathbb{C}}^2$* , Trans. Amer. Math. Soc. **349** (1997), no. 6, 2211–2228. MR 1407696
- CF15 P. Cascini and E. Floris, *On invariance of plurigenera for foliations on surfaces*, ArXiv e-prints (2015), arXiv e-print:1502.00817v2. Accepted for publication in Crelle Journal.
- CLN91 Dominique Cerveau and Alcides Lins Neto, *Holomorphic foliations in $\mathbf{CP}(2)$ having an invariant algebraic curve*, Ann. Inst. Fourier (Grenoble) **41** (1991), no. 4, 883–903. MR 1150571 (93b:32050)
- CLNL⁺07 Dominique Cerveau, Alcides Lins-Neto, Frank Loray, Jorge Vitório Pereira, and Frédéric Touzet, *Complex codimension one singular foliations and Godbillon-Vey sequences*, Mosc. Math. J. **7** (2007), no. 1, 21–54, 166. MR 2324555 (2010c:37107)
- CLV16 Gaël Cousin, Alcides Lins Neto, and Jorge Vitório Pereira, *Toward Effective Liouvillian Integration*, ArXiv e-prints (2016).
- CP06 S. C. Coutinho and Jorge Vitório Pereira, *On the density of algebraic foliations without algebraic invariant sets*, J. Reine Angew. Math. **594** (2006), 117–135. MR 2248154 (2007i:14010)
- CP14 Gaël Cousin and Jorge Vitório Pereira, *Transversely affine foliations on projective manifolds*, Math. Res. Lett. **21** (2014), no. 5, 985–1014. MR 3294560
- EK03 Eduardo Esteves and Steven Kleiman, *Bounds on leaves of one-dimensional foliations*, Bull. Braz. Math. Soc. (N.S.) **34** (2003), no. 1, 145–169, Dedicated to the 50th anniversary of IMPA. MR 1993042
- Gra08 Jeremy J. Gray, *Linear differential equations and group theory from Riemann to Poincaré*, second ed., Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2008. MR 2414794 (2009d:34002)
- GZ97 Alexis García Zamora, *Foliations in algebraic surfaces having a rational first integral*, Publ. Mat. **41** (1997), no. 2, 357–373. MR 1485488

- Har71 Godfrey H. Hardy, *The integration of functions of a single variable*, Hafner Publishing Co., New York, 1971, Reprint of the second edition, 1916, Cambridge Tracts in Mathematics and Mathematical Physics, No. 2. MR 0349924 (50 #2417)
- Hen91 Poincaré Henri, *Sur l'intégration algébrique des équations différentielles du premier ordre et du premier degré. I.*, Rend. Circ. Mat. Palermo **5** (1891), 161–191 (French).
- Hen97 ———, *Sur l'intégration algébrique des équations différentielles du premier ordre et du premier degré. II.*, Rend. Circ. Mat. Palermo **11** (1897), 193–239 (French).
- Kle56 Felix Klein, *Lectures on the icosahedron and the solution of equations of the fifth degree*, revised ed., Dover Publications, Inc., New York, N.Y., 1956, Translated into English by George Gavin Morrice. MR 0080930
- KM98 János Kollár and Shigefumi Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original. MR 1658959 (2000b:14018)
- LN02 Alcides Lins Neto, *Some examples for the Poincaré and Painlevé problems*, Ann. Sci. École Norm. Sup. (4) **35** (2002), no. 2, 231–266. MR 1914932 (2003j:34009)
- LPT16 Frank Loray, Jorge Vitório Pereira, and Frédéric Touzet, *Representations of quasi-projective groups, flat connections and transversely projective foliations*, J. Éc. polytech. Math. **3** (2016), 263–308. MR 3522824
- McQ08 Michael McQuillan, *Canonical models of foliations*, Pure Appl. Math. Q. **4** (2008), no. 3, part 2, 877–1012. MR 2435846 (2009k:14026)
- McQ15 ———, *Curves on surfaces of mixed characteristic*, Preprints (2015).
- Men00 Luís Gustavo Mendes, *Kodaira dimension of holomorphic singular foliations*, Bol. Soc. Brasil. Mat. (N.S.) **31** (2000), no. 2, 127–143. MR 1785264 (2001i:14015)
- MM80 Jean-François Mattei and Robert Moussu, *Holonomie et intégrales premières*, Ann. Sci. École Norm. Sup. (4) **13** (1980), no. 4, 469–523. MR 608290
- Pai97 Paul Painlevé, *Leçons sur la théorie analytique des équations différentielles professées à Stockholm (septembre, octobre, novembre 1895) sur l'invitation de S. M. le Roi de Suède et de Norwège.*, Paris: A. Hermann. 19 + 6 + 589 S. 4°. lith (1897)., 1897.
- Per02 Jorge Vitório Pereira, *On the Poincaré problem for foliations of general type*, Math. Ann. **323** (2002), no. 2, 217–226. MR 1913040 (2003e:32056)
- Per03 ———, *Integrabilidade de folheações holomorfas*, Publicações Matemáticas do IMPA. [IMPA Mathematical Publications], Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 2003, 24o Colóquio Brasileiro de Matemática. [24th Brazilian Mathematics Colloquium]. MR 2029287
- Per05 ———, *On the height of foliated surfaces with vanishing Kodaira dimension*, Publ. Mat. **49** (2005), no. 2, 363–373. MR 2177633 (2006m:32034)
- Pro03 Yuri G. Prokhorov, *On the Zariski decomposition problem*, Tr. Mat. Inst. Steklova **240** (2003), no. Biratsion. Geom. Lineĭn. Sist. Konechno Porozhdennye Algebry, 43–72. MR 1993748
- Rei88 Igor Reider, *Vector bundles of rank 2 and linear systems on algebraic surfaces*, Ann. of Math. (2) **127** (1988), no. 2, 309–316. MR 932299
- Ris70 Robert H. Risch, *The solution of the problem of integration in finite terms*, Bull. Amer. Math. Soc. **76** (1970), 605–608. MR 0269635
- Sch73 Hermann A. Schwarz, *Über die jenigen fälle, in welchen die gaussische hypergeometrische reihe eine algebraische function ihres vierten elementes darstellt.*, Journal für die reine und angewandte Mathematik **75** (1873), 292–335 (ger).
- Ser92 Fernando Serrano, *Fibred surfaces and moduli*, Duke Math. J. **67** (1992), no. 2, 407–421. MR 1177313
- Tou16 Frédéric Touzet, *On the structure of codimension 1 foliations with pseudoeffective conormal bundle*, Foliation theory in algebraic geometry, Simons Symp., Springer, Cham, 2016, pp. 157–216. MR 3644247

- vdE79 Arno van den Essen, *Reduction of singularities of the differential equation $Ady = Bdx$* , Équations différentielles et systèmes de Pfaff dans le champ complexe (Sem., Inst. Rech. Math. Avancée, Strasbourg, 1975), Lecture Notes in Math., vol. 712, Springer, Berlin, 1979, pp. 44–59. MR 548142
- vdPS03 Marius van der Put and Michael F. Singer, *Galois theory of linear differential equations*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 328, Springer-Verlag, Berlin, 2003. MR 1960772
- Yos97 Masaaki Yoshida, *Hypergeometric functions, my love*, Aspects of Mathematics, E32, Friedr. Vieweg & Sohn, Braunschweig, 1997, Modular interpretations of configuration spaces. MR 1453580

Jorge Vitório Pereira jvp@impa.br
 IMPA, Estrada Dona Castorina, 110, Horto, Rio de Janeiro, Brasil

Roberto Svaldi RSvaldi@dpmms.cam.ac.uk
 DPMMS, Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge, CB3 0WB, United Kingdom