# SINGLE AND DOUBLE PION PRODUCTION IN NP COLLISIONS AT 1.25 GEV WITH HADES.

A.K. Kurilkin<sup>1\*</sup>, G. Agakishiev<sup>\*</sup>, A. Balanda<sup>&</sup>, D. Belver<sup>⊎</sup>, A. Belyaev<sup>\*</sup>, A. Blanco<sup>\$</sup>, M. Böhmer<sup>•</sup>, J. L. Boyard<sup>↑</sup>, P. Cabanelas<sup>⊎</sup>, E. Castro<sup>⊎</sup>, S. Chernenko<sup>⋆</sup>, J. Díaz<sup>⊎</sup>, A. Dybczak<sup>&</sup>, E. Epple<sup>•</sup>, L. Fabbietti<sup>•</sup>, O. Fateev<sup>\*</sup>, P. Finocchiaro<sup>%</sup>, P. Fonte<sup>\$,a</sup>, J. Friese<sup>•</sup>, I. Fröhlich<sup>∓</sup>, T. Galatyuk<sup>∓</sup>, J. A. Garzón<sup>⊎</sup>, A. Gil<sup>□</sup>, M. Golubeva°, D. González-Díaz<sup>+</sup> F. Guber $^{\circ}$ , T. Hennino $^{\circ}$ , R. Holzmann $^{+}$ , P. Huck $^{\bullet}$ , A. Ierusalimov $^{\star}$ , I. Iori $^{\otimes,c}$ , A. Ivashkin $^{\circ}$ , M. Jurkovic, B. Kämpfer, T. Karavicheva, I. Koenig, W. Koenig, B. W. Kolb, A. Kopp<sup>×</sup>, G. Korcyl<sup>&</sup>, GK Kornakov<sup>⊎</sup>, R. Kotte<sup>±</sup>, A. Kozuch<sup>&,d</sup>, A. Krása<sup>∪</sup>, F. Krizek<sup>∪</sup> R. Krücken<sup>•</sup>, H. Kuc<sup>&</sup>, W. Kühn<sup>×</sup>, A. Kugler<sup>∪</sup>, A. Kurepin<sup>°</sup>, P. Kurilkin<sup>\*</sup>, P. Khlitz<sup>±</sup>. V. Ladygin\*, J. Lamas-Valverde<sup>⊎</sup>, S. Lang<sup>+</sup>, K. Lapidus°, T. Liu•, L. Lopes<sup>\$</sup>, M. Lorenz<sup>∓</sup>, L. Maier<sup>•</sup>, A. Mangiarotti<sup>\$</sup>, J. Markert<sup>∓</sup>, V. Metag<sup>×</sup>, B. Michalska<sup>&</sup>, J. Michel<sup>∓</sup>, C. Müntz<sup>∓</sup>, L. Naumann<sup>±</sup>, Y. C. Pachmayer<sup>∓</sup>, M. Palka<sup>∓</sup>, Y. Parpottas<sup>÷</sup>, V. Pechenov<sup>+</sup>. O. Pechenova<sup>∓</sup>, J. Pietraszko<sup>∓</sup>, W. Przygoda<sup>&</sup>, B. Ramstein<sup>∩</sup>, A. Reshetin<sup>°</sup>, J. Roskoss<sup>×</sup> A. Rustamov<sup>+</sup>, A. Sadovsky<sup>o</sup>, P. Salabura<sup>&</sup>, A. Schmah<sup>•</sup>, J. Siebenson<sup>•</sup>, Yu. G. Sobolev<sup>∪</sup>. S. Spataro<sup>×,e</sup>, H. Ströbele<sup>∓</sup>, J. Stroth<sup>∓,+</sup>, C. Sturm<sup>+</sup>, M. Sudol<sup>↑</sup>, A. Tarantola<sup>∓</sup>, K. Teilab<sup>∓</sup>. P. Tlusty<sup>∪</sup>, M. Traxler<sup>+</sup>, R. Trebacz<sup>&</sup>, H. Tsertos<sup>÷</sup>, T. Vasiliev<sup>\*</sup>, V. Wagner<sup>∪</sup>, M. Weber<sup>•</sup>. J. Wüstenfeld<sup>±</sup>, S. Yurevich<sup>+</sup>, Y. Zanevsky<sup>\*</sup>

- % Istituto Nazionale di Fisica Nucleare Laboratori Nazionali del Sud, 95125 Catania, Italy \$ LIP-Laboratório de Instrumentação e Física Experimental de Partículas , 3004-516 Coimbra, Portugal
- & Smoluchowski Institute of Physics, Jagiellonian University of Cracow, 30-059 Kraków, Poland + GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany
- <sup>±</sup> Institut für Strahlenphysik, Forschungszentrum Dresden-Rossendorf, 01314 Dresden, Germany \* Joint Institute of Nuclear Research, 141980 Dubna, Russia
  - <sup>∓</sup> Institut für Kernphysik, Goethe-Universität, 60438 Frankfurt, Germany
  - × II.Physikalisches Institut, Justus Liebig Universität Giessen, 35392 Giessen, Germany

    <sup>⊗</sup> Istituto Nazionale di Fisica Nucleare, Sezione di Milano, 20133 Milano, Italy
  - ° Institute for Nuclear Research, Russian Academy of Science, 117312 Moscow, Russia
  - Physik Department E12, Technische Universität München, 85748 München, Germany

     Department of Physics, University of Cyprus, 1678 Nicosia, Cyprus
  - ∩ Institut de Physique Nucléaire (UMR 8608), CNRS/IN2P3 Université Paris Sud, F-91406 Orsay Cedex, France
- Unuclear Physics Institute, Academy of Sciences of Czech Republic, 25068 Rez, Czech Republic □ Departamento de Física de Partículas, Univ. de Santiago de Compostela, 15706 Santiago de Compostela, Spain
  - □ Instituto de Física Corpuscular, Universidad de Valencia-CSIC, 46971 Valencia, Spain

    <sup>a</sup> Also at ISEC Coimbra, Coimbra, Portugal
    - <sup>b</sup> Also at Technische Universität Dresden, 01062 Dresden, Germany
    - <sup>c</sup> Also at Dipartimento di Fisica, Università di Milano, 20133 Milano, Italy
    - <sup>d</sup> Also at Panstwowa Wyzsza Szkola Zawodowa , 33-300 Nowy Sacz, Poland
    - <sup>e</sup> Also at Dipartimento di Fisica Generale, Universitá di Torino, 10125 Torino, Italy † E-mail: akurilkin@jinr.ru

#### Abstract

The preliminary results on charged pion production in np collisions at an incident beam energy of 1.25 GeV measured with HADES are presented. The np reactions were isolated in dp collisions at 1.25 GeV/u using the Forward Wall hodoscope, which allowed to register spectator protons. The results for  $np \to pp\pi^-$ ,  $np \to np\pi^+\pi^-$  and  $np \to d\pi^+\pi^-$  channels are compared with OPE calculations. A reasonable agreement between experimental results and the predictions of the OPE+OBE model is observed.

#### 1 Introduction

Pion production in nucleon-nucleon collisions has been the subject of considerable interest in nuclear and particle physics for many years. A number of experiments have been performed, spanning the energy region from threshold to many GeV's[1, 2]. The bulk of the experimental data has come from pp collision. The pn interaction data in the low and medium energy regions are scare despite of their importance not only for understanding the NN interactions but also for the interpretation of medium-energy heavy-ion interactions. The np interactions are studied by using deuteron-proton (dp) collisions with the deuteron either as the projectile or as the target. The main reason for this situation is due to the difficulty to create pure monoenergetic neutron beams.

One of the questions which is not answered at the present time is the contribution of isoscalar (I=0) partial waves to the inelastic np collision. The neutron-proton scattering amplitude contains both isoscalar (I=0) and isovector (I=1) parts and, while the isovector part is rather well known, even the order of magnitude of the total isoscalar cross section is badly determined. Usually, this cross-section is extracted from the difference of the total cross-sections of the pion production reactions:  $np \to pp\pi^-$  and  $pp \to pp\pi^0$ .

A study of the double pion production in the NN collisions is one way to obtain information about nucleon-nucleon, pion-nucleon, pion-pion interactions and for the investigation of resonances properties. The comparison of the double-pion production from np and pp interaction can bring new constraints on the recently reported  $e^+e^-$  excess in the np reaction [3]. Also the study of single and double pion production in np collisions at different energies is important for the determination of both the energy dependence of the total np cross section and the contribution of inelastic channels to np interactions. This paper presents preliminary results for the  $np \to pp\pi^-$ ,  $np \to np\pi^+\pi^-$  and  $np \to d\pi^+\pi^-$  channels at 1.25 GeV.

# 2 High Acceptance Di-Electron Spectrometer

The High Acceptance Di-Electron Spectrometer (HADES) is an unique apparatus installed at the heavy-ion synchrotron SIS18 at GSI Darmstadt [4]. It is designed for high-resolution and high-acceptance dielectron spectroscopy in hadron-hadron, hadron-nucleus and nucleus-nucleus reactions at beam energies in the range from 1A GeV to 2A GeV. The major part of the HADES physics program focuses on in-medium properties of the light vector mesons,  $\rho$ ,  $\omega$  and  $\phi$ .

The HADES spectrometer consists of 6 identical sectors covering the full azimuthal angle and polar angles from 18° to 85° relative to beam direction. Each sector of the spectrometer contains a Ring Imaging Cherenkov Detector (RICH) operating in a magnetic field,

inner Multi Wire Drift Chambers (MDCs) in front of magnetic field, outer MDCs behind the magnetic field, Time Of Flight (TOF and TOFino) detectors and a electromagnetic cascade detector (Pre-Shower). A detailed description of the HADES spectrometer can be found in [4]. Momentum measurement for charged particles is achieved by tracking the particles in front of and behind a toroidal field generated by six superconducting coils arranged around the beam axis. A hadron-blind Ring Imaging CHerenkov detector (RICH) placed around the target region is used for electron identification, together with TOF/TOFINO and an electromagnetic pre-shower detector (Pre-Shower). Particle identification is also provided using the correlations between time-of-flight and momentum of charged pions, protons and deuterons. Forward Wall (FW) scintillator hodoscope covering the polar angle between 1° and 7° was installed lately 2007 for tagging the spectator proton in dp reactions. With technical design features, HADES can obtain data with high quality and statistical significance.




Figure 1: Cut through two sectors of the HADES spectrometer. The magnet coils are projected onto the cut plane to visualize the toroidal magnetic field. A schematic view of the quasi-free n + p reaction is shown.

# 3 dp experiment

A deuteron beam of  $10^7$  particles/s with kinetic energy of 1.25 AGeV was incident on a liquid hydrogen cell with a length of 5 cm, corresponding to a interaction length of  $\rho d=0.35$   $g/cm^2$ . Quasi-free n-p reactions were selected at the trigger level by detection of fast spectator protons from the deuterium break-up in the FW. The detection of the spectator protons by the FW allowed to suppress the contribution of quasi-free p-p reactions [5]. The FW is an array which consists of nearly 300 scintillating cells with each 2.54 cm thickness. During the dp experiment it was located 7 m downstream the target. The estimated time resolution of the FW is about 500 ps; thus the estimated momentum resolution of the detected particles (protons) is  $\sim 11\%$ .

#### 4 Results

Fig. 2 exhibits the preliminary missing mass and invariant mass spectra for the  $np \rightarrow pp\pi^-$  channel. Both spectra are not efficiency corrected. The vertical line in Fig. 2.a) at 0.03  $GeV^2/c^4$  shows the criterion on the  $M_{miss}^2$  which was applied to remove events with an additional  $\pi^0$  production. The experimental  $M_{inv}$  distribution is shown in Fig.1.b). Two possible proton combinations were taken into account. The maximum of the distribution correspond to the  $\Delta(1232)$  resonance




Figure 2: Preliminary results for  $np \rightarrow pp\pi^-$  channel. a) - the  $pp\pi^-$  missing mass spectrum and b) the  $p\pi^-$  invariant mass spectrum. The dashed arrow in a) and the solid arrow in b) correspond to the squared mass of  $\pi^0$  and the mass of  $\Delta(1232)$  resonance, respectively.

Figs. 3. and 4. exhibit preliminary spectra of  $M_{miss}^2$  and  $M_{inv}$  for the np $\to d\pi^+\pi^-$  and np $\to np\pi^+\pi^-$  channels, respectively. The spectra are not efficiency corrected. Distributions of  $M_{miss}^2$  in Fig. 3.a) and Fig. 4.a) show that channels are separated correctly. Fig. 3.b) and 3.c) correspond to the  $M_{inv}$  spectra of  $\pi^+\pi^-$  and  $d\pi$ . Fig. 4.b), 4.c), 4.d), 4.e), 4.f) correspond to the  $M_{inv}$  spectra of  $\pi^+\pi^-$ ,  $p\pi^+$ ,  $p\pi^-$ ,  $n\pi^+$ ,  $n\pi^-$ , respectively.  $M_{inv}$  spectra of  $p\pi^+$  in Fig. 4.c) show that the np reaction should follow via np $\to \Delta^{++}n\pi^- \to np\pi^+\pi^-$ .




Figure 3: Preliminary results for np $\to d\pi^+\pi^-$  channel. a) - the  $d\pi^+\pi^-$  missing mass spectra, b) - the  $\pi^+\pi^-$  invariant mass spectra and c) -  $d\pi$  invariant mass spectra.

.




Figure 4: Preliminary results for  $np \rightarrow np\pi^+\pi^-$  channel. a) the  $p\pi^+\pi^-$  - missing mass spectra, b) the  $\pi^+\pi^-$  - invariant mass spectra, c) the  $p\pi^+$  - invariant mass spectra, d) the  $p\pi^-$  - invariant mass spectra, e) the  $n\pi^+$  - invariant mass spectra, f) the  $n\pi^-$  - invariant mass spectra. The solid arrow in a) corresponds to the squared neutron mass.

#### 5 Prediction of the Pluto and OPE+OBE models

The predictions of Pluto event generator [6] for the np $\to pp\pi^-$  and np $\to d\pi^+\pi^-$  channels are presented in Fig. 5. The spectra show events which belong to the geometric acceptance of HADES. The excitation of  $\Delta(1232)$  resonance only was included for the np $\to pp\pi^-$  process. The phase space was generated for the np $\to d\pi^+\pi^-$  channel. The behavior of  $M_{inv}$  spectra is qualitatively reproduced by Pluto event generator for both channels. However the  $\Delta(1232)$  resonance is not so clearly visible in the not corrected by the efficiency experimental distribution (see Fig. 2.b) ).




Figure 5: Prediction of Pluto for the  $np \rightarrow pp\pi^-$  and  $np \rightarrow d\pi^+\pi^-$  channels. Explanations are given in the text.




Figure 6: Predictions of OPE+OBE model calculations for  $np \rightarrow np\pi^+\pi^-$  channel.

Fig. 6 exhibits the prediction of OPE+OBE (one  $\pi$ -exchange and one barion-exchange) model [7] calculations for the  $np \to np\pi^+\pi^-$  channel. This model include also the interference of the main diagrams of reggeized one  $\pi$ -exchange model (OPER) for  $np \to np\pi^+\pi^-$  reaction at the energies 1-5 GeV [7]. The behavior of  $M_{inv}$  spectra (see Fig. 4) is qualitatively reproduced by predictions of the OPE+OBE model. However the experimental data require the correction on the efficiency.

## 6 Conclusion

The preliminary results on charged pion production in np collisions at 1.25 GeV obtained with HADES are presented. A reasonable agreement of preliminary experimental results and predictions of the Pluto event generator and OPE+OBE model is observed. More detailed information will be obtained through a comparison the efficiency corrected experimental spectra and the simulation results.

The work has been supported in part by the Russian Foundation for Basis Research grant No.~10-02-00087a.

## References

- [1] F. Shimizu, et al., Nucl. Phys. A 386, 571 (1982).
- [2] T. Tsuboyama, et al., Phys. Rev. C 62, 034001 (2000).
- [3] G. Agakishiev et al., HADES Collaboration, Phys. Lett. B 690, 118-122 (2010).
- [4] G. Agakishiev et al., HADES Collaboration, Eur. Phys. J. A 40, 45-59 (2009).
- [5] K. Lapidus, et al., HADES collaboration, ar.Xiv:0904.1128 [nucl-ex].
- [6] I. Fröhlich, et al., ar. Xiv: 0708.2382v2 [nucl-ex], (2007).
- [7] A.P. Ierusalimov et al., JINR Rapid Communications, 2[35]-89, 21-26 (1989).