Computing Mixed-strategy Nash Equilibria for Zero-Sum Game

Duc V. Nguyen, Tu Khue Tran

Abstract—This paper presents a preliminary exposition on the computation of Nash Equilibria in zero-sum games (more precisely, two-person finite constant-sum games)-an important class of problems studied in Game Theory. It first gives a brief overview of the field of Game Theory, Zero-sum Game, and Nash Equilibrium. A game-theoretic formulation of Zero-sum Games is carefully developed, followed by a mathematical analysis of Zero-sum Games as linear optimization problems. The paper then reports a computational experiment comparing the performance of two popular linear programming algorithms (the Simplex Algorithm and Interior Point Method) in solving for the solution (Nash Equilibria) of these games. A short discussion of General-sum Games - natural extensions of Zero-sum Games, as well as their mathematical, and computational properties concludes the paper.

I. Introduction

A. Game Theory

AME Theory is the mathematical study of strategic interactions amongst rational decision-makers. While game-theoretic analyses have appeared since the late 1700s, it was not until the publication of von Neumann and Morgenstern's canonical 'Theory of Games and Economic Behavirors' in 1944 that Game Theory became an independent scientific field. In the 1950s, the field enjoyed an explosion of interest due to its potential applications to global nuclear strategy, which brought about numerous developments in both its theoretical understanding and practical applications. In the decades since then, Game Theory has been applied across the social sciences, evolutionary biology, and even lends itself to numerous recent advances in machine learning.

The subject of study in Game Theory are 'games' - abstract mathematical models describing an empirical process of interactive decision-making. Each 'game' is often associated with a 'solution concepts', which is essentially the purported prediction of how a game would be play. As with any other form of mathematical model, different types of game, and solution concepts are constructed so as to be accurate and nuanced enough to capture nontrivial insights into real world interactions, while being simple enough to evade unnecessary complications.

B. Zero-Sum Game

Zero-sum Game is one of the most important class of problems in Game Theory, due both to its historical significance as one of the first 'games' formally analyzed, and its theoretical ties to linear programming. In essence, a Zero-sum Game models a strategic interaction where interests of players are diametrically opposed, that is, the gains of a player is necessarily a loss of the other. Theoretically, as we shall see, a Zero-sum Game is equivalent to a linear program problem. In real life, any type of interaction where there are two self-maximizing agents competing for a fixed amount of resources, each with a finite number of available actions (hence the precise delienation two-person finite constant-sum game) can be described as a Zero-sum Game.

1

C. Nash Equilibrium

Nash Equilibrium is arguably the most important solution concept in Game Theory. Popularized by John Nash, it describes a state in which each participant's strategy is optimal given the strategies chosen by others, and no player has an incentive to unilaterally deviate from their chosen strategy. Intuitively, such a state will always be reached, and mutually maintained by all players in a game assuming that they are *self-maximizing*, and *intelligent* (details of these assumptions will be discussed later in this paper). As a result, Nash Equilibrium is a *solution* to a game in a sense that it predicts what players would do.

II. GAME-THEORETICAL FORMULATION

A. Formulating a Zero-Sum Game

In its most general game-theoretic formulation, a zerosum game is a *strategic game* that is *strictly competitive*.

Definition 1 (Strategic Game). A *strategic game* is a triple $\langle N, (S_i), (\succeq_i) \rangle$ consisting of

- a finite set N of players;
- for each player $i \in N$ a nonempty set of actions S_i (also known as the *pure strategy space*);
- for each player $i \in N$ a preference relation \succsim_i on the set of outcomes $S = \times_{j \in N} S_j$.

2 OPTIMIZATION IN GAME THEORY

The preference relation \succsim_i captures the preference of the i^{th} player toward an outcome of the game, and also our assumptions of players' behavior in a game. Popular conventions assume the set of preference relations (\succsim_i) satisfies von Neumann and Morgenstern's four axioms of rationality [1]. These conditions ensure that each preference relations \succsim_i corresponds to a utility function $u_i(a) \ge u_i(b)$ whenever $a \succsim_i b$, and that players would act consistently to maximize the expected value of μ_i . Such is referred to as the Expected Utility Maximization [1], or Rationality Assumption. Another conventional assumption is the Intelligence Assumption, which considers players as fully informed decision-makers, who are capable of inferences.

The last assumption to make a game *zero-sum* is the *strictly competitive* assumption, which essentially requires that an outcome is preferable to one of the players must be undesirable to the other.

Definition 2 (Strictly Competitive Game). A strategic game $\langle N, (S_i), (\mu_i) \rangle$ is *strictly competitive* if for any $a, b \in S$ we have $a \succsim_i b$ if and only if $b \succsim_i a$.

The strictly competitive assumption imply that the sum of the corresponding utility functions that is $\sum_i \mu_i(s_{ij}) = 0$ for arbitrary outcome s_{ij} in S. In words, this means the utility for all players always sum to zero (hence the name zero-sum) whatever outcome s_{ij} in S. With this, we are now ready to rigorously define a zero-sum game, where we would start right-away with utility functions (instead of the game-theoretic preference relations) as per popular conventions. For simplicity, we will also define this for only a game with two players. Generalization of this two-person version to multiplayer is trivial, since we could consider a multiplayer Zero-sum Game as consisting of many smaller Zero-sum Games.

Definition 3 (Zero-Sum Game). A *zero-sum game* is a strictly competitive game $\langle \{I,II\},(X,Y),(\mu_I,\mu_{II})\rangle$ consisting of

- the set of two players $\{I, II\}$;
- the finite action sets X and Y of Player I and Player II respectively.
- the *utility function* μ_I for Player I, and μ_{II} for Player II.

In popular literature, a Zero-sum Game is often rep-

resented with just a finite utility matrix, where

	a_{11}	a_{12}	a_{13}	 a_{1n}
	a_{21}	a_{22}	a_{23}	 a_{2n}
$A = (a_{ij}) = [$	a_{31}	a_{32}	a_{33}	 a_{3n}
	a_{m1}	a_{m2}	a_{m3}	 a_{mn}

is the *utility matrix* for one of the player (say, Player I) with the a_{ij} component representing the utility $\mu_I(s_{ij})$ Player I receives when he plays his i^{th} action and Player II plays his j^{th} action. Since the utility matrix for Player II is then simply -A, the matrix A alone fully characterizes a Zero-sum Game. In this paper, we assume that the utility matrix for Player I is non-negative [2]. Given an utility matrix A of a Zero-sum Game, this condition can be ensured by adding a constant cto each element of the original utility matrix - an action analogous to requiring Player II pay an admission fee c to play the game. Note that doing this also does not alters the strategies p^* and q^* that maximize the expected payoff of Player I and II for any given game, since both seek to maximize their respective payoffs regardless. So, for our current purpose to analyze a game for its Nash Equilibrium, such an action does not 'fundamentally' change a game.

B. Strategy in a Zero-sum Game

Here we describe the strategy in which a player in a Zero-sum Game can adopt.

Definition 4 ((Mixed) Strategy). Let $G = \langle N, (A_i), (\mu_i) \rangle$ be a strategic game. A (mixed) strategy of player i is a probability distribution over A_i .

By this definition, in our zero-sum game, a strategy of Player I is a stochastic vector $\mathbf{p} = \begin{bmatrix} p_1 & p_2 & \cdots & p_m \end{bmatrix}^T$ where

$$\sum_{i} p_i = 1 \; ; \; p_i \ge 0, \; \text{with } i = 1, 2, \dots, m$$

Similarly, a strategy of Player II is a stochastic vector $\mathbf{q} = \begin{bmatrix} q_1 & q_2 & \cdots & q_n \end{bmatrix}^T$ where

$$\sum_{j=1}^{n} q_{j} = 1 \; ; \; q_{j} \ge 0 \text{ with } j = 1, 2, \dots, n$$

Given chosen strategies p and q, the probability that Player I plays the i^{th} strategy, and Player II plays the j^{th} strategy is p_iq_j . The expected utility to Player I given the chosen strategies being p and q are then

$$\sum_{i=1}^{m} \sum_{j=1}^{n} a_{ij} p_i q_j = \sum_{i=1}^{m} p_i \sum_{j=1}^{n} a_{ij} q_j = \mathbf{p}^T A \mathbf{q}.$$

¹Alternative constructions of preference relations are out of scope of this project, and are explored in the field of *Utility/Decision Theory*.

By symmetry, the expected utility of Player II is $-p^T Aq$.

C. Solution Concept

As is mentioned, each 'game' comes attached with a 'solution concept' that predicts how it would be played. In our zero-sum game, assuming that each player is intelligent and plays to maximize his/her profit, a game would be in equilibrium (that is, in a state where no player is incentivized to deviate from) if both players do not increase their utility by altering their action. This intuition is captured by the concept of the popular *Nash Equilibrium*.

Definition 5 (Nash Equilibrium). A pair of mixed strategies (p_*, q_*) is in a Nash Equilibrium if

$$p_*^T A q_* \ge p^T A q_*$$
 for all mixed strategy p
 $p_*^T A q_* \le p_*^T A q$ for all mixed strategy q

It could be shown that, in a Zero-sum Game, the *Nash Equilibrium* is equivalent to another class of solution concept - the *Minimax Equilibrium*.

Definition 6 (Minimax Equilibrium). In a zero-sum game, a pair of mixed strategies (p_*, q_*) is in a *Minimax Equilibrium* if

$$\boldsymbol{p}_*^T A \boldsymbol{q}_* = \max_{\boldsymbol{p}} \min_{\boldsymbol{q}} \boldsymbol{p}^T A \boldsymbol{q} = \min_{\boldsymbol{q}} \max_{\boldsymbol{p}} \boldsymbol{p}^T A \boldsymbol{q}$$

We call p_* the *maximin strategy* of Player I and q_* the *minimax strategy* of Player II.

The intuition behind this equilibrium is that Player I, assuming that Player II plays $\arg\min_{\boldsymbol{q}} \boldsymbol{p} A \boldsymbol{q}$ to minimize her loss, will play a strategy that maximize this value, that is

$$\max_{\boldsymbol{p}} \min_{\boldsymbol{q}} \boldsymbol{p}^T A \boldsymbol{q}.$$

The situation is in reverse for Player II. If it happens that the *maximin* strategy of Player I and the *minimax* strategy of Player II aligns, then nobody is motivated to change alter their actions. Thus, the game is in equilibrium.

Example (Rock-Paper-Scissor). The game of Rock-Paper-Scissor is a classic zero-sum game with the following expected utility matrix

$$A = \begin{array}{|c|c|c|c|c|c|} \hline & R & P & S \\ \hline R & 0 & 1 & -1 \\ \hline P & -1 & 0 & 1 \\ \hline S & 1 & -1 & 0 \\ \hline \end{array}$$

where, say, Player I receives 1 dollar if he/she wins, 0 if it is a tie, and lose 1 dollar if he/she wins. The *Minimax Equilibrium*, and by extension the *Nash Equilibrium*, for the game is the pair (p_*, q_*) where

 $p_* = q_* = \begin{bmatrix} 1/3 & 1/3 & 1/3 \end{bmatrix}^T$. Note that if the strategy of Player I favoring an action, say declaring Paper, over the other will be countered by a strategy that favors Scissor by Player II (the best strategy of Player II in this case is in fact to play exclusively Scissor!). This original game can be altered to a game with non-negative utility matrix for one of the player by requiring a 2 dollars admission fee from the other player where

		R	P	S
$A = \begin{cases} 1 & \text{if } A = A = \begin{cases} 1 & \text{if } A = A = A = A = A = A = A = A = A = A$	R	2	3	1
	P	1	2	3
	S	3	1	2

Assuming both players want to maximizes their utility, the *Nash Equilibrium* is still

$$\boldsymbol{p}_* = \boldsymbol{q}_* = \begin{bmatrix} 1/3 & 1/3 & 1/3 \end{bmatrix}^T$$
.

III. MATHEMATICAL ANALYSIS

A. Linear Programming

The key to determining the *Minimax Equilibrium*, and by extension, the *Nash Equilibrium* of a Zero-sum Game is to turn the minimizing - maximizing task of the two players involved into a *dual linear program*.

Definition 7 (Linear Duality). A *primal linear program* has the form

maximize
$$c^T x$$

subject to $Ax \le b$
 $x > 0$

while the associated dual linear program is to

minimize
$$b^T y$$

subject to $A^T y \ge c$
 $y > 0$

Specifically, the *minimax strategy* of Player II corresponds to a solution of a primal linear program, while the *maximin strategy* of Player I is a solution to the corresponding dual program. First, recognize that

$$u = \min_{\boldsymbol{q}} \boldsymbol{p}^T A \boldsymbol{q} = \min_{j} \sum_{i=1}^{m} a_{ij} p_i$$

where q is set to be 1 on the j^{th} component and 0 elsewhere. We thus have

$$u \le \sum_{i=1}^{m} a_{ij} p_i, \ \forall j = 1, 2, \dots, n$$

This implies

$$A^T \mathbf{p} \ge u\mathbf{e}, \ \mathbf{e} = \begin{bmatrix} 1 & 1 & \cdots & 1 \end{bmatrix}^T$$

Player I thus wants to

4

maximize
$$u$$
 subject to $A^T \mathbf{p} \ge u \mathbf{e}$ $\mathbf{p}^T \mathbf{e} = 1$ $\mathbf{p} \ge 0$

Letting $\mathbf{y} = \mathbf{p}/u$ yields $1/u = \mathbf{e}^T \mathbf{p}/u = \mathbf{e}^T \mathbf{y}$. We could thus reconstruct the above into a dual linear program:

minimize
$$e^T y$$

subject to $A^T y \ge e$ (1)
 $y \ge 0$

Similarly, Player II wants to

minimize
$$v$$
 subject to $A^T \mathbf{q} \leq v \mathbf{e}$ $\mathbf{q}^T \mathbf{e} = 1$ $\mathbf{q} \geq 0$

where $v = \max_{p} \mathbf{p}^{T} A \mathbf{q}$. Let $\mathbf{x} = q/v$, we could reconstruct this into a primal linear program:

maximize
$$e^T x$$

subject to $Ax \le e$ (2)
 $x \ge 0$

It could be seen that (2) and (1) are respectively two primal and dual linear programs. A classical in linear programming called the *Duality Theorem*, which states that the optimal value of the primal and dual linear program is the same, then asserts the existence of a *Nash Equilibrium* in every Zero-sum Game.

Theorem 1 (Duality Theorem). Suppose that both the primal and dual linear programming problem have feasible solutions. Then both have optimal solutions and

$$\max \boldsymbol{c}^T \boldsymbol{x} = \min \boldsymbol{b}^T \boldsymbol{y}$$

Given our primal-dual linear programs (2) and (1), there exist optimal solutions $\boldsymbol{x^*}$ and $\boldsymbol{y^*}$ and

$$\frac{1}{v^*} = e^T x^* = e^T y^* = \frac{1}{u^*}.$$

The strategies

$$p_* = u^* y^* \; ; \; q_* = v^* x^*$$

are thus the solution to the original Minimax - Maximin problem of Player I and Player II. In other words, (p_*, q_*) is in Minimax Equilibrium, and thus in Nash Equilibrium. This result, which states that the Minimax Equilibrium of a Zero-sum Game always exists, is also the result of the famed Minimax Theorem, proved by von

Neumann in 1928.

Theorem 2 (Minimax Theorem). Let A be an $m \times n$ matrix. There exist a stochastic n-vector p^* and a stochastic m-vector q^* such that

$$\boldsymbol{p}_*^T A \boldsymbol{q}_* = \max_{\boldsymbol{p}} \min_{\boldsymbol{q}} \boldsymbol{p}^T A \boldsymbol{q} = \min_{\boldsymbol{q}} \max_{\boldsymbol{p}} \boldsymbol{p}^T A \boldsymbol{q}$$

IV. ALGORITHMS & COMPARISON

With the linear program formulation and conversion of inequality constraints to equality constraints, we could solve the Nash equilibrium problem using the linear programming method, specifically the Simplex Algorithm and Interior Point Method (Predictor-Corrector).

A. Simplex Method

The simplex method could be thought of as a descent method that moves from one set of basic variables to another until the current basic variables are optimal. Specifically, as it advances from one basis to the adjacent basis, the simplex method removes and appends a single variable from each basis. This signifies transitioning between neighboring fundamentally viable solutions, which decreases the objective function. The simplex method traverses the feasible region along its boundaries.

Let c_B be the coefficients for the basic variables, c_B be coefficients for the non-basic variables, B be the coefficient matrix for the constraints of the basic variables, and N be the complementary matrix of B.

Algorithm 1 Simplex Algorithm

Solve for a dual feasible point

while Basic is not optimal: $c_N^T - c_B^T B^{-1} N < 0$ do

Choose a non-basic variable variable x_i that satisfies $\hat{c}_i < 0$ to enter the basic vector

Increased the entering variable long as all the other basic variables remain nonnegative

Choose a basic variable that reaches zero along the increase of the entering variable to remove from the basic vector.

Update B and x_B

end while

B. Interior Point Method

Unlike the Simplex Algorithm, where it moves along the edges of the feasible set to each vertex, the Interior Point Method (Primal-Dual Algorithm) starts at a feasible point in the feasible set and moves along the central path (μ) , where μ is the duality gap until μ reaches zero.

A solution to a linear programming problem must satisfy the complementary slackness conditions. The

condition on complementary slackness is equivalent to the condition on duality gap:

$$\boldsymbol{c}^T \boldsymbol{x} - \boldsymbol{b}^T \boldsymbol{y} = \boldsymbol{x}^T \boldsymbol{s} = n\mu$$

We would like to choose a search direction Δx , Δy , Δs such that $x + \Delta x$ is feasible, $y + \Delta y$ is feasible, and $(x + \Delta x)(y + \Delta y)$ satisfy the complementary slackness.

$$A\Delta x = \mathbf{b} - A\mathbf{x}$$

 $\Delta y + \Delta x = \mathbf{c} - A^T \mathbf{y} - \mathbf{s}$
 $(\mathbf{x} + \Delta \mathbf{x})(\mathbf{y} + \Delta \mathbf{y}) = \mu \mathbf{e}$

At each iteration, we update to reduce μ . If x, y, s are infeasible points, the problem could be modified to be equivalent to

$$A\Delta x = \mathbf{b} - A\mathbf{x} = \mathbf{r}_{P}$$
$$\Delta y + \Delta x = \mathbf{c} - A^{T}\mathbf{y} - \mathbf{s} = \mathbf{r}_{D}$$
$$S\Delta \mathbf{x} + X\Delta \mathbf{s} = \mu \mathbf{e} - XS\mathbf{e} - \Delta X\Delta S\mathbf{e}$$

where r_P is the residual for the primal constraints and r_d is the residuals for the dual constraints, $X = diag(x_i)$, $S = diag(s_i)$, $\Delta S = diag(\Delta s_i)$, $\Delta X = diag(\Delta x_i)$. In this case, we would also want to minimize the residuals.

The Predictor-Corrector Algorithm modifies the Primal-Dual Algorithm to reduce the number of iterations after each iteration. The 'predictor' step ignore $\mu e - \Delta X \Delta S e$ to find $\Delta \hat{X}$ and $\Delta \hat{X}$, and subsequently \hat{X} , \hat{X} , and μ . The 'corrector' step solves for Δx , Δy , Δs by substituting $\Delta \hat{X}$, $\Delta \hat{X}$, and μ into the equation:

$$A\Delta x = \mathbf{b} - A\mathbf{x} = \mathbf{r}_{P}$$

$$\Delta y + \Delta x = \mathbf{c} - A^{T}\mathbf{y} - \mathbf{s} = \mathbf{r}_{D}$$

$$S\Delta \mathbf{x} + X\Delta \mathbf{s} = \mu \mathbf{e} - XS\mathbf{e} - \Delta \hat{X}\Delta \hat{S}\mathbf{e}$$

Algorithm 2 Predictor-Corrector Algorithm

Calculate the initial points

for k = 0, 1, 2,... do

Solve for $\Delta \hat{X}$, $\Delta \hat{X}$

Substitute $\Delta \hat{X}$, $\Delta \hat{X}$ back to solve for Δx , Δy , Δs

Reduce μ : $\mu_{k+1} = \theta \mu_k$

end for

V. COMPUTATIONAL EXPERIMENT

Using the Simplex Algorithm (Dual Simplex) and Interior Point Method (Predictor-Corrector), we tested the computational aspects of solving the Nash equilibrium

for 1000 iterations, each on two experimental utility matrices (of size 10x10 and 100x100) created randomly.

The results align with the theory of the simplex algorithm and interior point methods, where simplex, with its time complexity in the worst case being exponential, tends to perform slower. While on larger problems, the interior point method is less stable.




Fig. 1: Compare Runtime on 10x10 utility matrix using Dual and Interior Point Method




Fig. 2: Compare Runtime on 100x100 utility matrix using Dual and Interior Point Method

VI. DISCUSSION

A natural extension of Zero-sum Game is to Generalsum Game, where, depending on the actions of players, the total utility of the two players might change. John Nash proves that in such a game, there always exist at least one Nash Equilibrium.

Theorem 3 (Nash's Theorem). In any game involving a finite number of players, where each player have mixed

strategy from a finite set of pure strategies, there exists at least one Nash equilibrium.

Nash's Theorem, however, is only existential, that is, it does not points out the in ways in which we can construct a mixed-strategy Nash Equilibrium in General-sum Game. Current approach includes formulating the game as a *linear complementarity problem*, then uses graph-search algorithm to seek for *best response strategies*, which form the *support* for the mixed-strategy. However, at the moment, even the state-of-the-art algorithm for this purpose - the Lemke-Howson, is not polynomial in worst case.

ACKNOWLEDGEMENT

We would like to thank Prof. Hieu Nguyen at Fulbright University Vietnam for providing us with a primer on linear programming and duality in MATH310: Optimization.

REFERENCES

- [1] R. B. Myerson, *Game Theory: Analysis of Conflict.* Harvard University Press, Mar. 2013. [Online]. Available: http://www.jstor.org/stable/10.2307/j.ctvjsf522
- [2] A. Ostaszewski, Advanced mathematical methods, ser. London School of Economics mathematics series. Cambridge; New York: Cambridge University Press, 1990, oCLC: 818873366.