

**STAT 2400**

**Introduction to Probability I**

Solutions to Sample Test #1 (C)

**Question 1:** (15 marks: 2.5 each)

How many different student lineups are there if

- (A) the students are lined up completely at random?

*Solution:*  $N(\cdot) = 20!$

- (B) the smallest student and tallest student of the class necessarily come first and last, respectively?

*Solution:*  $N(\cdot) = 18!$

- (C) Andrew, Carrie and Kathy will not be separated in the line?

*Solution:*  $N(\cdot) = 18! \times 3!$

- (D) Andrew, Carrie and Kathy will not be separated in the line, but Andrew is also the tallest student in the class and has to be last in line?

*Solution:*  $N(\cdot) = 17! \times 2$

- (E) Andrew and Carrie have been fighting and so, won't be next to each other in line;

*Solution:*  $N(\cdot) = 20! - 19! \times 2$

- (F) boys and girls should alternate in the line (assuming the class has 10 boys and 10 girls).

*Solution:*  $N(\cdot) = (10!)^2 \times 2$

**Question 2:** (5 marks: 2.5 each)

Let  $A$  and  $B$  be events of  $\Omega$  such that

$$\mathbb{P}(A) = 0.2, \quad \mathbb{P}(A \cup B) = 0.6 \quad \text{and} \quad \mathbb{P}(A \cap B) = 0.1.$$

- (A) Determine  $\mathbb{P}(B)$ .

*Solution:* From the general addition rule we have that

$$\mathbb{P}(B) = \mathbb{P}(A \cup B) - \mathbb{P}(A) + \mathbb{P}(A \cap B) = 0.5.$$

- (B) Determine  $\mathbb{P}(A \cup B^c)$ .

*Solution:* Again, using the general addition rule,

$$\begin{aligned} \mathbb{P}(A \cup B^c) &= \mathbb{P}(A) + \mathbb{P}(B^c) - \mathbb{P}(A \cap B^c) \\ &= \mathbb{P}(A) + \mathbb{P}(B^c) - [\mathbb{P}(A) - \mathbb{P}(A \cap B)] = \mathbb{P}(B^c) + \mathbb{P}(A \cap B) = 0.6. \end{aligned}$$

**Question 3:** (4 marks)

What is the probability the word ABRACADABRA is formed?

*Solution:* There are here

$$\binom{11}{5, 2, 1, 1, 2} = \frac{11!}{5!(2!)^2} = 83,160$$

equally likely words that can be formed when drawing the balls. Hence  $\mathbb{P}(\cdot) = 1/83,160$ .

**Question 4:** (4 marks)

Find  $\mathbb{P}(A_i)$  for  $i = 4, 5, \dots, 10$ .

*Solution:* In this case,

$$\mathbb{P}(A_i) = \frac{\binom{i-1}{3}(4)_4(6)_{i-4}}{(10)_i},$$

with the numerator coming from the fact that there are  $i - 1$  positions to choose from for the first three red balls (the fourth red ball needs to be in position  $i$ ) and that 4 red and  $i - 4$  black balls need to be selected. This formula leads to the following table of probabilities:

$i$	4	5	6	7	8	9	10
$\mathbb{P}(A_i)$	1/210	2/105	1/21	2/21	1/6	4/15	2/5
	0.0048	0.0190	0.0476	0.0952	0.1667	0.2667	0.4000

Actually, a much easier approach is to write this as

$$\mathbb{P}(A_i) = \frac{\binom{i-1}{3}}{\binom{10}{4}},$$

which comes from selecting the positions of the red balls when drawing all 10 balls. For the numerator, we need a red ball in position  $i$  and the three other reds in the first  $i - 1$  positions.

**Question 5:** (4 marks)

What is the probability that a double six is obtained at least once?

*Solution:* This uses the complementation rule

$$\mathbb{P}() = 1 - \mathbb{P}(\text{"no double six"}) = 1 - \left(\frac{35}{36}\right)^n,$$

since

$$\mathbb{P}(\text{"no double six"}) = \frac{N(\text{"no double six"})}{N(\Omega)} = \frac{35^n}{36^n}.$$

**Question 6:** (4 marks)

According to Barbara, what is the probability that she will be hired?

*Solution:* Let  $H_J$ ,  $H_M$  and  $H_B$  denote the events that John, Martin and Barbara are hired, respectively. Let also  $\mathbb{P}(H_J) = \mathbb{P}(H_M) = p$ . Then, it should be clear that

$$\mathbb{P}(H_J) + \mathbb{P}(H_M) + \mathbb{P}(H_B) = 1, \quad \text{and that} \quad \mathbb{P}(H_B) = \frac{3}{2}p.$$

From these, we have that

$$1 = \frac{7}{2}p, \quad \text{or } p = \frac{2}{7} \simeq 0.289.$$

Hence, for Barbara,

$$\mathbb{P}(H_B) = \frac{3}{2}p = \frac{3}{7} \simeq 0.429.$$

**Question 7:** (4 marks)

State Kolmogorov's axioms of probability.

*Solution:* The three axioms are:

- 1) The nonnegativity axiom: for any event  $E \subset \Omega$ ,

$$\mathbb{P}(E) \geq 0.$$

- 2) The certainty axiom:

$$\mathbb{P}(\Omega) = 1.$$

- 3) The countable additivity axiom: for  $A_1, A_2, \dots$  disjoint events of  $\Omega$ ,

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i).$$

**Question 8:** (8 marks: 4 each)

- (A) Prove Bonferroni's inequality.

*Solution:* We start with the general addition rule, which implies that

$$\mathbb{P}(A \cap B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cup B) \geq \mathbb{P}(A) + \mathbb{P}(B) - 1,$$

since  $\mathbb{P}(A \cup B) \leq 1$ .

(B) Show that  $\mathbb{P}(A \cap B) = \mathbb{P}(A \cup B) = 1$ .

*Solution:* Firstly,  $A \subset A \cup B$ , so that the domination principle implies

$$1 = \mathbb{P}(A) \leq \mathbb{P}(A \cup B),$$

implying, in turn, that  $\mathbb{P}(A \cup B) = 1$ .

Secondly, using Bonferroni's equality,

$$\mathbb{P}(A \cap B) \geq \mathbb{P}(A) + \mathbb{P}(B) - 1 = 1,$$

implying  $\mathbb{P}(A \cap B) = 1$ .

**Question 9:** (4 marks)

Prove that for  $0 \leq i \leq k \leq m \leq n$ ,

$$\binom{n}{m} \binom{m}{k} \binom{k}{i} = \binom{n}{i} \binom{n-i}{k-i} \binom{n-k}{m-k}.$$

*Solution:* Recall that

$$\binom{m'}{r} \binom{r}{k'} = \binom{m'}{k'} \binom{m'-k'}{r-k'}.$$

It suffices to use this equality twice to get the result. Indeed, applying this with  $m' = n$ ,  $r = m$  and  $k' = k$ , we have

$$\binom{n}{m} \binom{m}{k} = \binom{n}{k} \binom{n-k}{m-k},$$

so that

$$\binom{n}{m} \binom{m}{k} \binom{k}{i} = \binom{n}{k} \binom{k}{i} \binom{n-k}{m-k}.$$

Using the same argument with  $m' = n$ ,  $r = k$  and  $k' = i$  leads to the wanted result.

A proof based on a combinatorial argument is also acceptable here.