# Kaya clinical summary

Patient 1705
Generated 18 Oct 2026 14:11

Synthetic patient. Decision support for clinicians; not a diagnostic device and not clinically validated.

| | | | |
|---|---|---|---|
| **Patient** | 74 y, female; BMI 25.1; waist 80 cm | **Diabetes** | 2.9 years; HbA1c 8.5%; BP 126/75 |
| **Treatment** | Metformin | **Genetics** | PNPLA3 G alleles 2; TCF7L2 T alleles 0 |
| **Priority** | **MEDIUM** | **Why flagged** | Time in range 42%; Advanced fibrosis risk 36%; 6 alert(s) in the last 24 h |

## Glucose now and the next 2 hours

| Now | In 30 min | In 60 min | In 120 min | High within 2 h | Low within 2 h |
|---|---|---|---|---|---|
| **275** mg/dL | 266 (253-281) | 235 (218-251) | 192 (180-204) | 99% | 0% |




Last 3 days of CGM with the twin's forecast (median and 80% range) from the latest reading. Healthy range 70-180 mg/dL shaded.

## 14-day glucose profile

| Time in range | Above 180 | Below 70 | Variability (CV) | Mean glucose | Est. HbA1c |
|---|---|---|---|---|---|
| 42% | 58% | 0.0% | 18% | 196 mg/dL | 8.0% |

## Organ risk

| Organ | Finding | Evidence |
|---|---|---|
| **Liver** | Steatosis risk 19%, 26% at 12 months at current habits.<br>Advanced fibrosis 31.4%, 35.5% at 12 months. Last FibroScan 11.9 kPa; FIB-4 2.2.<br>Main drivers: waist 80 cm (lowers), HbA1c 8.5% (raises), ALT 35 U/L (raises). | Baseline: model validated on real NHANES FibroScan data (AUROC 0.83). 12-month drift from glucose habits: simulation. |
| **Eyes** | No retinopathy (grade 0). Two-year progression risk 6%. | Image grader validated on APTOS (kappa 0.89) and Indian IDRiD (0.71 after fine-tuning). Progression risk: simulation. |
| **Kidneys** | eGFR 65, UACR 12 mg/g. Albuminuria risk 30% now, 33% at 12 months. eGFR about 56 in 5 years (1.7 per year); reaches 60 in about 2.9 years at this rate.<br>Main drivers: HbA1c 8.5% (raises), diabetes 1.0 (raises), age 74y (raises). | Risk model validated on real NHANES data (AUROC 0.76 albuminuria, 0.85 reduced eGFR). eGFR path: simulation. |

## What the twin suggests

| Scenario (simulation) | Time in range | Liver fat risk, 12 m | eGFR in 5 y | Eye progression, 2 y |
|---|---|---|---|---|
| Current habits | 42% | 26% | 56 | 6% |
| Walk 20 min after meals | 43% | 18% | 57 | 4% |
| Smaller carb portions (-25%) | 43% | 23% | 57 | 5% |
| Medication optimised (-15 mg/dL) | 69% | 18% | 57 | 3% |
| All three together | 71% | 11% | 58 | 2% |

Effect sizes are literature-informed assumptions shown for transparency; this is a simulation to support a conversation, not a prediction of outcome.

## Alerts, last 24 hours

| When | Alert | Why |
|---|---|---|
| Wed 14 Jan, 20:45 | High glucose expected (100%) | Glucose 178 and steady, no recent meal logged. |
| Wed 14 Jan, 18:40 | High glucose expected (87%) | Glucose 178 and steady, no recent meal logged. |
| Wed 14 Jan, 12:05 | High glucose expected (93%) | Glucose 168 and steady, no recent meal logged. |
| Wed 14 Jan, 06:50 | High glucose expected (99%) | Glucose 177 and steady, no recent meal logged. |
| Wed 14 Jan, 04:50 | High glucose expected (87%) | Glucose 175 and steady, no recent meal logged. |
| Wed 14 Jan, 02:50 | High glucose expected (90%) | Glucose 179 and steady, no recent meal logged. |

Kaya is an open-source research prototype built on open and synthetic data (CGMacros, NHANES, APTOS, IDRiD). Forecasts and risks are estimates to support clinical judgement. Simulations are labelled and their assumptions are editable.