

Physics Research Note1

Jizheng Bo

April 15, 2026

Contents

Chapter 1 Proton Decay 的 Bayesian 计算	5
1.1 试运行	5
Appendix A Supplementary Calculations	11

Chapter 1

Proton Decay 的 Bayesian 计算

考虑形变势能:

$$V(r, \theta) = V_N(r, \theta) + V_c(r, \theta) + V_{so}(r, \theta) + \frac{\hbar^2}{2\mu r^2} \left(l + \frac{1}{2}\right)^2 \quad (1.1)$$

$$V_N(r, \theta) = -\frac{V_{N0}}{1 + \exp\{r - R_N(\theta)/a_N(\theta)\}} \quad (1.2)$$

$$R_N(\theta) = R_{N0}(1 + \beta_2 Y_{20}(\theta) + \beta_4 Y_{40}(\theta)) \quad (1.3)$$

$$a_N(\theta) = a_{N0} \sqrt{1 + \left(\frac{1}{R_N(\theta)} \frac{dR_N(\theta)}{d\theta}\right)^2} \times [1 + \tilde{\beta}_2 Y_{20}(\theta) + \tilde{\beta}_4 Y_{40}(\theta)] \quad (1.4)$$

$$V_{so} = V_{so0} \lambda_\pi^2 \frac{1}{r} \frac{d}{dr} \frac{1}{1 + \exp[(r - R_{so})/a_{so}]} \vec{\sigma} \cdot \vec{l} \quad (1.5)$$

$$\lambda_\pi^2 = 2fm^2 \quad (1.6)$$

$$V_c(r) = \begin{cases} \frac{Z_D Z_p e^2}{2R_{c0}} \frac{3R_{c0}^2 - r^2}{R_{c0}^2}, & r \leq R_{c0} \\ \frac{Z_D Z_p e^2}{r}, & r > R_{c0} \end{cases} \quad (1.7)$$

1.1 试运行

使用参数 (145Tm, 形变势能,wkb 积分)

```
#System
e2=1.43997 ; hbarc=197.3269718 ; amu=931.49432
zp=69 ; Ap=145 ; mass_excess_p=-27.58 ; mp=Ap*amu+mass_excess_p
z1=68 ; A1=144 ; mass_excess_1=-36.61 ; m1=A1*amu+mass_excess_1 #1-> daughter
z2=1 ; A2=1 ; mass_excess_2=7.288971064 ; m2=A2*amu+mass_excess_2
Q=mp-m1-m2
mu=m1*m2/(m1+m2)
P0=0.580
#Potential
```

```

#Deformation
beta2 = 0.231 # quadrupole
beta4 = -0.068 # hexadecapole

#Polarizaton
beta2tilde = 0
beta4tilde = 0

#spin-orbit
Vso0=6.2 ; rso=1.01*(A1)**(1/3) ; aso=0.75 ; lambda_pi=np.sqrt(2)
L=5; S=0.5; J=5.5

#Vn
a0=0.75 ; r0=1.27*A1**(1/3)-0.1 ; V0=55

#Vc
Rc0 = 1.21 * A1**(1/3)

# experiment data
y=np.array([3.16e-6]) # s
sigma=np.array([0.16e-6])

#a = v0,a0,r0,p0,rc0,beta2,beta4,vso0,rso,aso
def log_prior(a):
    v0,a0,r0,p0,rc0,beta2,beta4,vso0,rso,aso=a
    R=1

# Hyperparameters
v0_mean=55; a0_mean=0.75; r0_mean=6.556; p0_mean=0.580
rc0_mean=6.342; beta2_mean=0.231; beta4_mean=-0.068
vso0_mean=6.2; rso_mean=5.294; aso_mean=0.75

sigma_v0=1*R; sigma_a0=0.05*R; sigma_r0=0.1*R; sigma_p0=0.05*R
sigma_rc0=0.1*R; sigma_beta2=0.05*R; sigma_beta4=0.01*R
sigma_vso0=1*R; sigma_rso=0.1*R; sigma_aso=0.05*R

with multiprocessing.Pool() as pool:
    sampler = emcee.EnsembleSampler(nwalkers, M, log_posterior, args=[y,sigma],a=1.0,
    pool=pool)
    state = sampler.run_mcmc(initial_positions, 200)
    sampler.reset()

```

```
sampler.run_mcmc(state,10000)
```

获得 ^{145}Tm 的 Posterior 分布如下图所示:

(说明之前的 bug 出自于对 model 函数的解包错误, 而非采样参数错误)




Figure 1.1: Posterior distribution of the parameters for ^{145}Tm proton decay.

上图在 helium1 上计算, 耗时约 3.5d, 速度较慢

后续预计先调整图片样式, 然后做好代码优化加速, 最后可以选取具体核区正式计算
我们使用如下代码修改样式:

```
import prettyplease
from scipy.stats import norm
samples = sampler.get_chain(flat=True)

#a = v0,a0,r0,p0,rc0,beta2,beta4,vso0,rso,aso

labels = ["$V_0$", "$a_0$", "$r_0$", "$P_0$", "$R_{c0}$",
          "$\\beta_2$", "$\\beta_4$", "$V_{so0}$", "$R_{so}$", "$a_{so}$"]
fig, axes = prettyplease.corner(
    samples,
    labels=labels,
```

```

    bins=20,
    colors=['white', '#333333'],
    linewidth=1.0,
    fontsize=10,
    n_ticks=3,
    return_axes=True
)

# ----- 添加 1 区间竖线 -----
ndim = samples.shape[1]      # 参数个数

# 先验参数定义 (与 log_prior 保持一致)
prior_means = np.array([55, 0.75, 6.556, 0.580, 6.342, 0.231, -0.068, 6.2, 5.294, 0.75])
prior_sigmas = np.array([1, 0.05, 0.1, 0.05, 0.1, 0.05, 0.01, 1, 0.1, 0.05])

for i in range(ndim):
    ax = axes[i, i]          # 对角线子图
    data = samples[:, i]

    # 计算 16% 和 84% 分位数 (对应 1 区间边界)
    q16, q84 = np.percentile(data, [16, 84])

    # 绘制两条虚线竖线
    ax.axvline(q16, color='red', linestyle='--', linewidth=1.0, alpha=0.8)
    ax.axvline(q84, color='red', linestyle='--', linewidth=1.0, alpha=0.8)
    median = np.median(data)
    ax.axvline(median, color='blue', linestyle='--', linewidth=1.0, alpha=0.8)

# ----- 叠加先验曲线 -----
for i in range(ndim):
    ax = axes[i, i]
    xlim = ax.get_xlim()
    x_plot = np.linspace(xlim[0], xlim[1], 200)
    pdf = norm.pdf(x_plot, loc=prior_means[i], scale=prior_sigmas[i])
    ax.plot(x_plot, pdf, color='green', linestyle='--', linewidth=1.2, alpha=0.8)
    ax.set_xlim(xlim) # 恢复原范围

# plt.savefig('145Tm.pdf')
plt.savefig('test.pdf')
plt.show()

```

示例图为




Figure 1.2: 后续计算使用的后验图的示例，包括增加了 prior 线，以及 1σ , 2σ , median 的标定，还使用了灰色线图

Appendix A

Supplementary Calculations

Put detailed derivations that are too long for the main text here.