



THE UNIVERSITY OF TEXAS AT AUSTIN  
McCOMBS SCHOOL OF BUSINESS

# Probability review 1

---

## Lecture 3

STA 371G

# Announcements

- Homework 1 due Sunday night at 11:59 PM

## Announcements

- Homework 1 due Sunday night at 11:59 PM
- First (optional) R help session will be this Friday from 11 AM to 12 PM in GSB 3.138 (note temporary room change)

# Announcements

- Homework 1 due Sunday night at 11:59 PM
- First (optional) R help session will be this Friday from 11 AM to 12 PM in GSB 3.138 (note temporary room change)
- About grading

# Announcements

- Homework 1 due Sunday night at 11:59 PM
- First (optional) R help session will be this Friday from 11 AM to 12 PM in GSB 3.138 (note temporary room change)
- About grading
  - QUEST: two tries on each question for full credit, then decreasing credit for each subsequent try

# Announcements

- Homework 1 due Sunday night at 11:59 PM
- First (optional) R help session will be this Friday from 11 AM to 12 PM in GSB 3.138 (note temporary room change)
- About grading
  - QUEST: two tries on each question for full credit, then decreasing credit for each subsequent try
  - Learning Catalytics: answer 75% of in-class questions for full participation credit

What are the chances that two people *in this room* have the same birthday (month/day, not year)?



# Probability Theory

## *The Concept of Probability*

What do each of these have in common?



# Probability Theory

## *The Concept of Probability*

What do each of these have in common?

- Outcome of rolling a die

# Probability Theory

## *The Concept of Probability*

What do each of these have in common?

- Outcome of rolling a die
- S&P500 index at the end of January

# Probability Theory

## *The Concept of Probability*

What do each of these have in common?

- Outcome of rolling a die
- S&P500 index at the end of January
- Number of iPhone 7s to be sold over the next year

# Probability Theory

## *The Concept of Probability*

What do each of these have in common?

- Outcome of rolling a die
- S&P500 index at the end of January
- Number of iPhone 7s to be sold over the next year
- Number of unique visitors to Amazon.com over the next week

# Probability Theory

## *The Concept of Probability*

What do each of these have in common?

- Outcome of rolling a die
- S&P500 index at the end of January
- Number of iPhone 7s to be sold over the next year
- Number of unique visitors to Amazon.com over the next week
- Lifetime of your MacBook Air

# Probability Theory

## *The Concept of Probability*

What do each of these have in common?

- Outcome of rolling a die
- S&P500 index at the end of January
- Number of iPhone 7s to be sold over the next year
- Number of unique visitors to Amazon.com over the next week
- Lifetime of your MacBook Air

# Probability Theory

## *The Concept of Probability*

What do each of these have in common?

- Outcome of rolling a die
- S&P500 index at the end of January
- Number of iPhone 7s to be sold over the next year
- Number of unique visitors to Amazon.com over the next week
- Lifetime of your MacBook Air

We cannot predict any of these with certainty.

# Probability Theory

## *The Concept of Probability*

What do each of these have in common?

- Outcome of rolling a die
- S&P500 index at the end of January
- Number of iPhone 7s to be sold over the next year
- Number of unique visitors to Amazon.com over the next week
- Lifetime of your MacBook Air

We cannot predict any of these with certainty. But we can model them using **probability theory** and learn a lot about how these variables will behave.



# Probability Theory

## *Definitions*

### Definition

A **random variable** expresses the outcome of a random process as a number. It is denoted by an uppercase letter.

# Probability Theory

## *Definitions*

### Definition

A **random variable** expresses the outcome of a random process as a number. It is denoted by an uppercase letter.

- Random experiment → Selecting a student at random from the class

# Probability Theory

## Definitions

### Definition

A **random variable** expresses the outcome of a random process as a number. It is denoted by an uppercase letter.

- Random experiment  $\rightarrow$  Selecting a student at random from the class
- Random variable  $\rightarrow X$  : The day of their birthday  $(1, 2, \dots, 31)$

# Probability Theory

## Definitions

### Definition

A **random variable** expresses the outcome of a random process as a number. It is denoted by an uppercase letter.

- Random experiment  $\rightarrow$  Selecting a student at random from the class
- Random variable  $\rightarrow X$  : The day of their birthday (1, 2,  $\dots$ , 31)
- Random variable  $\rightarrow Y$  : Their height, in inches

# Probability Theory

## *Definitions*

### Definition

A **discrete random variable** is a random variable with a finite (or countably infinite) range.

# Probability Theory

## *Definitions*

### Definition

A **discrete random variable** is a random variable with a finite (or countably infinite) range.

A **continuous random variable** is a random variable with an interval (either finite or infinite) of real numbers for its range.

# Probability Theory

## *Definitions*

### Definition

A **discrete random variable** is a random variable with a finite (or countably infinite) range.

A **continuous random variable** is a random variable with an interval (either finite or infinite) of real numbers for its range.

- $X$  : Number of stocks on NYSE whose price change today (discrete)

# Probability Theory

## *Definitions*

### Definition

A **discrete random variable** is a random variable with a finite (or countably infinite) range.

A **continuous random variable** is a random variable with an interval (either finite or infinite) of real numbers for its range.

- $X$  : Number of stocks on NYSE whose price change today (discrete)
- $Y$  : Average price change of the stocks on NYSE (continuous)



# Probability Theory

## Definitions

### Definition

A **discrete random variable** is a random variable with a finite (or countably infinite) range.

A **continuous random variable** is a random variable with an interval (either finite or infinite) of real numbers for its range.

- $X$  : Number of stocks on NYSE whose price change today (discrete)
- $Y$  : Average price change of the stocks on NYSE (continuous)
- $Z$  : Number of people that lose money in the stock market today (discrete!)

# Probability Theory

## *Exercise*

Discrete or continuous?



# Probability Theory

## *Exercise*

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year



# Probability Theory

## *Exercise*

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year



# Probability Theory

## *Exercise*

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year (discrete)
- Lifetime of your MacBook Air, in years



# Probability Theory

## *Exercise*

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year (discrete)
- Lifetime of your MacBook Air, in years



# Probability Theory

## *Exercise*

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year (discrete)
- Lifetime of your MacBook Air, in years (continuous)
- Number of unique visitors to Amazon.com over the next week



# Probability Theory

## *Exercise*

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year (discrete)
- Lifetime of your MacBook Air, in years (continuous)
- Number of unique visitors to Amazon.com over the next week





# Probability Theory

## *Exercise*

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year (discrete)
- Lifetime of your MacBook Air, in years (continuous)
- Number of unique visitors to Amazon.com over the next week (discrete)
- S&P 500 index at the end of January



# Probability Theory

## *Exercise*

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year (discrete)
- Lifetime of your MacBook Air, in years (continuous)
- Number of unique visitors to Amazon.com over the next week (discrete)
- S&P 500 index at the end of January



# Probability Theory

## Exercise

Discrete or continuous?

- Number of iPhone 7s to be sold over the next year (discrete)
- Lifetime of your MacBook Air, in years (continuous)
- Number of unique visitors to Amazon.com over the next week (discrete)
- S&P 500 index at the end of January (continuous)



# Probability Theory

## *Definitions*

### Definition

**Probability** is the measure of the likelihood that a particular outcome (or set of outcomes) will be observed. Probability is a number always between 0 and 1.

# Probability Theory

## *Definitions*

### Definition

**Probability** is the measure of the likelihood that a particular outcome (or set of outcomes) will be observed. Probability is a number always between 0 and 1.

- $X$  : The month number of the student's birthday,  
 $P(X = 3) = 1/12$

# Probability Theory

## Definitions

### Definition

**Probability** is the measure of the likelihood that a particular outcome (or set of outcomes) will be observed. Probability is a number always between 0 and 1.

- $X$  : The month number of the student's birthday,  
 $P(X = 3) = 1/12$
- $Y$  : Lifetime of your MacBook,  $P(Y > 5 \text{ years}) = 0.05$

# Probability Distributions

Now that we have defined a random variable, how do we know what the probabilities are?

## Probability Distributions

Now that we have defined a random variable, how do we know what the probabilities are?

For example, what is the probability that your MacBook will break down after 5 years but before 7 years? That is,  $P(5 < X < 7) = ?$



# Probability Distributions

Now that we have defined a random variable, how do we know what the probabilities are?

For example, what is the probability that your MacBook will break down after 5 years but before 7 years? That is,  $P(5 < X < 7) = ?$

## Definition

The **probability distribution** of a random variable  $X$  is a description of the probabilities associated with the possible values of  $X$ .

# Probability Distributions

Now that we have defined a random variable, how do we know what the probabilities are?

For example, what is the probability that your MacBook will break down after 5 years but before 7 years? That is,  $P(5 < X < 7) = ?$

## Definition

The **probability distribution** of a random variable  $X$  is a description of the probabilities associated with the possible values of  $X$ .

Discrete random variable  $\rightarrow$  Probability Mass Function (p.m.f.)

# Probability Distributions

Now that we have defined a random variable, how do we know what the probabilities are?

For example, what is the probability that your MacBook will break down after 5 years but before 7 years? That is,  $P(5 < X < 7) = ?$

## Definition


The **probability distribution** of a random variable  $X$  is a description of the probabilities associated with the possible values of  $X$ .

Discrete random variable  $\rightarrow$  Probability Mass Function (p.m.f.)

Continuous random variable  $\rightarrow$  Probability Density Function (p.d.f.)

## Discrete Random Variables

Michael Beuoy's model for NFL games gives the following probabilities for the Super Bowl (positive margin = Patriots win):



## Discrete Random Variables

Let  $M$  be a random variable representing the Patriots' margin of victory.

Patriots' margin of victory $m$	Probability $P(M = m)$
—28	0.024
—27	0.006
—26	0.002
—25	0.004
$\vdots$	$\vdots$
27	0.008
28	0.033

# Discrete Random Variables

## Definition

The **expected value** and **variance** of a random variable  $X$  are the long run mean and variance if we draw values repeatedly from  $X$ .

# Discrete Random Variables

## Definition

The **expected value** and **variance** of a random variable  $X$  are the long run mean and variance if we draw values repeatedly from  $X$ .


$$\begin{aligned} E(M) &= \sum_{\text{all margins } m} m \cdot P(M = m) \\ &= (-28)(0.024) + (-27)(0.006) + \dots \end{aligned}$$

$$\begin{aligned} \text{Var}(M) &= \sum_{\text{all margins } m} (m - E(M))^2 \cdot P(M = m) \\ &= (-28 - E(M))^2(0.024) + (-27 - E(M))^2(0.006) + \dots \end{aligned}$$

## Normal Distributions (the “bell curve”)

The **normal distribution** is a common continuous probability distribution; we will denote a normal distribution with mean  $\mu$  and variance  $\sigma^2$  by  $N(\mu, \sigma^2)$ .





## Continuous Random Variables

Let  $V$  be a random variable indicating the SAT Verbal score of a randomly selected student in the US.

$$P(\text{SAT} > 600) =$$

## Continuous Random Variables


Let  $V$  be a random variable indicating the SAT Verbal score of a randomly selected student in the US.

$$P(\text{SAT} > 600) = \int_{600}^{800} P(V = v) dv$$

## Continuous Random Variables

Let  $V$  be a random variable indicating the SAT Verbal score of a randomly selected student in the US.

$$P(\text{SAT} > 600) = \int_{600}^{800} P(V = v) dv$$



## Continuous Random Variables


Let  $V$  be a random variable indicating the SAT Verbal score of a randomly selected student in the US.

$$P(\text{SAT} = 600) =$$

## Continuous Random Variables

Let  $V$  be a random variable indicating the SAT Verbal score of a randomly selected student in the US.

$$P(\text{SAT} = 600) = 0$$



## Continuous Random Variables

The formulas for expected value and variance are the same for continuous random variables, but they involve integrals instead of sums:

Discrete random variable  $X$

$$E(X) = \sum_x xP(X = x)$$

$$\text{Var}(X) = \sum_x (x - E(X))^2 P(X = x)$$

Continuous random variable  $Y$   
with density function  $f(y)$

$$E(Y) = \int_y yf(y) dy$$

$$\text{Var}(Y) = \int_y (y - E(Y))^2 f(y)$$



# R Simulation of expected value

## *R Simulation*

Let's simulate rolling dice and see that the expected value really does represent the long-run average!

```
# Simulate rolling a die 1 time
sample(c(1, 2, 3, 4, 5, 6), 1, replace=T)
# Simulate rolling a die 4 times
sample(c(1, 2, 3, 4, 5, 6), 4, replace=T)
# Take the average
mean(sample(c(1, 2, 3, 4, 5, 6), 4, replace=T))
# Let's increase the number of dice
mean(sample(c(1, 2, 3, 4, 5, 6), 100, replace=T))
```



## Joint probability – back to birthdays

If two events are *independent* (i.e., knowing the outcome of one tells you nothing about the outcome of the other), then the probability of two events happening at the same time is

$$P(A \text{ and } B) = P(A)P(B).$$

## Joint probability – back to birthdays

If two events are *independent* (i.e., knowing the outcome of one tells you nothing about the outcome of the other), then the probability of two events happening at the same time is

$$P(A \text{ and } B) = P(A)P(B).$$

Let's figure out the probability of no one sharing a birthday. Let's call  $B_1$  the first person's birthday,  $B_2$  the second person's birthday, etc.

## Joint probability – back to birthdays

If two events are *independent* (i.e., knowing the outcome of one tells you nothing about the outcome of the other), then the probability of two events happening at the same time is

$$P(A \text{ and } B) = P(A)P(B).$$

Let's figure out the probability of no one sharing a birthday. Let's call  $B_1$  the first person's birthday,  $B_2$  the second person's birthday, etc.

$$\begin{aligned} P(\text{none shared}) &= P(B_2 \neq B_1) \cdot P(B_3 \neq B_1, B_2) \cdot P(B_4 \neq B_1, B_2, B_3) \cdots \\ &= \frac{364}{365} \cdot \frac{363}{365} \cdot \frac{362}{365} \cdots \end{aligned}$$

## Joint probability – back to birthdays

If two events are *independent* (i.e., knowing the outcome of one tells you nothing about the outcome of the other), then the probability of two events happening at the same time is

$$P(A \text{ and } B) = P(A)P(B).$$

Let's figure out the probability of no one sharing a birthday. Let's call  $B_1$  the first person's birthday,  $B_2$  the second person's birthday, etc.

$$\begin{aligned} P(\text{none shared}) &= P(B_2 \neq B_1) \cdot P(B_3 \neq B_1, B_2) \cdot P(B_4 \neq B_1, B_2, B_3) \cdots \\ &= \frac{364}{365} \cdot \frac{363}{365} \cdot \frac{362}{365} \cdots \end{aligned}$$

When  $n = 70$  the probability of no shared birthdays is just 0.01%!