

Sum Law

Claim: $\lim_{x \rightarrow a} [f(x) - g(x)] = M - N$

Proof. Provided that $\lim_{x \rightarrow c} f(x) = M$, $\lim_{x \rightarrow c} g(x) = N$, and $N \neq 0$. Using the ε - δ definition of a limit, we will directly prove that:

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{M}{N}$$

The definition tell us, that for every $\varepsilon > 0$, there exists a $\delta > 0$ such that if $0 < |x - a| < \delta$, then $|f(x) - L| < \varepsilon$.

Suppose $0 < |x - a| < \delta$. Let us analyze the distance between the function and its limit value:

$$|f(x)/g(x) - M/N| = |f(x)(N - M) + M(f(x) - g(x))|/|g(x)N|$$

Then using the triangle inequality, we can estimate each part as follows:

$$\left| \frac{f(x)(N - M) + M(f(x) - g(x))}{g(x)N} \right| \leq \frac{|f(x)||N - M| + |M||f(x) - g(x)|}{|g(x)N|}$$

Hence, for every $\varepsilon > 0$, there exists a $\delta > 0$ such that if $0 < |x - c| < \delta$, then $|f(x) - g(x) - (M - N)| < \varepsilon$. Thus, we can conclude:

$$\lim_{x \rightarrow c} [f(x) - g(x)] = M - N.$$

To put a lower bound on $|g(x)|$, choose $\delta_1 = \frac{|N|^2}{2}$ such that if $|g(x) - N| < \delta_1$, then:

$$|g(x)||N| > \frac{|N|^2}{2} \Rightarrow \frac{1}{|g(x)||N|} < \frac{2}{|N|^2}$$

Then we can represent the total as:

$$\frac{|f(x)||N - M| + |M||f(x) - g(x)|}{|g(x)||N|} < (|f(x)||N - M| + |M||f(x) - g(x)|) \cdot \frac{2}{|N|^2} < \epsilon$$

Since the total expression is to be bounded by ϵ , choose $\delta_2 = \frac{\epsilon|N|}{4}$. We can bound each part individually as follows:

$$|f(x)||N - M| < \frac{\epsilon|N|}{4} \quad \text{and} \quad |M||f(x) - g(x)| < \frac{\epsilon|N|}{4}$$

Bounding $|f(x)|$, choose $\delta_3 = 1 + |L|$ such that:

$$|f(x)| < 1 + |L|$$

Since $|f(x)|$ is bounded. Choose $\delta_4 = \epsilon|N|/4(1 + |L|)$, it follows that:

$$|f(x)||N - M| < \frac{\epsilon|N|}{4} \Rightarrow |N - M| < \frac{\epsilon|N|}{4(1 + |L|)}$$

Now choose $\delta = \min \left(\frac{2}{|N|}, \quad \frac{\epsilon|N|}{4|M|}, \quad 1 + |L|, \quad \frac{\epsilon|N|}{4(1+|L|)} \right)$ such that:

$$\frac{|f(x)||N - M| + |M||f(x) - g(x)|}{|g(x)N|} < \left[(1 + |L|) \cdot \frac{\epsilon|N|}{4(1 + |L|)} + |M| \cdot \frac{\epsilon|N|}{4|M|} \right] \cdot \frac{2}{|N|}$$

Simplifying the right side:

$$= \left[\frac{\epsilon|N|}{4} + \frac{\epsilon|N|}{4} \right] \cdot \frac{2}{|N|} = \frac{\epsilon|N|}{2} \cdot \frac{2}{|N|} = \epsilon$$

Hence, for every $\epsilon > 0$, there exists a $\delta > 0$ such that if $0 < |x - c| < \delta$, then:

$$\left| \frac{f(x)}{g(x)} - \frac{L}{M} \right| < \epsilon$$

Thus, we conclude:

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{L}{M}$$

□