

Aprendizaje Reforzado

David Peña Peralta

2024-02-09

Table of contents

Preface	3
1 Introduction	6
2 Summary	7
References	8

Preface

This is a Quarto book.

To learn more about Quarto books visit <https://quarto.org/docs/books>.

```
import numpy as np
```

```
x = 10
```

% Vista preliminar cuerpo

$$\begin{aligned}\nu(x) &= \max_{a \in A(x)} \{c(x, a) + \nu(f(x, a))\} \\ &= \max_{a \in A(x)} \{a^2 + x^2 + E[\nu(x + a + \xi)]\}\end{aligned}$$

Para $\nu(x) = ax^2 + b$

$$\begin{aligned}\nu(x) &= \max_{a \in A(x)} \{c(x, a) + \beta E[\nu(f(x, a))]\} \\ &= \max_{a \in A(x)} \{A^2 + x^2 + \beta(E[a(f^2(x, a))] + b)\} \\ &= \max_{a \in A(x)} \{A^2 + x^2 + \beta(aE[f^2(x, a)] + b)\}\end{aligned}$$

Notemos que

$$\begin{aligned}E[f^2(x, a)] &= E[(x + A + \xi)^2] \\ &= E[x^2 + A^2 + \xi^2 + 2xA + 2x\xi + 2\xi A] \\ &= x^2 + A^2 + E[\xi^2] + 2xA + 2xE[\xi] + 2AE[\xi] \\ &= x^2 + A^2 + d + 2xA\end{aligned}$$

Entonces

$$\begin{aligned}
ax^2 + b &= \max_{a \in A(x)} \{A^2 + x^2 + \beta [a(x^2 + A^2 + d + 2xA) + b]\} \\
&= \max_{a \in A(x)} \{A^2 + x^2 + \beta a(x^2 + A^2 + d + 2xA) + \beta b\} \\
&= \max_{a \in A(x)} \{A^2 + x^2 + \beta ax^2 + \beta aA^2 + \beta ad + 2\beta axA + \beta b\} \\
&= \max_{a \in A(x)} \{A^2(\beta a + 1) + 2\beta axA + x^2 + \beta ax^2 + \beta ad + \beta b\}
\end{aligned}$$

Definimos

$$w(x, A) = A^2(\beta a + 1) + 2\beta axA + x^2 + \beta ax^2 + \beta ad + \beta b,$$

entonces

$$\partial_A w = 2A(\beta a + 1) + 2\beta ax.$$

Si $\partial_A w = 0$, entonces

$$A = -\frac{\beta ax}{\beta a + 1}$$

Entonces

$$\begin{aligned}
\nu(x) &= (\beta ax)^2 - 2\frac{(\beta ax)^2}{\beta a + 1} + x^2 + \beta ax^2 + \beta ad + \beta b \\
&= x^2 \left([\beta a]^2 - 2\frac{(\beta a)^2}{\beta a + 1} + 1 + \beta a \right) + \beta ad + \beta b
\end{aligned}$$

Entonces

$$\begin{aligned}
a &= [\beta a]^2 - 2\frac{(\beta a)^2}{\beta a + 1} + 1 + \beta a \\
b &= \beta ad + \beta b,
\end{aligned}$$

de forma rapida

$$b = \frac{\beta ad}{1 - \beta},$$

entonces queda pendiente calcular a

$$a = [\beta a]^2 - 2 \frac{(\beta a)^2}{\beta a + 1} + 1 + \beta a.$$

$$\begin{aligned} 0 &= (\beta a)^2 \left(1 - \frac{2}{\beta a + 1} \right) + 1 + (\beta - 1) a \\ &= (\beta a)^2 (\beta a + 1 - 2) + \beta a + 1 + (a\beta - a) (\beta a + 1) \\ &= (\beta a)^2 (\beta a - 1) + \beta a + 1 + [(a\beta)^2 + a\beta - \beta a^2 - a] \\ &= (\beta a)^3 + 2a\beta + 1 - \beta a^2 - a \\ &= \beta^3 a^3 - \beta a^2 + (2\beta - 1) a + 1 \end{aligned}$$

1 Introduction

This is a book created from markdown and executable code.

See Knuth (1984) for additional discussion of literate programming.

2 Summary

In summary, this book has no content whatsoever.

References

Knuth, Donald E. 1984. “Literate Programming.” *Comput. J.* 27 (2): 97–111. <https://doi.org/10.1093/comjnl/27.2.97>.