





S17 Figure (page 1 of 4). Training response against phenotype, all legs, Reactome. All 38 Reactome set−outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Nominal p, uncorrected: chance alone returns 86.0 of the 4 x 430 Reactome set−outcome pairs. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.




S17 Figure (page 2 of 4). Training response against phenotype, all legs, Reactome. All 38 Reactome set−outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Nominal p, uncorrected: chance alone returns 86.0 of the 4 x 430 Reactome set−outcome pairs. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.




S17 Figure (page 3 of 4). Training response against phenotype, all legs, Reactome. All 38 Reactome set−outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Nominal p, uncorrected: chance alone returns 86.0 of the 4 x 430 Reactome set−outcome pairs. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.




S17 Figure (page 4 of 4). Training response against phenotype, all legs, Reactome. All 38 Reactome set−outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Nominal p, uncorrected: chance alone returns 86.0 of the 4 x 430 Reactome set−outcome pairs. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.




S18 Figure (page 1 of 4). BFR minus HLRT, within participant, Reactome. All 44 Reactome set−outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Nominal p, uncorrected: chance alone returns 86.0 of the 4 x 430 Reactome set−outcome pairs. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within collection and outcome. Data: set_association sheet of 06_set_association.xlsx.



S18 Figure (page 2 of 4). BFR minus HLRT, within participant, Reactome. All 44 Reactome set−outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Nominal p, uncorrected: chance alone returns 86.0 of the 4 x 430 Reactome set−outcome pairs. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within collection and outcome. Data: set_association sheet of 06_set_association.xlsx.



S18 Figure (page 3 of 4). BFR minus HLRT, within participant, Reactome. All 44 Reactome set−outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Nominal p, uncorrected: chance alone returns 86.0 of the 4 x 430 Reactome set−outcome pairs. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within collection and outcome. Data: set_association sheet of 06_set_association.xlsx.



S18 Figure (page 4 of 4). BFR minus HLRT, within participant, Reactome. All 44 Reactome set−outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Nominal p, uncorrected: chance alone returns 86.0 of the 4 x 430 Reactome set−outcome pairs. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within collection and outcome. Data: set_association sheet of 06_set_association.xlsx.