





S23 Figure. Pathways by leg, GOBP. Colour is NES, dot size the share of the pathway in its leading edge. Pathways chosen from set_tests (01_run_fgsea_and_fry.xlsx): the collapse−surviving fgsea sets at FDR < 0.05 in either training contrast. NES and leading edges from the compareCluster run.



S24 Figure. Enrichment map, GOBP. Nodes are pathways, pies split by leg, joined when their proteins overlap. Pathways chosen from set_tests (01_run_fgsea_and_fry.xlsx): the collapse−surviving fgsea sets at FDR < 0.05 in either training contrast. NES and leading edges from the compareCluster run.



S25 Figure. Pathway tree, GOBP. Pathways clustered by shared proteins, each cluster named by its most frequent words. Pathways chosen from set_tests (01_run_fgsea_and_fry.xlsx): the collapse−surviving fgsea sets at FDR < 0.05 in either training contrast. NES and leading edges from the compareCluster run.



S26 Figure. Pathways and their proteins, GOBP. The five strongest pathways and their leading−edge proteins; pies show which leg each protein leads in. Shared proteins are named in S21 of 07_leading_edge. Pathways chosen from set_tests (01_run_fgsea_and_fry.xlsx): the collapse−surviving fgsea sets at FDR < 0.05 in either training contrast. NES and leading edges from the compareCluster run.