




BFR
HLRT
KEGG_Legacy: Lysine Degradation
vs vl_csa r = +0.38 p 0.002 q 0.19
BFR n=32 r=+0.27 p=0.135
HLRT n=33 r=+0.49 p=0.004
KEGG_Legacy: Colorectal Cancer
vs rf_echo r = −0.35 p 0.0046 q 0.43
BFR n=32 r=−0.50 p=0.004
HLRT n=33 r=−0.25 p=0.154
KEGG_Legacy: WNT Signaling Pathway
vs rf_echo r = −0.31 p 0.011 q 0.47
BFR n=32 r=−0.42 p=0.016
HLRT n=33 r=−0.22 p=0.216
KEGG_Legacy: Chemokine Signaling Pathway
vs rf_echo r = −0.29 p 0.017 q 0.47
BFR n=32 r=−0.32 p=0.078
HLRT n=33 r=−0.29 p=0.107
KEGG_Legacy: Hypertrophic Cardiomyopathy Hcm
vs rf_echo r = +0.29 p 0.022 q 0.47
BFR n=32 r=+0.41 p=0.021
HLRT n=33 r=+0.13 p=0.479
KEGG_Legacy: Arrhythmogenic Right Ventricular Cardiomyopathy Arvc
vs vl_csa r = −0.28 p 0.023 q 0.41
BFR n=32 r=−0.13 p=0.476
HLRT n=33 r=−0.40 p=0.022
KEGG_Legacy: Aminoacyl tRNA Biosynthesis
vs vl_csa r = +0.28 p 0.025 q 0.41
BFR n=32 r=+0.25 p=0.168
HLRT n=33 r=+0.34 p=0.055
KEGG_Legacy: Pyruvate Metabolism
vs vl_csa r = +0.28 p 0.026 q 0.41
BFR n=32 r=+0.16 p=0.384
HLRT n=33 r=+0.39 p=0.024
KEGG_Legacy: Adherens Junction
vs vl_csa r = −0.28 p 0.026 q 0.41
BFR n=32 r=−0.27 p=0.131
HLRT n=33 r=−0.36 p=0.039
KEGG_Legacy: Cysteine And Methionine Metabolism
vs vl_csa r = +0.28 p 0.026 q 0.41
BFR n=32 r=+0.22 p=0.226
HLRT n=33 r=+0.35 p=0.044
KEGG_Legacy: TGF Beta Signaling Pathway
vs rf_echo r = −0.27 p 0.027 q 0.47
BFR n=32 r=−0.20 p=0.271
HLRT n=33 r=−0.31 p=0.083
KEGG_Legacy: Purine Metabolism
vs rf_csa r = −0.26 p 0.037 q 0.91
BFR n=32 r=−0.37 p=0.038
HLRT n=33 r=−0.14 p=0.437
change in phenotype
change in set score, T2 − T1, one point per leg

S11. Pooled association, all legs, KEGG_Legacy (page 2 of 2)



S12. Differential association, BFR minus HLRT, KEGG_Legacy (page 1 of 2)




BFR − HLRT
KEGG_Legacy: Colorectal Cancer
vs rf_echo r = −0.38 p 0.035 q 0.33
KEGG_Legacy: Apoptosis
vs rf_csa r = −0.37 p 0.037 q 0.87
KEGG_Legacy: Long Term Potentiation
vs rf_echo r = −0.37 p 0.037 q 0.33
KEGG_Legacy: B Cell Receptor Signaling Pathway
vs rf_echo r = −0.37 p 0.037 q 0.33
KEGG_Legacy: Tight Junction
vs rf_echo r = +0.37 p 0.037 q 0.33
KEGG_Legacy: T Cell Receptor Signaling Pathway
vs rf_echo r = −0.37 p 0.039 q 0.33
KEGG_Legacy: Dilated Cardiomyopathy
vs vl_echo r = +0.37 p 0.04 q 0.78
KEGG_Legacy: Starch And Sucrose Metabolism
vs vl_echo r = −0.36 p 0.041 q 0.78
KEGG_Legacy: Amyotrophic Lateral Sclerosis Als
vs rf_csa r = −0.36 p 0.043 q 0.87
KEGG_Legacy: Glioma
vs rf_echo r = −0.36 p 0.043 q 0.34
KEGG_Legacy: TGF Beta Signaling Pathway
vs vl_csa r = +0.36 p 0.046 q 0.98
change in phenotype, BFR minus HLRT
change in set score, T2 − T1, BFR minus HLRT, one point per participant