





BFR
HLRT
Reactome: Mitochondrial tRNA Aminoacylation
vs vl_csa r = +0.38 p 0.0017 q 0.74
BFR n=32 r=+0.37 p=0.038
HLRT n=33 r=+0.42 p=0.016
Reactome: Cytoprotection By HMOX1
vs vl_echo r = −0.37 p 0.0026 q 0.96
BFR n=32 r=−0.19 p=0.306
HLRT n=33 r=−0.56 p=0.001
Reactome: Sars Cov 2 Infection
vs rf_echo r = +0.35 p 0.004 q 0.97
BFR n=32 r=+0.42 p=0.019
HLRT n=33 r=+0.32 p=0.069
Reactome: Sars Cov 1 Infection
vs rf_echo r = +0.33 p 0.0067 q 0.97
BFR n=32 r=+0.36 p=0.042
HLRT n=33 r=+0.36 p=0.041
Reactome: Mitochondrial tRNA Aminoacylation
vs rf_csa r = +0.33 p 0.0067 q 1
BFR n=32 r=+0.35 p=0.051
HLRT n=33 r=+0.32 p=0.067
Reactome: DNA Repair
vs rf_csa r = −0.33 p 0.0074 q 1
BFR n=32 r=−0.28 p=0.123
HLRT n=33 r=−0.35 p=0.049
Reactome: Regulation Of Hsf1 Mediated Heat Shock Response
vs rf_echo r = −0.33 p 0.0077 q 0.97
BFR n=32 r=−0.45 p=0.011
HLRT n=33 r=−0.22 p=0.217
Reactome: Rhoq GTPase Cycle
vs rf_echo r = +0.32 p 0.01 q 0.97
BFR n=32 r=+0.18 p=0.328
HLRT n=33 r=+0.49 p=0.004
Reactome: Mitochondrial Translation
vs rf_csa r = +0.31 p 0.012 q 1
BFR n=32 r=+0.29 p=0.104
HLRT n=33 r=+0.30 p=0.089
Reactome: Protein Ubiquitination
vs vl_echo r = +0.31 p 0.012 q 0.96
BFR n=32 r=+0.33 p=0.066
HLRT n=33 r=+0.31 p=0.078
Reactome: Translation
vs rf_csa r = +0.31 p 0.012 q 1
BFR n=32 r=+0.33 p=0.064
HLRT n=33 r=+0.30 p=0.091
Reactome: Mitochondrial Translation Elongation
vs rf_csa r = +0.29 p 0.018 q 1
BFR n=32 r=+0.29 p=0.108
HLRT n=33 r=+0.28 p=0.114
change in phenotype
change in set score, T2 − T1, one point per leg



BFR
HLRT
Reactome: Sars Cov 1 Modulates Host Translation Machinery
vs rf_echo r = +0.29 p 0.018 q 0.97
BFR n=32 r=+0.33 p=0.062
HLRT n=33 r=+0.29 p=0.097
Reactome: Cellular Response To Heat Stress
vs rf_echo r = −0.29 p 0.02 q 0.97
BFR n=32 r=−0.32 p=0.078
HLRT n=33 r=−0.23 p=0.197
Reactome: Sars Cov Infections
vs rf_echo r = +0.28 p 0.023 q 0.97
BFR n=32 r=+0.37 p=0.037
HLRT n=33 r=+0.21 p=0.251
Reactome: Metabolism Of Steroids
vs vl_echo r = −0.28 p 0.024 q 0.96
BFR n=32 r=−0.30 p=0.097
HLRT n=33 r=−0.36 p=0.039
Reactome: tRNA Aminoacylation
vs vl_csa r = +0.28 p 0.026 q 1
BFR n=32 r=+0.20 p=0.267
HLRT n=33 r=+0.34 p=0.056
Reactome: Peroxisomal Protein Import
vs vl_echo r = +0.28 p 0.026 q 0.96
BFR n=32 r=+0.25 p=0.172
HLRT n=33 r=+0.28 p=0.112
Reactome: Clathrin Mediated Endocytosis
vs vl_echo r = −0.28 p 0.027 q 0.96
BFR n=32 r=−0.46 p=0.009
HLRT n=33 r=−0.13 p=0.486
Reactome: Antimicrobial Mechanism Of IFN Stimulated Genes
vs vl_echo r = +0.27 p 0.028 q 0.96
BFR n=32 r=+0.17 p=0.341
HLRT n=33 r=+0.47 p=0.006
Reactome: Iron Uptake And Transport
vs vl_csa r = +0.27 p 0.032 q 1
BFR n=32 r=+0.06 p=0.746
HLRT n=33 r=+0.49 p=0.004
Reactome: Sars Cov 2 Modulates Host Translation Machinery
vs rf_echo r = +0.27 p 0.033 q 0.97
BFR n=32 r=+0.32 p=0.075
HLRT n=33 r=+0.21 p=0.242
Reactome: Late Endosomal Microautophagy
vs rf_csa r = −0.26 p 0.034 q 1
BFR n=32 r=−0.25 p=0.160
HLRT n=33 r=−0.22 p=0.212
Reactome: Metabolism Of Steroids
vs vl_csa r = +0.26 p 0.035 q 1
BFR n=32 r=+0.09 p=0.637
HLRT n=33 r=+0.48 p=0.004
change in phenotype
change in set score, T2 − T1, one point per leg


BFR
HLRT
Reactome: Cytoprotection By HMOX1
vs rf_csa r = +0.26 p 0.039 q 1
BFR n=32 r=+0.11 p=0.549
HLRT n=33 r=+0.33 p=0.058
Reactome: Sars Cov 1 Host Interactions
vs rf_echo r = +0.26 p 0.039 q 0.97
BFR n=32 r=+0.39 p=0.028
HLRT n=33 r=+0.19 p=0.291
Reactome: Response To Elevated Platelet Cytosolic Ca2
vs vl_echo r = +0.26 p 0.04 q 0.96
BFR n=32 r=+0.08 p=0.677
HLRT n=33 r=+0.41 p=0.018
Reactome: Cdc42 GTPase Cycle
vs rf_echo r = +0.26 p 0.04 q 0.97
BFR n=32 r=+0.02 p=0.894
HLRT n=33 r=+0.47 p=0.006
Reactome: Deubiquitination
vs rf_echo r = −0.26 p 0.04 q 0.97
BFR n=32 r=−0.59 p=0.000
HLRT n=33 r=+0.09 p=0.608
Reactome: Rab Gefs Exchange GTP For GDP On Rabs
vs vl_echo r = −0.26 p 0.04 q 0.96
BFR n=32 r=−0.48 p=0.006
HLRT n=33 r=−0.03 p=0.848
Reactome: Formation Of The Dystrophin Glycoprotein Complex DGC
vs vl_csa r = −0.25 p 0.041 q 1
BFR n=32 r=−0.19 p=0.289
HLRT n=33 r=−0.28 p=0.116
Reactome: Signaling By Retinoic Acid
vs vl_echo r = +0.25 p 0.043 q 0.96
BFR n=32 r=+0.12 p=0.498
HLRT n=33 r=+0.38 p=0.029
Reactome: Intra Golgi And Retrograde Golgi To Er Traffic
vs vl_csa r = +0.25 p 0.043 q 1
BFR n=32 r=−0.06 p=0.757
HLRT n=33 r=+0.53 p=0.002
Reactome: Activation Of The mRNA Upon Binding Of The Cap Binding Complex And Eifs And Subsequent Binding To 43s
vs rf_echo r = +0.25 p 0.046 q 0.97
BFR n=32 r=+0.30 p=0.091
HLRT n=33 r=+0.25 p=0.159
Reactome: G2 M DNA Damage Checkpoint
vs rf_csa r = −0.25 p 0.046 q 1
BFR n=32 r=−0.29 p=0.103
HLRT n=33 r=−0.19 p=0.290
Reactome: Rab Gefs Exchange GTP For GDP On Rabs
vs vl_csa r = +0.25 p 0.047 q 1
BFR n=32 r=+0.12 p=0.494
HLRT n=33 r=+0.38 p=0.029
change in phenotype
change in set score, T2 − T1, one point per leg



BFR
HLRT
Reactome: Cell Surface Interactions At The Vascular Wall
vs vl_csa r = −0.25 p 0.047 q 1
BFR n=32 r=−0.33 p=0.062
HLRT n=33 r=−0.18 p=0.321
Reactome: Class I Mhc Mediated Antigen Processing Presentation
vs vl_echo r = +0.25 p 0.047 q 0.96
BFR n=32 r=+0.27 p=0.129
HLRT n=33 r=+0.20 p=0.275
change in phenotype
change in set score, T2 − T1, one point per leg



BFR − HLRT
Reactome: Copii Mediated Vesicle Transport
vs vl_echo r = −0.46 p 0.0081 q 0.96
Reactome: Slc Mediated Transmembrane Transport
vs rf_csa r = −0.46 p 0.0083 q 0.84
Reactome: Cargo Concentration In The Er
vs vl_echo r = −0.45 p 0.01 q 0.96
Reactome: Cellular Senescence
vs vl_echo r = +0.43 p 0.015 q 0.96
Reactome: Esr Mediated Signaling
vs rf_echo r = −0.43 p 0.016 q 0.93
Reactome: Arachidonate Metabolism
vs vl_echo r = −0.42 p 0.018 q 0.96
Reactome: Rhov GTPase Cycle
vs vl_csa r = −0.42 p 0.018 q 0.99
Reactome: Signaling By Fgfr
vs vl_echo r = +0.41 p 0.02 q 0.96
Reactome: Er To Golgi Anterograde Transport
vs vl_echo r = −0.41 p 0.02 q 0.96
Reactome: Transport To The Golgi And Subsequent Modification
vs vl_echo r = −0.41 p 0.02 q 0.96
Reactome: Cooperation Of Pdcl Phlp1 And Tric Cct In G Protein Beta Folding
vs vl_echo r = −0.41 p 0.021 q 0.96
Reactome: G Alpha Q Signalling Events
vs rf_echo r = −0.41 p 0.022 q 0.93
change in phenotype, BFR minus HLRT
change in set score, T2 − T1, BFR minus HLRT, one point per participant

S12. Differential association, BFR minus HLRT, Reactome (page 2 of 4)



S12. Differential association, BFR minus HLRT, Reactome (page 3 of 4)




BFR − HLRT
Reactome: Signaling By WNT
vs rf_echo r = −0.36 p 0.045 q 0.93
Reactome: Metabolism Of Steroids
vs vl_echo r = −0.36 p 0.046 q 0.96
Reactome: Signaling By Fgfr2
vs vl_echo r = +0.36 p 0.047 q 0.96
Reactome: Regulation Of Hsf1 Mediated Heat Shock Response
vs rf_echo r = −0.36 p 0.047 q 0.93
Reactome: Cytosolic tRNA Aminoacylation
vs vl_csa r = −0.35 p 0.047 q 0.99
Reactome: Striated Muscle Contraction
vs rf_echo r = +0.35 p 0.048 q 0.93
Reactome: Fc Epsilon Receptor Fceri Signaling
vs rf_echo r = −0.35 p 0.05 q 0.93
change in phenotype, BFR minus HLRT
change in set score, T2 − T1, BFR minus HLRT, one point per participant