



A

BFR_Post-Pre, 10 of 88 survivors




B

HLRT_Post-Pre, 10 of 68 survivors




C

Modality_x_Time_Interaction, 10 of 10 survivors




D

BFR_Post-HLRT_Post, 10 of 52 survivors




S1 Figure. Top fgsea sets, all collections. (A) BFR_Post-Pre. (B) HLRT_Post-Pre. (C) Modality_x_Time_Interaction. (D) BFR_Post-HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is the collection. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.


A




B



C




D




S2 Figure. Top fgsea sets, Hallmark. (A) BFR_Post-Pre. (B) HLRT_Post-Pre. (C) Modality_x_Time_Interaction. (D) BFR_Post-HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is —log₁₀ FDR. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.


A



B




C




S3 Figure. Top fgsea sets, Reactome. (A) BFR_Post-Pre. (B) HLRT_Post-Pre. (C) BFR_Post-HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is $-\log_{10}$ FDR. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.


A



B




C



S4 Figure. Top fgsea sets, KEGG_Legacy. (A) BFR_Post-Pre. (B) HLRT_Post-Pre. (C) BFR_Post-HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is $-\log_{10}$ FDR. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.


A

BFR_Post-Pre, 10 of 50 survivors




B

HLRT_Post-Pre, 10 of 39 survivors




C

Modality_x_Time_Interaction, 7 of 7 survivors




D

BFR_Post-HLRT_Post, 10 of 27 survivors




S5 Figure. Top fgsea sets, GOBP. (A) BFR_Post-Pre. (B) HLRT_Post-Pre. (C) Modality_x_Time_Interaction. (D) BFR_Post-HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is $-\log_{10}$ FDR. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.


A




B




C



D



S6 Figure. Top fgsea sets, GO_Slim. (A) BFR_Post-Pre. (B) HLRT_Post-Pre. (C) Modality_x_Time_Interaction. (D) BFR_Post-HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is $-\log_{10}$ FDR. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.



S7 Figure. Collapse, before and after. Sets at FDR 0.05 per collection and contrast, before and after collapsePathways, which keeps a set only if it stands alone given a stronger set's leading edge. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.