







S16 Figure (page 1 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 2 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 3 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 4 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 5 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 6 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 7 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 8 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 9 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 10 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 11 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S16 Figure (page 12 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 13 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 14 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



change in phenotype















S16 Figure (page 15 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S16 Figure (page 16 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




change in phenotype




change in set score, T2 – T1, one point per leg


S16 Figure (page 17 of 17). Training response against phenotype, all legs, GOBP. All 204 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 1 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set–outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 2 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 3 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 4 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 5 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S17 Figure (page 6 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 7 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 8 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



change in phenotype















S17 Figure (page 9 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S17 Figure (page 10 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




change in phenotype




change in set score, T2 - T1, one point per leg


S17 Figure (page 11 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 12 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S17 Figure (page 13 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S17 Figure (page 14 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S17 Figure (page 15 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




change in phenotype




S17 Figure (page 16 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




change in phenotype



S17 Figure (page 17 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S17 Figure (page 18 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S17 Figure (page 19 of 19). BFR minus HLRT, within participant, GOBP. All 227 GOBP set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 68.3 of the 1366 GOBP sets. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.