



Training response against phenotype, all legs, rf_echo (page 1 of 10)




Training response against phenotype, all legs, rf_echo (page 2 of 10)






BFR
HLRT
NOS1 P29475
vs rf_echo t = −2.57 p 0.012 q 1
BFR n=32 r=−0.02
HLRT n=33 r=−0.41
CLYBL Q8N0X4
vs rf_echo t = −2.57 p 0.012 q 1
BFR n=32 r=−0.24
HLRT n=33 r=−0.24
SUOX P51687
vs rf_echo t = −2.56 p 0.013 q 1
BFR n=32 r=−0.32
HLRT n=33 r=−0.31
PSMC3 P17980
vs rf_echo t = +2.55 p 0.013 q 1
BFR n=32 r=−0.12
HLRT n=33 r=+0.56
UBE2V1 Q13404
vs rf_echo t = +2.54 p 0.013 q 1
BFR n=32 r=+0.26
HLRT n=33 r=+0.16
HEBP2 Q9Y5Z4
vs rf_echo t = +2.54 p 0.013 q 1
BFR n=32 r=+0.10
HLRT n=33 r=+0.37
ANP32A P39687
vs rf_echo t = +2.53 p 0.014 q 1
BFR n=32 r=+0.27
HLRT n=33 r=+0.42
CD151 P48509
vs rf_echo t = −2.50 p 0.015 q 1
BFR n=32 r=−0.30
HLRT n=33 r=−0.39
SHMT2 P34897
vs rf_echo t = −2.48 p 0.016 q 1
BFR n=32 r=−0.20
HLRT n=33 r=−0.36
DHRS7C A6NNS2
vs rf_echo t = +2.48 p 0.016 q 1
BFR n=32 r=+0.14
HLRT n=33 r=+0.45
COPS6 Q7L5N1
vs rf_echo t = −2.48 p 0.016 q 1
BFR n=32 r=−0.34
HLRT n=33 r=−0.43
LAMTOR1 Q6IAA8
vs rf_echo t = +2.48 p 0.016 q 1
BFR n=32 r=+0.19
HLRT n=33 r=+0.31
change in phenotype
change in protein, log2, T2 − T1, one point per leg



BFR
HLRT
UBE2N P61088
vs rf_echo t = +2.47 p 0.016 q 1
BFR n=32 r=+0.01
HLRT n=33 r=+0.43
CLASP1 Q7Z460
vs rf_echo t = +2.47 p 0.016 q 1
BFR n=32 r=+0.24
HLRT n=33 r=+0.35
HAX1 O00165
vs rf_echo t = −2.46 p 0.016 q 1
BFR n=32 r=−0.01
HLRT n=33 r=−0.61
USP14 P54578
vs rf_echo t = +2.44 p 0.017 q 1
BFR n=32 r=+0.10
HLRT n=33 r=+0.40
SYAP1 Q96A49
vs rf_echo t = +2.41 p 0.019 q 1
BFR n=32 r=+0.13
HLRT n=33 r=+0.34
FEN1 P39748
vs rf_echo t = +2.40 p 0.019 q 1
BFR n=32 r=+0.37
HLRT n=33 r=+0.15
SRSF1 Q07955
vs rf_echo t = +2.37 p 0.021 q 1
BFR n=32 r=+0.28
HLRT n=33 r=+0.24
DARS1 P14868
vs rf_echo t = −2.35 p 0.021 q 1
BFR n=32 r=−0.17
HLRT n=33 r=−0.45
ISCU Q9H1K1
vs rf_echo t = +2.35 p 0.022 q 1
BFR n=32 r=+0.07
HLRT n=33 r=+0.27
PSMB4 P28070
vs rf_echo t = −2.35 p 0.022 q 1
BFR n=32 r=−0.24
HLRT n=33 r=−0.43
RPL22 P35268
vs rf_echo t = +2.34 p 0.022 q 1
BFR n=32 r=+0.44
HLRT n=33 r=+0.24
GNA11 P29992
vs rf_echo t = +2.34 p 0.022 q 1
BFR n=32 r=+0.33
HLRT n=33 r=+0.35
change in phenotype
change in protein, log2, T2 − T1, one point per leg



BFR
HLRT
ACE P12821
vs rf_echo t = +2.32 p 0.023 q 1
BFR n=32 r=+0.31
HLRT n=33 r=+0.23
AKR1C2 P52895
vs rf_echo t = −2.31 p 0.024 q 1
BFR n=32 r=−0.35
HLRT n=33 r=−0.26
APIP Q96GX9
vs rf_echo t = +2.30 p 0.025 q 1
BFR n=32 r=+0.08
HLRT n=33 r=+0.31
NUBPL Q8TB37
vs rf_echo t = −2.29 p 0.025 q 1
BFR n=32 r=+0.18
HLRT n=33 r=−0.46
OTUD4 Q01804
vs rf_echo t = −2.28 p 0.026 q 1
BFR n=32 r=−0.18
HLRT n=33 r=−0.41
NDUFA1 O15239
vs rf_echo t = −2.28 p 0.026 q 1
BFR n=32 r=−0.12
HLRT n=33 r=−0.28
MPC2 O95563
vs rf_echo t = −2.27 p 0.026 q 1
BFR n=32 r=−0.03
HLRT n=33 r=−0.49
COPZ1 P61923
vs rf_echo t = +2.27 p 0.027 q 1
BFR n=32 r=+0.09
HLRT n=33 r=+0.32
HSDL1 Q3SXM5
vs rf_echo t = −2.26 p 0.027 q 1
BFR n=32 r=+0.05
HLRT n=33 r=−0.53
ELOC Q15369
vs rf_echo t = +2.25 p 0.028 q 1
BFR n=32 r=+0.07
HLRT n=33 r=+0.33
ETFA P13804
vs rf_echo t = −2.24 p 0.028 q 1
BFR n=32 r=−0.23
HLRT n=33 r=−0.35
RPS9 P46781
vs rf_echo t = +2.24 p 0.028 q 1
BFR n=32 r=+0.26
HLRT n=33 r=+0.30
change in phenotype
change in protein, log2, T2 − T1, one point per leg



BFR
HLRT
RTCB Q9Y3I0
vs rf_echo t = +2.23 p 0.029 q 1
BFR n=32 r=+0.33
HLRT n=33 r=+0.42
PGK2 P07205
vs rf_echo t = +2.23 p 0.03 q 1
BFR n=32 r=+0.26
HLRT n=33 r=+0.22
LRRC57 Q8N9N7
vs rf_echo t = +2.22 p 0.03 q 1
BFR n=32 r=+0.32
HLRT n=33 r=+0.22
UBE2H P62256
vs rf_echo t = −2.21 p 0.03 q 1
BFR n=32 r=−0.22
HLRT n=33 r=−0.33
FKBP4 Q02790
vs rf_echo t = −2.21 p 0.031 q 1
BFR n=32 r=−0.18
HLRT n=33 r=−0.54
UFL1 O94874
vs rf_echo t = +2.20 p 0.031 q 1
BFR n=32 r=+0.24
HLRT n=33 r=+0.40
MTHFS P49914
vs rf_echo t = −2.20 p 0.031 q 1
BFR n=32 r=−0.31
HLRT n=33 r=−0.22
BRCC3 P46736
vs rf_echo t = −2.19 p 0.032 q 1
BFR n=32 r=−0.21
HLRT n=33 r=−0.29
PTCD3 Q96EY7
vs rf_echo t = −2.19 p 0.032 q 1
BFR n=32 r=−0.13
HLRT n=33 r=−0.40
MRPS18B Q9Y676
vs rf_echo t = −2.18 p 0.033 q 1
BFR n=32 r=−0.10
HLRT n=33 r=−0.18
RRAD P55042
vs rf_echo t = +2.17 p 0.033 q 1
BFR n=32 r=+0.15
HLRT n=33 r=+0.40
CNIH4 Q9P003
vs rf_echo t = +2.17 p 0.034 q 1
BFR n=32 r=+0.16
HLRT n=33 r=+0.40
change in phenotype
change in protein, log2, T2 − T1, one point per leg



BFR
HLRT
SMS P52788
vs rf_echo t = −2.16 p 0.034 q 1
BFR n=32 r=−0.25
HLRT n=33 r=−0.28
ITGA1 P56199
vs rf_echo t = −2.15 p 0.035 q 1
BFR n=32 r=−0.33
HLRT n=33 r=−0.00
UBE2M P61081
vs rf_echo t = −2.15 p 0.035 q 1
BFR n=32 r=−0.27
HLRT n=33 r=−0.25
CRYL1 Q9Y2S2
vs rf_echo t = −2.15 p 0.035 q 1
BFR n=32 r=−0.12
HLRT n=33 r=−0.27
EIF4A2 Q14240
vs rf_echo t = +2.15 p 0.035 q 1
BFR n=32 r=+0.21
HLRT n=33 r=+0.24
VPS33A Q96AX1
vs rf_echo t = −2.14 p 0.036 q 1
BFR n=32 r=−0.15
HLRT n=33 r=−0.39
CD93 Q9NPY3
vs rf_echo t = −2.14 p 0.036 q 1
BFR n=32 r=−0.14
HLRT n=33 r=−0.24
NRP1 O14786
vs rf_echo t = −2.14 p 0.036 q 1
BFR n=32 r=−0.23
HLRT n=33 r=−0.22
RPS24 P62847
vs rf_echo t = +2.14 p 0.036 q 1
BFR n=32 r=+0.33
HLRT n=33 r=+0.08
HMBS P08397
vs rf_echo t = +2.13 p 0.036 q 1
BFR n=32 r=+0.03
HLRT n=33 r=+0.44
CERT1 Q9Y5P4
vs rf_echo t = +2.13 p 0.037 q 1
BFR n=32 r=+0.01
HLRT n=33 r=+0.34
BIN1 O00499
vs rf_echo t = +2.13 p 0.037 q 1
BFR n=32 r=+0.06
HLRT n=33 r=+0.40
change in phenotype
change in protein, log2, T2 − T1, one point per leg



BFR
HLRT
FKBP5 Q13451
vs rf_echo t = −2.12 p 0.037 q 1
BFR n=32 r=−0.39
HLRT n=33 r=−0.16
PGP A6NDG6
vs rf_echo t = −2.12 p 0.038 q 1
BFR n=32 r=−0.41
HLRT n=33 r=−0.31
RPN1 P04843
vs rf_echo t = +2.12 p 0.038 q 1
BFR n=32 r=+0.24
HLRT n=33 r=+0.22
FBXO7 Q9Y3I1
vs rf_echo t = +2.11 p 0.038 q 1
BFR n=32 r=+0.16
HLRT n=33 r=+0.31
ATP6V1D Q9Y5K8
vs rf_echo t = +2.11 p 0.038 q 1
BFR n=32 r=+0.01
HLRT n=33 r=+0.37
VAPB O95292
vs rf_echo t = +2.11 p 0.039 q 1
BFR n=32 r=+0.41
HLRT n=33 r=+0.18
GNG2 P59768
vs rf_echo t = −2.11 p 0.039 q 1
BFR n=32 r=−0.21
HLRT n=33 r=−0.28
TRAPPC1 Q9Y5R8
vs rf_echo t = −2.10 p 0.039 q 1
BFR n=32 r=−0.43
HLRT n=33 r=−0.21
ATIC P31939
vs rf_echo t = +2.10 p 0.039 q 1
BFR n=32 r=+0.18
HLRT n=33 r=+0.33
RPS21 P63220
vs rf_echo t = +2.10 p 0.04 q 1
BFR n=32 r=+0.18
HLRT n=33 r=+0.24
POC1B Q8TC44
vs rf_echo t = +2.08 p 0.041 q 1
BFR n=32 r=−0.00
HLRT n=33 r=+0.39
EXOC1 Q9NV70
vs rf_echo t = −2.08 p 0.041 q 1
BFR n=32 r=−0.14
HLRT n=33 r=−0.30
change in phenotype
change in protein, log2, T2 − T1, one point per leg



BFR
HLRT
MAPK9 P45984
vs rf_echo t = −2.08 p 0.042 q 1
BFR n=32 r=−0.32
HLRT n=33 r=−0.11
GET4 Q7L5D6
vs rf_echo t = −2.07 p 0.042 q 1
BFR n=32 r=−0.18
HLRT n=33 r=−0.28
HNRNPH2 P55795
vs rf_echo t = +2.07 p 0.042 q 1
BFR n=32 r=+0.05
HLRT n=33 r=+0.41
EMC6 Q9BV81
vs rf_echo t = +2.07 p 0.042 q 1
BFR n=32 r=+0.39
HLRT n=33 r=+0.14
DRG1 Q9Y295
vs rf_echo t = +2.06 p 0.043 q 1
BFR n=32 r=+0.11
HLRT n=33 r=+0.27
BLVRB P30043
vs rf_echo t = +2.05 p 0.044 q 1
BFR n=32 r=+0.17
HLRT n=33 r=+0.23
PCYT1A P49585
vs rf_echo t = −2.05 p 0.044 q 1
BFR n=32 r=−0.13
HLRT n=33 r=−0.37
SRPRB Q9Y5M8
vs rf_echo t = +2.05 p 0.045 q 1
BFR n=32 r=+0.27
HLRT n=33 r=+0.29
RPS16 P62249
vs rf_echo t = +2.04 p 0.045 q 1
BFR n=32 r=+0.21
HLRT n=33 r=+0.42
PFN2 P35080
vs rf_echo t = +2.04 p 0.045 q 1
BFR n=32 r=+0.12
HLRT n=33 r=+0.37
POLDIP2 Q9Y2S7
vs rf_echo t = −2.04 p 0.045 q 1
BFR n=32 r=−0.18
HLRT n=33 r=−0.32
MRPL46 Q9H2W6
vs rf_echo t = −2.04 p 0.045 q 1
BFR n=32 r=−0.07
HLRT n=33 r=−0.29
change in phenotype
change in protein, log2, T2 − T1, one point per leg



BFR
HLRT
COA3 Q9Y2R0
vs rf_echo t = −2.03 p 0.046 q 1
BFR n=32 r=+0.22
HLRT n=33 r=−0.58
LUC7L2 Q9Y383
vs rf_echo t = −2.03 p 0.046 q 1
BFR n=32 r=−0.29
HLRT n=33 r=−0.35
CHCHD7 Q9BUK0
vs rf_echo t = −2.03 p 0.047 q 1
BFR n=32 r=−0.21
HLRT n=33 r=−0.27
HSPD1 P10809
vs rf_echo t = −2.02 p 0.047 q 1
BFR n=32 r=−0.19
HLRT n=33 r=−0.18
PLCL1 Q15111
vs rf_echo t = −2.02 p 0.047 q 1
BFR n=32 r=−0.31
HLRT n=33 r=−0.23
ACAA1 P09110
vs rf_echo t = −2.01 p 0.048 q 1
BFR n=32 r=−0.15
HLRT n=33 r=−0.26
ERGIC1 Q969X5
vs rf_echo t = +2.01 p 0.049 q 1
BFR n=32 r=−0.01
HLRT n=33 r=+0.38
USP19 O94966
vs rf_echo t = −2.01 p 0.049 q 1
BFR n=32 r=−0.32
HLRT n=33 r=−0.37
RALA P11233
vs rf_echo t = −2.00 p 0.049 q 1
BFR n=32 r=+0.02
HLRT n=33 r=−0.48
S100A4 P26447
vs rf_echo t = +2.00 p 0.049 q 1
BFR n=32 r=+0.23
HLRT n=33 r=+0.37
SMTNL1 A8MU46
vs rf_echo t = +2.00 p 0.05 q 1
BFR n=32 r=+0.04
HLRT n=33 r=+0.30
CALU O43852
vs rf_echo t = −2.00 p 0.05 q 1
BFR n=32 r=−0.17
HLRT n=33 r=−0.24
change in phenotype
change in protein, log2, T2 − T1, one point per leg



BFR − HLRT
PA2G4 Q9UQ80
vs rf_echo t = +4.22 p 0.00017 q 0.33
SMS P52788
vs rf_echo t = −4.14 p 0.00022 q 0.33
GNG2 P59768
vs rf_echo t = −3.48 p 0.0014 q 1
SYAP1 Q96A49
vs rf_echo t = +3.34 p 0.002 q 1
CALU O43852
vs rf_echo t = −3.31 p 0.0022 q 1
PPCS Q9HAB8
vs rf_echo t = −3.19 p 0.0031 q 1
RPS21 P63220
vs rf_echo t = +3.05 p 0.0045 q 1
TMEM70 Q9BUB7
vs rf_echo t = −2.95 p 0.0057 q 1
RNH1 P13489
vs rf_echo t = −2.90 p 0.0065 q 1
MT−ND2 P03891
vs rf_echo t = −2.81 p 0.0082 q 1
MAIP1 Q8WWC4
vs rf_echo t = −2.81 p 0.0083 q 1
NANS Q9NR45
vs rf_echo t = −2.79 p 0.0085 q 1
change in phenotype, BFR minus HLRT
change in protein, log2, T2 − T1, BFR minus HLRT, one point per participant



BFR − HLRT
ATPAF1 Q5TC12
vs rf_echo t = −2.79 p 0.0086 q 1
TIMM50 Q3ZCQ8
vs rf_echo t = −2.78 p 0.0088 q 1
RAD23B P54727
vs rf_echo t = +2.76 p 0.0091 q 1
MAOB P27338
vs rf_echo t = +2.76 p 0.0092 q 1
DHRS7 Q9Y394
vs rf_echo t = −2.75 p 0.0094 q 1
TIMM8B Q9Y5J9
vs rf_echo t = +2.74 p 0.0098 q 1
EMD P50402
vs rf_echo t = +2.70 p 0.011 q 1
SPTB P11277
vs rf_echo t = +2.71 p 0.011 q 1
STOM P27105
vs rf_echo t = +2.68 p 0.011 q 1
HSD17B12 Q53GQ0
vs rf_echo t = +2.66 p 0.012 q 1
VAPB O95292
vs rf_echo t = +2.64 p 0.012 q 1
LPIN1 Q14693
vs rf_echo t = −2.64 p 0.012 q 1
change in phenotype, BFR minus HLRT
change in protein, log2, T2 − T1, BFR minus HLRT, one point per participant



BFR − HLRT
RPS25 P62851
vs rf_echo t = +2.64 p 0.012 q 1
SDHAF2 Q9NX18
vs rf_echo t = −2.63 p 0.013 q 1
HEXA P06865
vs rf_echo t = −2.62 p 0.013 q 1
VAMP3 Q15836
vs rf_echo t = −2.60 p 0.014 q 1
DRG1 Q9Y295
vs rf_echo t = +2.57 p 0.015 q 1
COX5A P20674
vs rf_echo t = +2.56 p 0.015 q 1
VDAC1 P21796
vs rf_echo t = +2.55 p 0.016 q 1
PSMG2 Q969U7
vs rf_echo t = +2.53 p 0.016 q 1
BLVRB P30043
vs rf_echo t = +2.52 p 0.017 q 1
SOD1 P00441
vs rf_echo t = −2.49 p 0.018 q 1
EMC2 Q15006
vs rf_echo t = +2.48 p 0.018 q 1
DYNC1LI2 O43237
vs rf_echo t = +2.47 p 0.019 q 1
change in phenotype, BFR minus HLRT
change in protein, log2, T2 − T1, BFR minus HLRT, one point per participant



BFR − HLRT
AKR1C2 P52895
vs rf_echo t = −2.47 p 0.019 q 1
HMBS P08397
vs rf_echo t = +2.46 p 0.019 q 1
HEBP1 Q9NRV9
vs rf_echo t = +2.46 p 0.019 q 1
NDUFB5 O43674
vs rf_echo t = +2.45 p 0.019 q 1
DYNLT1 P63172
vs rf_echo t = −2.44 p 0.02 q 1
CMPK1 P30085
vs rf_echo t = −2.43 p 0.02 q 1
GSN P06396
vs rf_echo t = −2.41 p 0.022 q 1
CIAO2A Q9H5X1
vs rf_echo t = −2.39 p 0.023 q 1
CHMP4B Q9H444
vs rf_echo t = −2.38 p 0.023 q 1
MIPEP Q99797
vs rf_echo t = −2.38 p 0.023 q 1
ARMC1 Q9NVT9
vs rf_echo t = −2.36 p 0.024 q 1
CDV3 Q9UKY7
vs rf_echo t = +2.35 p 0.024 q 1
change in phenotype, BFR minus HLRT
change in protein, log2, T2 − T1, BFR minus HLRT, one point per participant



BFR − HLRT
MAPK3 P27361
vs rf_echo t = −2.36 p 0.025 q 1
CRYL1 Q9Y2S2
vs rf_echo t = −2.34 p 0.026 q 1
KPNA3 O00505
vs rf_echo t = −2.33 p 0.026 q 1
FUS P35637
vs rf_echo t = −2.32 p 0.026 q 1
NSUN4 Q96CB9
vs rf_echo t = −2.32 p 0.027 q 1
ANXA5 P08758
vs rf_echo t = −2.31 p 0.027 q 1
PFDN5 Q99471
vs rf_echo t = −2.29 p 0.028 q 1
SLMAP Q14BN4
vs rf_echo t = +2.27 p 0.03 q 1
ULK3 Q6PHR2
vs rf_echo t = −2.26 p 0.03 q 1
FKBP5 Q13451
vs rf_echo t = −2.26 p 0.03 q 1
KHSRP Q92945
vs rf_echo t = +2.26 p 0.03 q 1
NDUFB10 O96000
vs rf_echo t = +2.25 p 0.031 q 1
change in phenotype, BFR minus HLRT
change in protein, log2, T2 − T1, BFR minus HLRT, one point per participant


BFR − HLRT
MVD P53602
vs rf_echo t = +2.24 p 0.031 q 1
CCDC43 Q96MW1
vs rf_echo t = +2.24 p 0.031 q 1
SRL Q86TD4
vs rf_echo t = +2.24 p 0.032 q 1
LCN2 P80188
vs rf_echo t = +2.24 p 0.032 q 1
TBCEL Q5QJ74
vs rf_echo t = +2.22 p 0.033 q 1
EIF3G O75821
vs rf_echo t = +2.22 p 0.033 q 1
FKBP3 Q00688
vs rf_echo t = +2.22 p 0.033 q 1
NCKAP1 Q9Y2A7
vs rf_echo t = −2.22 p 0.033 q 1
COMT P21964
vs rf_echo t = −2.22 p 0.033 q 1
BABAM2 Q9NXR7
vs rf_echo t = +2.22 p 0.033 q 1
CLASP1 Q7Z460
vs rf_echo t = +2.22 p 0.033 q 1
OCIAD1 Q9NX40
vs rf_echo t = +2.21 p 0.034 q 1
change in phenotype, BFR minus HLRT
change in protein, log2, T2 − T1, BFR minus HLRT, one point per participant


BFR − HLRT
JPH1 Q9HDC5
vs rf_echo t = +2.20 p 0.034 q 1
PSMC2 P35998
vs rf_echo t = +2.20 p 0.034 q 1
GNB2 P62879
vs rf_echo t = −2.20 p 0.035 q 1
ASMTL O95671
vs rf_echo t = −2.20 p 0.035 q 1
PPP3CC P48454
vs rf_echo t = −2.19 p 0.035 q 1
TPP2 P29144
vs rf_echo t = +2.19 p 0.036 q 1
COPS6 Q7L5N1
vs rf_echo t = −2.19 p 0.036 q 1
LDHD Q86WU2
vs rf_echo t = +2.18 p 0.036 q 1
TOLLIP Q9H0E2
vs rf_echo t = −2.18 p 0.036 q 1
IARS1 P41252
vs rf_echo t = −2.17 p 0.037 q 1
DHRS7B Q6IAN0
vs rf_echo t = +2.17 p 0.037 q 1
HSPB6 O14558
vs rf_echo t = −2.17 p 0.037 q 1
change in phenotype, BFR minus HLRT
change in protein, log2, T2 − T1, BFR minus HLRT, one point per participant


BFR − HLRT
MYOC Q99972
vs rf_echo t = +2.16 p 0.038 q 1
PRKAA1 Q13131
vs rf_echo t = +2.15 p 0.039 q 1
STIM1 Q13586
vs rf_echo t = +2.15 p 0.039 q 1
PGP A6NDG6
vs rf_echo t = −2.14 p 0.04 q 1
GCLM P48507
vs rf_echo t = +2.14 p 0.04 q 1
SRP9 P49458
vs rf_echo t = −2.13 p 0.04 q 1
LXN Q9BS40
vs rf_echo t = +2.13 p 0.041 q 1
COQ10A Q96MF6
vs rf_echo t = −2.12 p 0.041 q 1
MT−ATP6 P00846
vs rf_echo t = −2.12 p 0.042 q 1
LMAN2 Q12907
vs rf_echo t = +2.12 p 0.042 q 1
SRP72 O76094
vs rf_echo t = +2.11 p 0.042 q 1
RPS24 P62847
vs rf_echo t = +2.11 p 0.042 q 1
change in phenotype, BFR minus HLRT
change in protein, log2, T2 − T1, BFR minus HLRT, one point per participant


BFR − HLRT
NACA E9PAV3
vs rf_echo t = +2.10 p 0.043 q 1
CPTP Q5TA50
vs rf_echo t = −2.10 p 0.044 q 1
CDK5RAP3 Q96JB5
vs rf_echo t = −2.10 p 0.044 q 1
CD5L O43866
vs rf_echo t = +2.10 p 0.044 q 1
MTARC2 Q969Z3
vs rf_echo t = +2.09 p 0.044 q 1
PPP2R3A Q06190
vs rf_echo t = −2.09 p 0.044 q 1
PGK2 P07205
vs rf_echo t = +2.09 p 0.045 q 1
PSMA3 P25788
vs rf_echo t = +2.08 p 0.045 q 1
FEN1 P39748
vs rf_echo t = +2.07 p 0.046 q 1
UACA Q9BZF9
vs rf_echo t = −2.07 p 0.046 q 1
ELOC Q15369
vs rf_echo t = +2.07 p 0.046 q 1
OTUD6B Q8N6M0
vs rf_echo t = +2.07 p 0.046 q 1
change in phenotype, BFR minus HLRT
change in protein, log2, T2 − T1, BFR minus HLRT, one point per participant


BFR − HLRT
COQ9 O75208
vs rf_echo t = −2.06 p 0.047 q 1
USP19 O94966
vs rf_echo t = −2.06 p 0.047 q 1
ASB11 Q8WXH4
vs rf_echo t = +2.05 p 0.048 q 1
LCLAT1 Q6UWP7
vs rf_echo t = +2.05 p 0.048 q 1
PCBP1 Q15365
vs rf_echo t = +2.05 p 0.049 q 1
APOOL Q6UXV4
vs rf_echo t = +2.04 p 0.049 q 1
LDHAL6B Q9BYZ2
vs rf_echo t = −2.04 p 0.049 q 1
CAND1 Q86VP6
vs rf_echo t = +2.04 p 0.049 q 1
change in phenotype, BFR minus HLRT
change in protein, log2, T2 − T1, BFR minus HLRT, one point per participant