


BFR minus HLRT, within participant


Spearman, differential analysis, 32 paired participants




GOBP: Prostanoid
Metabolism vs rf_echo
 $r = -0.65$ $p = 8.9e-05$




GOBP: Reg. Membrane
Depolarization vs rf_echo
 $r = +0.58$ $p = 0.00063$




GO_Slim: Cytoskel. Org. vs rf_echo
 $r = +0.44$ $p = 0.013$




GO_Slim: Circulatory
System Process vs vl_csa
 $r = -0.44$ $p = 0.013$




Hallmark: Heme
Metabolism vs vl_echo
 $r = +0.47$ $p = 0.007$




Hallmark: UPR vs rf_csa
 $r = +0.41$ $p = 0.02$




KEGG_Legacy: Hypertrophic
Cardiomyopathy
Hcm vs rf_echo
 $r = +0.46$ $p = 0.0081$




KEGG_Legacy: WNT Signaling vs rf_echo
 $r = -0.45$ $p = 0.011$



Reactome: Copii Mediated
Vesicle
Transport vs vl_echo
 $r = -0.52$ $p = 0.0024$




Reactome: Cargo
Concentration
In The Er vs vl_echo
 $r = -0.48$ $p = 0.0055$



change in set score, T2 - T1, one point per leg

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled




Strongest 2 set-outcome pairs per collection of 270 reaching nominal p. Lines are fitted within each arm; the header r is the pooled correlation that selected the panel. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO


Nominal hits relative to chance, by collection

paired Wilcoxon per set, 5 collections, uncorrected p


Pre to post, BFR




Pre to post, HLRT




Baseline BFR vs HLRT (control)



Post BFR vs HLRT




Change, BFR vs HLRT



0 2 4 6 8
observed nominal hits / chance expectation
Bar length is observed nominal hits divided by the count that collection returns under the null. Red clears 1, grey does not

Change, BFR vs HLRT


singscore per set, 32 paired participants, AUC with direction fixed



Strongest 2 sets per collection of 43 reaching nominal p. Red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test

Post BFR vs HLRT


singscore per set, 32 paired participants, AUC with direction fixed



Strongest 2 sets per collection of 94 reaching nominal p. Red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test. Nominal p, uncorrected.

Pre to post, BFR


singscore per set, 32 paired legs, AUC with direction fixed



Strongest 2 sets per collection of 391 reaching nominal p. Red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test. Nominal

Pre to post, HLRT

singscore per set, 33 paired legs, AUC with direction fixed



Strongest 2 sets per collection of 389 reaching nominal p. Red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test. Nominal p, un