





Training response against phenotype, all legs, rf_echo (page 1 of 10). All 113 proteins whose Spearman correlation with rf_echo reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, rf_echo (page 2 of 10). All 113 proteins whose Spearman correlation with rf_echo reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, rf_echo (page 3 of 10). All 113 proteins whose Spearman correlation with rf_echo reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, rf_echo (page 4 of 10). All 113 proteins whose Spearman correlation with rf_echo reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, rf_echo (page 5 of 10). All 113 proteins whose Spearman correlation with rf_echo reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, rf_echo (page 6 of 10). All 113 proteins whose Spearman correlation with rf_echo reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, rf_echo (page 7 of 10). All 113 proteins whose Spearman correlation with rf_echo reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, rf_echo (page 8 of 10). All 113 proteins whose Spearman correlation with rf_echo reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, rf_echo (page 9 of 10). All 113 proteins whose Spearman correlation with rf_echo reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, rf_echo (page 10 of 10). All 113 proteins whose Spearman correlation with rf_echo reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




BFR minus HLRT, within participant, rf_echo (page 1 of 10). All 116 proteins whose Spearman correlation with rf_echo reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within analysis and outcome. Data: correlations sheet of 03_protein_association.xlsx.




BFR minus HLRT, within participant, rf_echo (page 2 of 10). All 116 proteins whose Spearman correlation with rf_echo reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within analysis and outcome. Data: correlations sheet of 03_protein_association.xlsx.




BFR minus HLRT, within participant, rf_echo (page 3 of 10). All 116 proteins whose Spearman correlation with rf_echo reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within analysis and outcome. Data: correlations sheet of 03_protein_association.xlsx.




BFR minus HLRT, within participant, rf_echo (page 4 of 10). All 116 proteins whose Spearman correlation with rf_echo reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within analysis and outcome. Data: correlations sheet of 03_protein_association.xlsx.




BFR minus HLRT, within participant, rf_echo (page 5 of 10). All 116 proteins whose Spearman correlation with rf_echo reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within analysis and outcome. Data: correlations sheet of 03_protein_association.xlsx.




BFR minus HLRT, within participant, rf_echo (page 6 of 10). All 116 proteins whose Spearman correlation with rf_echo reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within analysis and outcome. Data: correlations sheet of 03_protein_association.xlsx.




BFR minus HLRT, within participant, rf_echo (page 7 of 10). All 116 proteins whose Spearman correlation with rf_echo reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within analysis and outcome. Data: correlations sheet of 03_protein_association.xlsx.



BFR minus HLRT, within participant, rf_echo (page 8 of 10). All 116 proteins whose Spearman correlation with rf_echo reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within analysis and outcome. Data: correlations sheet of 03_protein_association.xlsx.



BFR minus HLRT, within participant, rf_echo (page 9 of 10). All 116 proteins whose Spearman correlation with rf_echo reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within analysis and outcome. Data: correlations sheet of 03_protein_association.xlsx.



BFR minus HLRT, within participant, rf_echo (page 10 of 10). All 116 proteins whose Spearman correlation with rf_echo reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within analysis and outcome. Data: correlations sheet of 03_protein_association.xlsx.