


A

All collections


1990 sets | 418 significant | rho 0.83 | 8 discordant



B

Collapse survivors


126 sets | 126 significant | rho 0.86 | 3 discordant



C

Discordant

8 sets | 8 significant | 8 discordant




genes ● 50 ● 150 ● 300 significant in ● Both ● BFR_Post-Pre ● HLRT_Post-Pre

S10 Figure. Set-level agreement between the two training responses. fgsea NES for each of 1990 sets in BFR_Post-Pre (x) against HLRT_Post-Pre (y), BH within contrast. (A) Every set. (B) Sets that survived collapsePathways in either contrast. (C) The 8 discordant sets, on opposite sides of zero and significant in one arm only, rescaled so each can be named. Dashed line is identity; grey points reach neither threshold; point size is gene count. Data: nes_scatter sheet of 03_enrich_scatter_fgsea.xlsx.

A

Hallmark and GO Slim


100 sets | 25 significant | rho 0.81 | 0 discordant



B

Up in both


14 sets | 14 significant | 0 discordant



C

Down in both

11 sets | 11 significant | 0 discordant



genes ● 50 ● 150 ● 300 significant in ● Both ● BFR_Post-Pre ● HLRT_Post-Pre

S11 Figure. Set-level agreement within Hallmark and GO Slim. (A) All 100 Hallmark and GO Slim sets, whose members do not nest. (B, C) The significant ones rescaled by direction so each can be named. Axes, colours and point size as in S10 Figure. Data: nes_scatter sheet of 03_enrich_scatter_fgsea.xlsx.