







S13 Figure (page 1 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 2 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 3 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 4 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 5 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 6 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 7 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 8 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 9 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 10 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 11 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 12 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 13 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 14 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 15 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 16 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 17 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 18 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 19 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 20 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 21 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 22 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 23 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 24 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 25 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 26 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 27 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 28 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 29 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 30 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 31 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 32 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S13 Figure (page 33 of 33). ROC curves for set scores, Pre to post, BFR. ROC curves for all 391 sets whose singscore separates the groups at nominal p, across 32 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 1 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 2 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 3 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose sing score separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 4 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 5 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 6 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 7 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 8 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose sing score separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 9 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 10 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 11 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 12 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 13 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 14 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 15 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 16 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 17 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singoscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 18 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 19 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 20 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 21 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 22 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 23 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 24 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose sing score separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 25 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 26 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 27 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 28 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 29 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 30 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 31 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singoscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 32 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S14 Figure (page 33 of 33). ROC curves for set scores, Pre to post, HLRT. ROC curves for all 389 sets whose singscore separates the groups at nominal p, across 33 paired legs, by collection then p. AUC has its direction fixed: red separates higher in post, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S15 Figure (page 1 of 8). ROC curves for set scores, Post BFR vs HLRT. ROC curves for all 94 sets whose singscore separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S15 Figure (page 2 of 8). ROC curves for set scores, Post BFR vs HLRT. ROC curves for all 94 sets whose singscore separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S15 Figure (page 3 of 8). ROC curves for set scores, Post BFR vs HLRT. ROC curves for all 94 sets whose singscore separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S15 Figure (page 4 of 8). ROC curves for set scores, Post BFR vs HLRT. ROC curves for all 94 sets whose sing score separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S15 Figure (page 5 of 8). ROC curves for set scores, Post BFR vs HLRT. ROC curves for all 94 sets whose singscore separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S15 Figure (page 6 of 8). ROC curves for set scores, Post BFR vs HLRT. ROC curves for all 94 sets whose singscore separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S15 Figure (page 7 of 8). ROC curves for set scores, Post BFR vs HLRT. ROC curves for all 94 sets whose singscore separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S15 Figure (page 8 of 8). ROC curves for set scores, Post BFR vs HLRT. ROC curves for all 94 sets whose singscore separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 1 of 4). ROC curves for set scores, Change, BFR vs HLRT. ROC curves for all 43 sets whose singscore separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 2 of 4). ROC curves for set scores, Change, BFR vs HLRT. ROC curves for all 43 sets whose sing score separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 3 of 4). ROC curves for set scores, Change, BFR vs HLRT. ROC curves for all 43 sets whose singscore separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S16 Figure (page 4 of 4). ROC curves for set scores, Change, BFR vs HLRT. ROC curves for all 43 sets whose singscore separates the groups at nominal p, across 32 paired participants, by collection then p. AUC has its direction fixed: red separates higher in BFR, blue lower. Shading is area under the curve, dashed line chance. p from the paired Wilcoxon signed-rank test, q from BH within collection and task. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_auc sheet of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 1 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 2 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 3 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 4 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p . Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 5 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S17 Figure (page 6 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 7 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p . Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




change in set score, T2 – T1, one point per leg


S17 Figure (page 8 of 23). Training response against phenotype, all legs. All 270 set–outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 9 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 10 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 11 of 23). Training response against phenotype, all legs. All 270 set–outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 12 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 13 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 14 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p . Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 15 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 16 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 17 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p . Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 18 of 23). Training response against phenotype, all legs. All 270 set–outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 19 of 23). Training response against phenotype, all legs. All 270 set–outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 20 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 21 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 22 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S17 Figure (page 23 of 23). Training response against phenotype, all legs. All 270 set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), by collection then p . Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 1 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 2 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 3 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 4 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 5 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S18 Figure (page 6 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 7 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S18 Figure (page 8 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.















change in phenotype



S18 Figure (page 9 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




change in phenotype




change in set score, T2 - T1, one point per leg


S18 Figure (page 10 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 11 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p . Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 12 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 13 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 14 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 15 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 16 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 17 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p . Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 18 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 19 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p . Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 20 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 21 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 22 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S18 Figure (page 23 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.




S18 Figure (page 24 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S18 Figure (page 25 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p . Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S18 Figure (page 26 of 26). BFR minus HLRT against phenotype, within participant. All 303 set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), by collection then p. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430. Data: set_association and set_by_arm sheets of 05_classify_and_associate_sets.xlsx.



S19 Figure. Nominal hits relative to chance, by collection. Paired Wilcoxon per set across 5 collections, uncorrected p. Bar length is observed nominal hits divided by the count that collection returns under the null; red clears 1, grey does not. Labels give observed of tested. Data: chance_expectation sheet of 05_classify_and_associate_sets.xlsx.