


Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT

change in phenotype




All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT




All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT

change in phenotype




All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

—●— BFR —●— HLRT



All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT




All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT




All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT



All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT




All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT




All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

—●— BFR —●— HLRT



All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT




All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs


Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT


GOBP: Protein
Neddylation vs rf_echo
r = -0.25 p 0.044 q 1




GOBP: Positive
Reg. Pattern
Recognition
Receptor
Signaling vs vl_echo
r = -0.25 p 0.044 q 0.99




GOBP: Reg.
Inflammatory
Response vs vl_echo
r = +0.25 p 0.044 q 0.99




GOBP: Tissue
Homeostasis vs rf_csa
r = -0.25 p 0.045 q 0.89




GOBP: Production
Of Molecular
Mediator
Involved In
Inflammatory
Response vs rf_csa
r = +0.25 p 0.045 q 0.89




GOBP: Lipoprotein
Metabolism vs rf_csa
r = -0.25 p 0.045 q 0.89




GOBP: Intrinsic
Apoptotic
Signaling
In Resp. to
Endoplasmic
Reticulum
Stress vs rf_echo
r = -0.25 p 0.045 q 1




GOBP: Protein
Polyubiquitination vs rf_csa
r = -0.25 p 0.045 q 0.89




GOBP: Positive Reg.
DNA Repair vs rf_echo
r = +0.25 p 0.045 q 1




GOBP: Central Nervous
System Neuron
Differentiation vs vl_echo
r = +0.25 p 0.046 q 0.99




GOBP: Reg. Protein
Depolymerization vs vl_csa
r = -0.25 p 0.046 q 0.96




GOBP: Skeletal System
Morphogenesis vs vl_csa
r = -0.25 p 0.046 q 0.96




GOBP: RNA Mod. vs vl_echo
r = -0.25 p 0.047 q 0.99




GOBP: tRNA
Metabolism vs rf_csa
r = +0.25 p 0.047 q 0.89



GOBP: Mesoderm
Development vs vl_echo
r = -0.25 p 0.047 q 0.99



GOBP: Myoblast
Differentiation vs rf_csa
r = -0.25 p 0.047 q 0.89




change in set score, T2 - T1, one point per leg

All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT




All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

● BFR ● HLRT




All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

● BFR ● HLRT



All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

—●— BFR —●— HLRT



All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

Training response against phenotype, all legs

Spearman, pooled analysis, 65 legs, both arms pooled

BFR HLRT



All 270 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the pooled correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.