






S17 Figure (page 1 of 4). Training response against phenotype, all legs, Reactome. All 38 Reactome set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 21.5 of the 430 Reactome sets. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.



S17 Figure (page 2 of 4). Training response against phenotype, all legs, Reactome. All 38 Reactome set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 21.5 of the 430 Reactome sets. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.




S17 Figure (page 3 of 4). Training response against phenotype, all legs, Reactome. All 38 Reactome set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 21.5 of the 430 Reactome sets. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.



Reactome: Apoptotic Cleavage Of Cellular Proteins

vs rf_csa $r = -0.25$ $p = 0.049$ $q = 1$


Method	n	r	p
BFR	32	-0.22	0.226
HLRT	33	-0.26	0.151




Reactome: Organelle Biogenesis And Maintenance

vs vl_csa $r = +0.25$ $p = 0.049$ $q = 1$


Method	n	r	p
BFR	32	+0.01	0.946
HLRT	33	+0.52	0.002



S17 Figure (page 4 of 4). Training response against phenotype, all legs, Reactome. All 38 Reactome set-outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 21.5 of the 430 Reactome sets. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.




S18 Figure (page 1 of 4). BFR minus HLRT, within participant, Reactome. All 44 Reactome set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p, uncorrected: chance alone returns 21.5 of the 430 Reactome sets. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.



S18 Figure (page 2 of 4). BFR minus HLRT, within participant, Reactome. All 44 Reactome set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 21.5 of the 430 Reactome sets. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.



S18 Figure (page 3 of 4). BFR minus HLRT, within participant, Reactome. All 44 Reactome set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 21.5 of the 430 Reactome sets. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.



S18 Figure (page 4 of 4). BFR minus HLRT, within participant, Reactome. All 44 Reactome set-outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within collection and outcome. Nominal p , uncorrected: chance alone returns 21.5 of the 430 Reactome sets. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.