


BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT




All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT




All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT




All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT




All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

—●— BFR —●— HLRT




All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT




All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT

change in phenotype




All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants


BFR HLRT



All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants



All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants



change in phenotype




All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT




All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT



All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT




All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT



All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT



All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT



All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants




All 303 set–outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants

BFR HLRT



All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.

BFR minus HLRT, within participant

Spearman, differential analysis, 32 paired participants



All 303 set-outcome pairs reaching nominal p, by collection then p. Lines are fitted within each arm; the header r is the differential correlation, q from BH within collection and outcome. Nominal p, uncorrected. Each collection is read against its own expectation: GOBP 68.3 of 1366, GO_Slim 3.0 of 59, Hallmark 2.1 of 41, KEGG_Legacy 4.7 of 94, Reactome 21.5 of 430.