



BFR
HLRT
KEGG_Legacy: Aminoacyl tRNA Biosynthesis
vs vl_csa r = +0.37 p 0.0023 q 0.22
BFR n=32 r=+0.31 p=0.084
HLRT n=33 r=+0.54 p=0.001
KEGG_Legacy: WNT Signaling Pathway
vs rf_echo r = −0.36 p 0.0037 q 0.23
BFR n=32 r=−0.48 p=0.006
HLRT n=33 r=−0.26 p=0.138
KEGG_Legacy: Colorectal Cancer
vs rf_echo r = −0.34 p 0.005 q 0.23
BFR n=32 r=−0.46 p=0.008
HLRT n=33 r=−0.25 p=0.166
KEGG_Legacy: Lysine Degradation
vs vl_csa r = +0.34 p 0.0058 q 0.27
BFR n=32 r=+0.25 p=0.169
HLRT n=33 r=+0.45 p=0.010
KEGG_Legacy: Adherens Junction
vs vl_csa r = −0.31 p 0.013 q 0.41
BFR n=32 r=−0.28 p=0.125
HLRT n=33 r=−0.40 p=0.023
KEGG_Legacy: Hypertrophic Cardiomyopathy Hcm
vs rf_echo r = +0.29 p 0.019 q 0.52
BFR n=32 r=+0.40 p=0.023
HLRT n=33 r=+0.15 p=0.413
KEGG_Legacy: Cysteine And Methionine Metabolism
vs vl_csa r = +0.29 p 0.02 q 0.48
BFR n=32 r=+0.25 p=0.174
HLRT n=33 r=+0.36 p=0.043
KEGG_Legacy: Chemokine Signaling Pathway
vs rf_echo r = −0.28 p 0.022 q 0.52
BFR n=32 r=−0.28 p=0.126
HLRT n=33 r=−0.33 p=0.062
KEGG_Legacy: Renal Cell Carcinoma
vs vl_echo r = +0.28 p 0.024 q 0.68
BFR n=32 r=+0.10 p=0.571
HLRT n=33 r=+0.44 p=0.011
KEGG_Legacy: Erbb Signaling Pathway
vs rf_echo r = −0.27 p 0.028 q 0.53
BFR n=32 r=−0.41 p=0.021
HLRT n=33 r=−0.17 p=0.334
KEGG_Legacy: Oxidative Phosphorylation
vs rf_csa r = +0.27 p 0.032 q 0.89
BFR n=32 r=+0.16 p=0.373
HLRT n=33 r=+0.33 p=0.062
KEGG_Legacy: Purine Metabolism
vs rf_csa r = −0.27 p 0.033 q 0.89
BFR n=32 r=−0.39 p=0.030
HLRT n=33 r=−0.15 p=0.411
change in phenotype
change in set score, T2 − T1, one point per leg

S11. Pooled association, all legs, KEGG_Legacy (page 2 of 2)



S12. Differential association, BFR minus HLRT, KEGG_Legacy (page 1 of 2)



S12. Differential association, BFR minus HLRT, KEGG_Legacy (page 2 of 2)

