




S11. Pooled association, all legs, Reactome (page 1 of 3)





BFR
HLRT
Reactome: Rhoq GTPase Cycle
vs rf_echo r = +0.30 p 0.016 q 0.93
BFR n=32 r=+0.12 p=0.513
HLRT n=33 r=+0.49 p=0.004
Reactome: Mitochondrial Translation Elongation
vs rf_csa r = +0.29 p 0.018 q 0.99
BFR n=32 r=+0.32 p=0.075
HLRT n=33 r=+0.22 p=0.208
Reactome: Cytoprotection By HMOX1
vs rf_csa r = +0.28 p 0.023 q 0.99
BFR n=32 r=+0.09 p=0.625
HLRT n=33 r=+0.39 p=0.024
Reactome: Sars Cov 1 Host Interactions
vs rf_echo r = +0.28 p 0.025 q 0.93
BFR n=32 r=+0.39 p=0.029
HLRT n=33 r=+0.22 p=0.224
Reactome: Sars Cov Infections
vs rf_echo r = +0.28 p 0.026 q 0.93
BFR n=32 r=+0.36 p=0.044
HLRT n=33 r=+0.21 p=0.242
Reactome: Peroxisomal Protein Import
vs vl_echo r = +0.27 p 0.028 q 0.98
BFR n=32 r=+0.26 p=0.147
HLRT n=33 r=+0.25 p=0.160
Reactome: Late Endosomal Microautophagy
vs rf_csa r = −0.27 p 0.03 q 0.99
BFR n=32 r=−0.26 p=0.155
HLRT n=33 r=−0.24 p=0.178
Reactome: Deubiquitination
vs rf_echo r = −0.27 p 0.03 q 0.93
BFR n=32 r=−0.61 p=0.000
HLRT n=33 r=+0.05 p=0.773
Reactome: Response To Elevated Platelet Cytosolic Ca2
vs vl_echo r = +0.27 p 0.03 q 0.98
BFR n=32 r=+0.28 p=0.123
HLRT n=33 r=+0.28 p=0.116
Reactome: Sars Cov 1 Modulates Host Translation Machinery
vs rf_echo r = +0.27 p 0.03 q 0.93
BFR n=32 r=+0.37 p=0.038
HLRT n=33 r=+0.27 p=0.131
Reactome: Intra Golgi And Retrograde Golgi To Er Traffic
vs vl_csa r = +0.27 p 0.03 q 1
BFR n=32 r=−0.06 p=0.761
HLRT n=33 r=+0.57 p=0.001
Reactome: Cell Surface Interactions At The Vascular Wall
vs vl_csa r = −0.27 p 0.033 q 1
BFR n=32 r=−0.36 p=0.043
HLRT n=33 r=−0.22 p=0.211
change in phenotype
change in set score, T2 − T1, one point per leg



BFR
HLRT
Reactome: Copii Mediated Vesicle Transport
vs vl_csa r = +0.26 p 0.034 q 1
BFR n=32 r=+0.08 p=0.648
HLRT n=33 r=+0.42 p=0.015
Reactome: Signaling By Retinoic Acid
vs vl_echo r = +0.26 p 0.034 q 0.98
BFR n=32 r=+0.14 p=0.431
HLRT n=33 r=+0.39 p=0.024
Reactome: Mitotic Prophase
vs vl_csa r = +0.26 p 0.035 q 1
BFR n=32 r=+0.46 p=0.009
HLRT n=33 r=+0.09 p=0.619
Reactome: Esr Mediated Signaling
vs rf_echo r = −0.26 p 0.036 q 0.93
BFR n=32 r=−0.34 p=0.055
HLRT n=33 r=−0.21 p=0.242
Reactome: Antimicrobial Mechanism Of IFN Stimulated Genes
vs vl_echo r = +0.26 p 0.036 q 0.98
BFR n=32 r=+0.17 p=0.352
HLRT n=33 r=+0.44 p=0.012
Reactome: Cdc42 GTPase Cycle
vs rf_echo r = +0.25 p 0.042 q 0.93
BFR n=32 r=+0.04 p=0.817
HLRT n=33 r=+0.47 p=0.007
Reactome: Metabolism Of Steroids
vs vl_echo r = −0.25 p 0.043 q 0.98
BFR n=32 r=−0.25 p=0.172
HLRT n=33 r=−0.34 p=0.055
Reactome: G2 M DNA Damage Checkpoint
vs rf_csa r = −0.25 p 0.044 q 0.99
BFR n=32 r=−0.32 p=0.072
HLRT n=33 r=−0.16 p=0.382
Reactome: Sars Cov 2 Modulates Host Translation Machinery
vs rf_echo r = +0.25 p 0.044 q 0.93
BFR n=32 r=+0.34 p=0.057
HLRT n=33 r=+0.20 p=0.275
Reactome: Copii Mediated Vesicle Transport
vs rf_csa r = +0.25 p 0.045 q 0.99
BFR n=32 r=+0.21 p=0.259
HLRT n=33 r=+0.24 p=0.180
Reactome: Apoptotic Cleavage Of Cellular Proteins
vs rf_csa r = −0.25 p 0.049 q 0.99
BFR n=32 r=−0.22 p=0.226
HLRT n=33 r=−0.26 p=0.148
Reactome: Foxo Mediated Transcription
vs rf_echo r = −0.24 p 0.05 q 0.93
BFR n=32 r=−0.32 p=0.074
HLRT n=33 r=−0.08 p=0.643
change in phenotype
change in set score, T2 − T1, one point per leg


BFR − HLRT
Reactome: Copii Mediated Vesicle Transport
vs vl_echo r = −0.52 p 0.0028 q 0.97
Reactome: Cargo Concentration In The Er
vs vl_echo r = −0.48 p 0.0055 q 0.97
Reactome: L1cam Interactions
vs vl_csa r = −0.48 p 0.0064 q 0.96
Reactome: G Alpha Q Signalling Events
vs rf_echo r = −0.45 p 0.011 q 0.91
Reactome: Mitotic Prophase
vs vl_csa r = +0.44 p 0.013 q 0.96
Reactome: Response To Elevated Platelet Cytosolic Ca2
vs vl_csa r = −0.43 p 0.014 q 0.96
Reactome: Slc Mediated Transmembrane Transport
vs rf_csa r = −0.43 p 0.014 q 0.84
Reactome: Plasma Lipoprotein Assembly Remodeling And Clearance
vs rf_csa r = −0.43 p 0.015 q 0.84
Reactome: Signaling By Insulin Receptor
vs rf_echo r = −0.43 p 0.015 q 0.91
Reactome: Cargo Trafficking To The Periciliary Membrane
vs vl_echo r = −0.42 p 0.016 q 0.97
Reactome: Response To Elevated Platelet Cytosolic Ca2
vs vl_echo r = +0.42 p 0.017 q 0.97
Reactome: Chromosome Maintenance
vs vl_csa r = +0.42 p 0.018 q 0.96
change in phenotype, BFR minus HLRT
change in set score, T2 − T1, BFR minus HLRT, one point per participant


BFR − HLRT
Reactome: Bacterial Infection Pathways
vs rf_csa r = +0.41 p 0.02 q 0.84
Reactome: Apoptosis
vs vl_csa r = −0.41 p 0.02 q 0.96
Reactome: Rhov GTPase Cycle
vs vl_csa r = −0.41 p 0.021 q 0.96
Reactome: Regulation Of Insulin Secretion
vs rf_csa r = −0.41 p 0.021 q 0.84
Reactome: Cytoprotection By HMOX1
vs vl_echo r = −0.41 p 0.021 q 0.97
Reactome: Hcmv Late Events
vs vl_csa r = +0.41 p 0.022 q 0.96
Reactome: Mitochondrial tRNA Aminoacylation
vs vl_csa r = +0.41 p 0.022 q 0.96
Reactome: Signaling By Fgfr
vs vl_echo r = +0.41 p 0.022 q 0.97
Reactome: Organelle Biogenesis And Maintenance
vs vl_echo r = −0.41 p 0.022 q 0.97
Reactome: Extra Nuclear Estrogen Signaling
vs rf_echo r = −0.40 p 0.024 q 0.91
Reactome: Esr Mediated Signaling
vs rf_echo r = −0.40 p 0.025 q 0.91
Reactome: Mitf M Regulated Melanocyte Development
vs rf_echo r = −0.39 p 0.027 q 0.91
change in phenotype, BFR minus HLRT
change in set score, T2 − T1, BFR minus HLRT, one point per participant

S12. Differential association, BFR minus HLRT, Reactome (page 3 of 4)




BFR − HLRT
Reactome: Hcmv Early Events
vs rf_csa r = +0.36 p 0.041 q 0.84
Reactome: Signaling By Fgfr2
vs vl_echo r = +0.36 p 0.042 q 0.97
Reactome: Peroxisomal Protein Import
vs vl_csa r = +0.36 p 0.042 q 0.96
Reactome: Intra Golgi And Retrograde Golgi To Er Traffic
vs vl_echo r = −0.36 p 0.043 q 0.97
Reactome: Aurka Activation By Tpx2
vs rf_echo r = +0.36 p 0.045 q 0.91
Reactome: Recruitment Of Mitotic Centrosome Proteins And Complexes
vs rf_echo r = +0.36 p 0.045 q 0.91
Reactome: Autophagy
vs rf_echo r = +0.35 p 0.049 q 0.91
change in phenotype, BFR minus HLRT
change in set score, T2 − T1, BFR minus HLRT, one point per participant