
Complex subunits, summary by leg
BFR
HLRT
fry FDR 0.93
fry FDR 0.93
fry FDR 0.93
fry FDR 0.93
fry FDR 0.93
log2 change, post − pre
0.4
0.2
0.0
−0.2
CI
(n = 41/44)
CII
(n = 4/4)
CIII
(n = 10/10)
CIV
(n = 16/21)
CV
(n = 17/21)


Assembly factors and other OXPHOS proteins, by leg
BFR
HLRT
log2 change, post − pre
0.50
0.25
0.00
−0.25
CI
(n = 14/22)
CII
(n = 2/4)
CIII
(n = 5/6)
CIV
(n = 18/30)
CV
(n = 4/5)
Other
(n = 2/4)


Complex subunits by leg
BFR
HLRT
CI (41), fry interaction FDR 0.93
CII (4), fry interaction FDR 0.93
CIII (10), fry interaction FDR 0.93
CIV (16), fry interaction FDR 0.93
CV (17), fry interaction FDR 0.93
log2 change, post − pre
0.4
0.2
0.0
−0.2
NDUFA2 NDUFA11 MT−ND2 MT−ND5 NDUFC1 MT−ND3 NDUFS3 MT−ND1 NDUFS4 NDUFA9 MT−ND4 NDUFS5 NDUFV1 NDUFA3 NDUFB8 NDUFB11 NDUFS1 NDUFA1 NDUFB10 NDUFS7 NDUFB3 NDUFA5 NDUFA10 NDUFB1 NDUFAB1 NDUFA8 NDUFS6 NDUFA12 NDUFV2 NDUFA13 NDUFB7 NDUFB4 NDUFB9 NDUFS8 NDUFB6 NDUFA6 NDUFC2 NDUFS2 NDUFB5 NDUFA7 NDUFV3
SDHD SDHC SDHA SDHB
UQCR10 MT−CYB UQCR11 UQCRFS1 UQCRC1 UQCRC2 UQCRH UQCRB UQCRQ CYC1
COX7C MT−CO1 COX6A2 COX7A1 MT−CO3 COX7A2 NDUFA4 COX7B COX5B COX7A2L COX6B1 COX4I1 MT−CO2 COX5A COX6C COX8A
ATP5MF DMAC2L ATP5F1C ATP5MPL MT−ATP6 ATP5F1A ATP5ME ATP5F1B ATP5MG ATP5PB MT−ATP8 ATP5PD ATP5IF1 ATP5PF ATP5PO ATP5F1D ATP5MD


Complex assembly factors by leg
BFR
HLRT
CI (14)
0.50
0.25
0.00
−0.25
TMEM70
NDUFAF4
DMAC1
TIMMDC1
NDUFAF3
NUBPL
NDUFAF1
AIFM1
NDUFAF6
ACAD9
NDUFAF2
NDUFAF7
NDUFAF5
TMEM126A
CII (2)
SDHAF4
SDHAF2
CIII (5)
log2 change, post − pre
UQCC3
UQCC1
UQCC2
LYRM7
BCS1L
CIV (18)
CMC1
COX19
SCO1
COX14
SCO2
TACO1
COA7
COX11
SURF1
COA4
COA6
CMC2
COX20
SMIM20
COX17
COA3
TIMM21
COA5
CV (4)
TMEM70
ATPAF2
ATPAF1
FMC1