

S1 Figure. Ten strongest collapse survivors, all collections. (A) BFR_Post−Pre. (B) HLRT_Post−Pre. (C) Modality_x_Time_Interaction. (D) BFR_Post−HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is the collection. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.



S2 Figure. Ten strongest collapse survivors, Hallmark. (A) BFR_Post−Pre. (B) HLRT_Post−Pre. (C) Modality_x_Time_Interaction. (D) BFR_Post−HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is −log10 FDR. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.

S3 Figure. Ten strongest collapse survivors, Reactome. (A) BFR_Post−Pre. (B) HLRT_Post−Pre. (C) BFR_Post−HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is −log10 FDR. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.



S4 Figure. Ten strongest collapse survivors, KEGG_Legacy. (A) BFR_Post−Pre. (B) HLRT_Post−Pre. (C) BFR_Post−HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is −log10 FDR. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.



S5 Figure. Ten strongest collapse survivors, GOBP. (A) BFR_Post−Pre. (B) HLRT_Post−Pre. (C) Modality_x_Time_Interaction. (D) BFR_Post−HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is −log10 FDR. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.



S6 Figure. Ten strongest collapse survivors, GO_Slim. (A) BFR_Post−Pre. (B) HLRT_Post−Pre. (C) Modality_x_Time_Interaction. (D) BFR_Post−HLRT_Post. Up to ten collapsePathways survivors per contrast, lowest adjusted p first; FDR is BH within contrast. Point size is the number of measured genes in the set; colour is −log10 FDR. Contrasts with no survivor are omitted. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.



S7 Figure. Significant sets before and after collapse. Sets at FDR 0.05 per collection and contrast, before and after collapsePathways, which makes a set the child of another significant set when it shows no enrichment of its own ($p > 0.05$) either outside or inside that set's genes. Data: set_tests sheet of 01_run_fgsea_and_fry.xlsx.