





S17 Figure (page 1 of 2). Training response against phenotype, all legs, KEGG_Legacy. All 17 KEGG_Legacy set−outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Nominal p, uncorrected: chance alone returns 18.8 of the 4 x 94 KEGG_Legacy set−outcome pairs. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.



S17 Figure (page 2 of 2). Training response against phenotype, all legs, KEGG_Legacy. All 17 KEGG_Legacy set−outcome pairs whose Spearman correlation reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first. Nominal p, uncorrected: chance alone returns 18.8 of the 4 x 94 KEGG_Legacy set−outcome pairs. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within collection and outcome. Data: set_association and set_by_arm sheets of 06_set_association.xlsx.



S18 Figure (page 1 of 2). BFR minus HLRT, within participant, KEGG_Legacy. All 21 KEGG_Legacy set−outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Nominal p, uncorrected: chance alone returns 18.8 of the 4 x 94 KEGG_Legacy set−outcome pairs. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within collection and outcome. Data: set_association sheet of 06_set_association.xlsx.



S18 Figure (page 2 of 2). BFR minus HLRT, within participant, KEGG_Legacy. All 21 KEGG_Legacy set−outcome pairs whose Spearman correlation reaches nominal p in the differential analysis (32 paired participants), smallest p first. Nominal p, uncorrected: chance alone returns 18.8 of the 4 x 94 KEGG_Legacy set−outcome pairs. One point per participant: the change in their BFR leg minus the change in their HLRT leg, on both axes. The line is a least−squares fit; the header r is the differential correlation, the one tested. q from BH within collection and outcome. Data: set_association sheet of 06_set_association.xlsx.