





S8 Figure. Protein volcanoes with collapse-surviving fgsea sets ringed. (A) Modality_x_Time_Interaction. (B) BFR_Post-Pre. (C) HLRT_Post-Pre. (D) BFR_Post-HLRT_Post. Points are proteins, coloured when BH FDR < 0.05 within the contrast, with the five lowest-FDR proteins named. The ring shows up to twelve fgsea sets that survived collapsePathways, lowest adjusted p first, in either direction, coloured by NES; ticks mark their leading-edge proteins. Protein colour and counts read protein-level BH FDR, the ring set-level fgsea FDR: two separate claims. Data: protein_results and set_tests sheets of 01_run_fgsea_and_fry.xlsx.

A

BFR_Post-Pre


Labels ranked by pi-score; colours and counts use BH FDR



B

HLRT_Post-Pre

Labels ranked by pi-score; colours and counts use BH FDR



S9 Figure. Protein volcanoes labelled by pi-score. (A) BFR_Post-Pre. (B) HLRT_Post-Pre. As S8 Figure, but the five proteins named are those with the smallest pi-score, $P.Value^{|logFC|}$ (Xiao et al. 2014). Pi controls no error rate, so no protein named here is a discovery. The ring shows up to twelve fgsea sets that survived collapsePathways, lowest adjusted p first, in either direction, coloured by NES; ticks mark their leading-edge proteins. Protein colour and counts read protein-level BH FDR, the ring set-level fgsea FDR: two separate claims. Data: protein_results and set_tests sheets of 01_run_fgsea_and_fry.xlsx.