













Training response against phenotype, all legs, vl_csa (page 3 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, vl_csa (page 4 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, vl_csa (page 5 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, vl_csa (page 6 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, vl_csa (page 7 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, vl_csa (page 8 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, vl_csa (page 9 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.







Training response against phenotype, all legs, vl_csa (page 11 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




Training response against phenotype, all legs, vl_csa (page 12 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.



Training response against phenotype, all legs, vl_csa (page 13 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.



Training response against phenotype, all legs, vl_csa (page 14 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.





NAPG Q99747

vs vl_csa $r = +0.24$ $p = 0.05$ $q = 0.82$

	BFR	n	r	p
BFR	n=32	$r = +0.15$	$p = 0.424$	
HLRT	n=33	$r = +0.37$	$p = 0.037$	

TARS3 A2RTX5

vs vl_csa $r = +0.24$ $p = 0.05$ $q = 0.82$


	BFR	n	r	p
BFR	n=32	$r = +0.12$	$p = 0.511$	
HLRT	n=33	$r = +0.42$	$p = 0.017$	

DCXR Q7Z4W1

vs vl_csa $r = +0.24$ $p = 0.05$ $q = 0.82$


	BFR	n	r	p
BFR	n=32	$r = +0.29$	$p = 0.108$	
HLRT	n=33	$r = +0.24$	$p = 0.184$	

change in phenotype




change in protein, log2, T2 - T1, one point per leg


Training response against phenotype, all legs, vl_csa (page 16 of 16). All 183 proteins whose Spearman correlation with vl_csa reaches nominal p in the pooled analysis (65 legs, both arms pooled), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the pooled correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




BFR minus HLRT, within participant, vl_csa (page 1 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




BFR minus HLRT, within participant, vl_csa (page 2 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




BFR minus HLRT, within participant, vl_csa (page 3 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




BFR minus HLRT, within participant, vl_csa (page 4 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




BFR minus HLRT, within participant, vl_csa (page 5 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




BFR minus HLRT, within participant, vl_csa (page 6 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




BFR minus HLRT, within participant, vl_csa (page 7 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




BFR minus HLRT, within participant, vl_csa (page 8 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




BFR minus HLRT, within participant, vl_csa (page 9 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.




BFR minus HLRT, within participant, vl_csa (page 10 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.



BFR minus HLRT, within participant, vl_csa (page 11 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.



BFR minus HLRT, within participant, vl_csa (page 12 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.



BFR minus HLRT, within participant, vl_csa (page 13 of 13). All 146 proteins whose Spearman correlation with vl_csa reaches nominal p in the differential analysis (32 paired participants), smallest p first; chance alone returns 152 of 3,042 proteins. Lines are fitted within each arm; the header r is the differential correlation and the inset gives it within each arm. q from BH within analysis and outcome. Data: correlations and protein_by_arm sheets of 03_protein_association.xlsx.