



All 15 modules by 44 biopsies, grouped by arm and time point; column labels are subjects. Fill: module eigengene. Colours stop at the 99th percentile of |value|, 0.46. Rows follow module size. Grey in the trait tracks: not measured (T3 has no traits).


Contrast overview: modules



All 15 modules by the nine contrasts. Fill: eigengene fold change from limma; status: p and BH q within contrast. Colours stop at the 99th percentile of |value|, 0.4. Rows follow module size. Table: c_data/module_contrasts.xlsx, eigengene_tests.

Module eigengenes across the nine contrasts


limma on eigengenes, protein design, subject block (correlation 0.135); BH within contrast



Fill: moderated t of each eigengene in each contrast. Label: nominal p where below 0.05. Black border: BH < 0.05. Table: c_data/module_contrasts.xlsx.

Modules as protein sets across the nine contrasts


limma::fry on limpa's matrix, protein design, subject block (correlation 0.204)



Fill: −log10 fry p, positive when the module moved up. fry's FDR runs over the 15 modules within each contrast. Label: nominal p where below 0.05.
Black border: BH < 0.05. Table: c_data/module_contrasts.xlsx.

Module membership against protein significance

Spearman between each member's kME and its protein−level moderated t, per module and contrast



Fill: rho; positive when the module's most central proteins moved up most. Members share a module, so the p describes rather than tests. Label: nominal p where below 0.05. Black border: BH < 0.05. Table: c_data/module_contrasts.xlsx.

Membership against the primary contrast

one point per module member; Spearman rho in each header



Line: least−squares fit, for reference. Table: c_data/module_contrasts.xlsx, membership_contrast.