

a Does a feature follow a trait differently in HR and LR?
HR slope minus LR slope (limma, feature ~ group x trait). Number: % of features at p < 0.05 (white: 5, chance);
bold below: count at FDR < 0.05. The composite score defined the groups and is left out
Proteins
Set scores
Modules
fCSA mixed 3 3 3 3 6 2 0 1 3 2 2 3 0 0 0 0 23 0
fCSA type1 5 2 2 2 7 2 2 1 3 1 2 1 0 0 0 0 15 0
fCSA type2 4 4 4 4 7 3 0 3 5 7 2 6 0 0 8 0 15 0
mCSA 6 5 7 3 2 4 2 3 4 6 1 1 4 1 0 0 0 0 0 8
leg press 1RM 4 10 6 3 4 1 2 8 11 44 1 9 1 1 0 31 8 1 0 8 0
leg ext 1RM 5 4 2 2 1 2 1 2 2 1 2 2 0 8 0 0 0 0
volume load (kg) 3 6 1 8 1 3 10 0 0 23
T1 level
T1, gain
T2 - T1
T2 level
T3 - T2
T3 level
% at p < 0.05
20%
15%
10%
5%
0%
b The strongest screen, within each group
leg press 1RM (% change), T2 - T1, 1,422 set scores; 44 of 44 FDR hits change sign
Cardiac muscle contraction
Regulation of the force of ...
Actin myosin filaments sliding
Actin mediated cell contrac...
Muscle organ morphogenesis
Cardiac muscle tissue morph...
Spearman rho in LR
Spearman rho in HR
HR - LR slope
FDR < 0.05
p < 0.05 only
not significant
c Cardiac muscle tissue morphogenesis, person by person
HR slope minus LR slope: t = -9.4, p < 0.001, FDR < 0.001
leg press 1RM (% change)
set score, T2 - T1
HR: rho -0.95 (n = 8)
LR: rho 0.95 (n = 8)