

a Training signature rank
fry: each arm's Π < 0.05 training sets in the other arm's ranking
LR training sets in HRs ranking
up (n = 13), 69% same way, FDR 0.069
down (n = 7), 86% same way, FDR 0.091
HR training sets in LRs ranking
up (n = 11), 82% same way, FDR 0.25
down (n = 14), 79% same way, FDR 0.25
0
500
1000
1500
rank in the other arm's training contrast, up to down (n = 1900)
b Acute signature rank
fry: each arm's Π < 0.05 acute sets in the other arm's ranking
LR acute sets in HRs ranking
up (n = 21), 86% same way, FDR 0.0072
down (n = 12), 83% same way, FDR 0.0072
HR acute sets in LRs ranking
up (n = 44), 66% same way, FDR 0.42
down (n = 18), 89% same way, FDR 0.42
0
500
1000
1500
rank in the other arm's acute contrast, up to down (n = 1900)
c Where training-responsive proteins go after the bout
p < 0.05 sets; change from own T1 (log2) ± SE; % same way, fry FDR; † T2 shared
Measured in HR
Measured in LR
Changed by training in HR
Changed by training in LR
selection window
8% same way FDR 0.0051 †
11% same way FDR 0.0051 †
90% same way FDR 0.14
27% same way FDR 0.64
74% same way FDR 0.25
46% same way FDR 0.92
82% same way FDR 0.016
30% same way FDR 0.13
76% same way FDR 0.026
45% same way FDR 0.57
3% same way FDR 1.1e-05 †
10% same way FDR 0.0011 †
T1
T2
T3
decreased in training
increased in training