

Sample model overview: modules




All 12 modules, one row each. Moderated t of the within and between terms, abundance ~ timepoint + between + within, subject blocked. Colours stop at the 99th percentile of |t|, 2.4. Rows follow module size. Grey: not tested. Table: c_data/04_associate.xlsx, model.

Change score overview, training window: modules




All 12 modules, one row each. Spearman rho between the eigengene (change, T1 to T2) and each outcome. Rows follow module size. Grey: not tested. Table: c_data/04_associate.xlsx, change_score.

Change score overview, baseline window: modules



All 12 modules, one row each. Spearman rho between the eigengene (T1 level) and each outcome. Rows follow module size. Grey: not tested. Table: c_data/04_associate.xlsx, change_score.


Change score overview, acute window: modules



All 12 modules, one row each. Spearman rho between the eigengene (change, T2 to T3) and each outcome. Rows follow module size. Grey: not tested. Table: c_data/04_associate.xlsx, change_score.

fibres_mixed, between−person term


1 of 12 modules at p < 0.05, 0 at BH < 0.05. Page 1 of 1.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


4 of 12 modules at p < 0.05, 0 at BH < 0.05. Page 1 of 1.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


2 of 12 modules at p < 0.05, 0 at BH < 0.05. Page 1 of 1.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


1 of 12 modules at p < 0.05, 0 at BH < 0.05. Page 1 of 1.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


1 of 12 modules at p < 0.05, 0 at BH < 0.05. Page 1 of 1.



limma, abundance ~ timepoint + between + within, subject blocked, on the T1 and T2 biopsies with the trait measured. Lines join each subject's two biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, BH q within trait and term, and the biopsies and subjects where the feature was observed. Table: c_data/04_associate.xlsx, model.

Change score, training window


5 of 240 module−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 1 of 1.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


13 of 240 module−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 1 of 2.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


13 of 240 module−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 2 of 2.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, acute window

7 of 240 module−outcome pairs at p < 0.05, 0 at BH < 0.05. Page 1 of 1.



One point per subject. Lines are a least−squares fit per arm, drawn as a guide; the tests are Spearman. Header: rho and p pooled over both arms, BH q within window and outcome, and the subjects used. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.