
Module eigengenes across the nine contrasts


limma on eigengenes, protein design, subject block (correlation 0.095); BH within contrast



Fill: moderated t of each eigengene in each contrast. Label: nominal p where below 0.05. Black border: BH < 0.05. Table: c_data/02_contrasts.xlsx.

Modules as protein sets across the nine contrasts

limma::fry on the imputed matrix, protein design, subject block (correlation 0.176)



Fill: −log10 fry p, positive when the module moved up. fry's FDR runs over the 12 modules within each contrast. Label: nominal p where below 0.05.
Black border: BH < 0.05. Table: c_data/02_contrasts.xlsx.

Module membership against protein significance


Spearman between each member's kME and its protein−level moderated t, per module and contrast



Fill: rho; positive when the module's most central proteins moved up most. Members share a module, so the p describes rather than tests. Label: nominal p where below 0.05. Black border: BH < 0.05. Table: c_data/02_contrasts.xlsx.

Module membership against phenotype association

Spearman between each member's kME and its protein rho with the phenotype over training



Fill: rho; positive when central proteins track the phenotype most positively. Descriptive, as above. Label: nominal p where below 0.05. Black border: BH < 0.05. Table: c_data/02_contrasts.xlsx.

Membership against the primary contrast

one point per module member; Spearman rho in each header



Line: least−squares fit, for reference. Table: c_data/02_contrasts.xlsx, membership_contrast.