
fibres_mixed, between−person term


1 of 12 modules at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, between−person term


1 of 12 modules at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

leg_ext_1rm, within−person term


2 of 12 modules at p < 0.05, 0 at BH < 0.05.



limma on 29 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

mcsa, between−person term


4 of 12 modules at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

type1_share, within−person term


1 of 12 modules at p < 0.05, 0 at BH < 0.05.



limma on 30 biopsies, abundance ~ timepoint + between + within, subject blocked. Lines join each subject's T1 and T2 biopsies: the within term reads their slopes, the between term the spread of subjects. Header: effect per trait unit, p, and BH q within trait and term. Table: c_data/04_associate.xlsx, model.

Change score, acute window


7 of 240 module−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


13 of 240 module−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, baseline window


13 of 240 module−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.

Change score, training window

5 of 240 module−outcome pairs at p < 0.05, 0 at BH < 0.05.



One point per subject; a line per arm. Header: Spearman rho and p pooled over both arms, and BH q within window and outcome. Corner: each arm's rho and exact p. Arm is the median split of comp_hypertrophy, so pooled correlations with it and with fibre area change partly carry the group split. Table: c_data/04_associate.xlsx, change_score and change_score_by_arm.