

a Does a feature follow a trait differently in HR and LR?
HR slope minus LR slope (limma, feature ~ group x trait). Number: % of features at p < 0.05 (white: 5, chance);
bold below: count at FDR < 0.05. The composite score defined the groups and is left out
Proteins
Set scores
Modules
fCSA mixed
fCSA type1
fCSA type2
mCSA
leg press 1RM
leg ext 1RM
volume load (kg)
T1 level
T1, gain
T2 - T1
T2 level
T3 - T2
T3 level
% at p < 0.05
20%
15%
10%
5%
0%
b The strongest screen, within each group
leg press 1RM (% change), T2 - T1, 1,422 set scores; 44 of 44 FDR hits change sign
Cardiac muscle contraction
Regulation of the force of ...
Actin myosin filament sliding
Actin mediated cell contrac...
Muscle organ morphogenesis
Cardiac muscle tissue morph...
Spearman rho in LR
Spearman rho in HR
HR - LR slope
FDR < 0.05
p < 0.05 only
not significant
c Cardiac muscle tissue morphogenesis, person by person
HR slope minus LR slope: t = -9.4, p < 0.001, FDR < 0.001
leg press 1RM (% change)
set score, T2 - T1
HR: rho -0.95 (n = 8)
LR: rho 0.95 (n = 8)
d Proteins that share a pattern of trait links
12 largest of 50 profile groups (762 of 2,059 proteins), mean limma t per screen; row: group (proteins) top over-represented set
1 (113) Spindle organization
2 (75) no set at FDR < 0.05
3 (73) no set at FDR < 0.05
4 (68) no set at FDR < 0.05
5 (62) Oxidative phosphorylation
6 (62) no set at FDR < 0.05
7 (55) Clathrin mediated endocytosis
8 (52) no set at FDR < 0.05
9 (51) Respiratory electron transport
10 (51) no set at FDR < 0.05
11 (50) Striated muscle contraction
12 (50) no set at FDR < 0.05
mean t